

Table 3. Cotton variety performance in north central Alabama, Shelby Co. 2010*.						
Producer:	Phillip Barber					
ACES Cooperators:	R. Colquitt, W. Griffith, D. Monks, K. Glass, and J. Brasher					
Variety	Lint turnout	Lint yield (lbs/acre)	Mic	Length (32nd)	Strength (g/tex)	Uniformity (%)
ST 4288B2RF	0.44	1114	5.1	33.9	25.9	82.8
DP 1048B2RF	0.47	1091	5.2	36.8	26.4	86.1
DP 1050B2RF	0.47	1090	5.0	36.2	27.3	86.2
DP 1028B2RF	0.47	1088	5.0	35.5	27.7	85.3
DP 1034B2RF	0.46	1063	5.1	35.8	27.3	86.4
DP 0935B2RF	0.45	1012	5.1	35.2	28.4	84.3
PHY 565WRF	0.44	971	5.0	35.5	31.1	85.9
DP 0949B2RF	0.47	966	5.2	35.8	29.9	84.9
PHY 375WRF	0.47	960	5.0	35.2	26.9	84.6
ST 5458B2RF	0.43	955	5.3	36.2	28.0	84.3
ST 5288B2RF	0.43	954	5.3	36.5	26.1	84.4
FM 1740B2F	0.44	934	4.8	35.8	27.3	85.7
PHY 485WRF	0.44	881	5.3	34.6	28.7	85.1
*Note: Plots were 8, 38 inch rows wide X approximately 1400 ft. in length no-till planted with 2 strips per variety. Planting date was May 7; harvest date was September 25, 2010.						
<i>Funding from Cotton Incorporated and the Alabama Cotton Commission was used in conducting this trial. We appreciate the support that Alabama cotton producers provide to our educational programs each year.</i>						