

<b>Table 2. Cotton variety performance in western central Alabama, Dallas Co. 2010*.</b>						
<b>Producer:</b>	Jay Minter					
<b>ACES Cooperators:</b> R. Yates, D. Monks, K. Glass, and J. Brasher						
<b>Variety</b>	<b>Lint turnout</b>	<b>Lint yield (lbs/acre)</b>	<b>Mic</b>	<b>Length (32nd)</b>	<b>Strength (g/tex)</b>	<b>Uniformity (%)</b>
DP 09R555B2R2	0.45	829	4.5	35.8	30.3	82.0
ST 5458B2RF	0.43	804	4.3	34.6	29.5	79.6
DP 0949B2RF	0.43	774	4.4	34.9	28.1	80.8
DG 2570B2RF	0.43	768	4.2	33.6	27.4	81.0
PHY 375WRF	0.44	761	3.9	34.6	27.2	81.5
DP 1034B2RF	0.44	750	4.0	34.9	27.3	81.9
DP 1048B2RF	0.43	745	4.3	35.5	26.1	81.0
DP 0935B2RF	0.42	730	4.1	34.2	27.5	80.2
ST 4288B2RF	0.38	718	4.0	34.9	27.4	79.1
ST 5288B2RF	0.41	702	4.1	34.6	26.4	80.5
PHY 565WRF	0.42	683	3.8	36.2	29.5	84.4
PHY 485WRF	0.42	678	4.1	33.3	27.9	81.0
DP 555BR	0.43	672	4.6	32.6	25.6	77.7
DP 1050B2RF	0.45	672	4.2	35.2	26.8	81.7
*Note: Plots were 4, 30 inch rows wide X approximately 1200 ft. in length no-till planted as single strips. Planting date was April 20; harvest date was November 5, 2010.						
<i>Funding from Cotton Incorporated and the Alabama Cotton Commission was used in conducting this trial. We appreciate the support that Alabama cotton producers provide to our educational programs each year.</i>						