



# Homeowner Trees and Ornamentals

Insect Control Recommendations for 2011

## INSECT CONTROL

---

### Homeowner Trees and Ornamentals: Insect Control

---

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>ARBORVITAE CEDAR</b>			
<b>Bagworms</b>			
<i>General Comments: See ANR-191, "<a href="#">Controlling Bagworms on Ornamentals</a>." Homeowners can handpick bags. Insecticides can be used to control bagworms if applied early in the spring (May) or summer (June) while caterpillars are small.</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	<i>Bacillus thuringiensis</i> AMERICAN BRAND THURICIDE CONCENTRATE	---	Use according to label directions.
	SAFER® CATERPILLAR KILLER	1-4 t.	
	bifenthrin ORTHO BUG B GON MAX LAWN AND GARDEN INSECT KILLER CONCENTRATE	---	
	carbaryl FERTILOME LIQUID CARBARYL GARDEN Spray	4 t.	
	HI-YIELD 10% CARBARYL GARDEN AND PET DUST	See label.	
	HI-YIELD 5% CARBARYL GARDEN AND PET DUST		
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	Concentrate, 1 T.	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.

---

**NOTE:** Before using any insecticide on a species **NOT** listed on the label, spray a small inconspicuous portion of the plant's foliage, wait 48 hours, and see if any signs of spray damage appear. Apply spray only if no signs of spray damage appear. Rates apply only to formulations listed in the table. If other formulations are used, apply at rates specified on the container.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>ARBORVITAE CEDAR (cont.)</b>			
<b>Bagworms (cont.)</b>			
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO- USE	2 T.	
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	Apply when bagworm larvae begin to hatch.
	lambda-cyhalothrin ENFORCER FLEA SPRAY FOR YARDS CONCENTRATE VI	0.5 T.	Apply when bagworm larvae begin to hatch.
	malathion HI-YIELD 5% MALATHION DUST	---	Treat in spring or early summer when bagworms are small. Repeat in 10 to 14 days if necessary.
	HI-YIELD 55% MALATHION SPRAY	---	
	SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE	See label.	
	ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	---	
	neem BIONEEM	6 T.	
	permethrin HI-YIELD KILL-A-BUG II	2 T.	
	HI-YIELD KILL-A-BUG II RTU	---	
	HI-YIELD PERMETHRIN CONCENTRATE LAWN, GARDEN, PET AND LIVE- STOCK INSECT SPRAY	1.5 t.	
	SPECTRICIDE ANT SHIELD HOME BARRIER INSECT KILLER	See label.	
	pyrethrins + potassium salts of fatty acids CONCERN MULTI-PURPOSE INSECT KILLER CONCENTRATE	See label.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>ARBORVITAE CEDAR (cont.)</b>			
<b>Bagworms (cont.)</b>			
	spinosad FERTI-LOME BORER, BAGWORM, LEAFMINER AND TENT CATERPILLAR SPRAY	4 T.	
<b>BOXELDER</b>			
<b>Boxelder Bugs</b>			
<i>Boxelder bugs are more of a nuisance when insects congregate on buildings and indoors in the fall than they are a pest of plants. Bugs can be vacuumed when they enter buildings. See the North Carolina Cooperative Extension Service Web publication "<a href="#">Boxelder Bug</a>" for more information on the bug's biology and control. The following insecticide recommendations are for outdoor use.</i>			
	carbaryl HI-YIELD 10% CARBARYL GARDEN AND PET DUST	---	
	HI-YIELD 5% CARBARYL GARDEN AND PET DUST	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	Concentrate, 1 T.	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray cyfluthrin into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has completely dried.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	See label.	
	lambda-cyhalothrin BONIDE BEETLE KILLER	0.5 T.	
	malathion ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	1 T.	
	permethrin HI-YIELD PERMETHIN CONCENTRATE LAWN, GARDEN, PET, AND LIVESTOCK INSECT SPRAY	1.5 t.	
	HI-YIELD KILL-A-BUG II	2 T.	
	HI-YIELD KILL-A-BUG II RTU SPECTRACIDE ANT SHIELD HOME BARRIER INSECT KILLER	See label.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>BOXELDER (cont.)</b>			
<b>Boxelder Bugs (cont.)</b>			
	pyrethrins + potassium salts of fatty acids SAFER BRAND BUG PATROL LAWN AND LANDSCAPE INSECTICIDE CONCENTRATE	See label.	
	pyrethrins + piperonyl butoxide RAID HOUSE AND GARDEN BUG KILLER	See label.	
	rotenone + pyrethrin BONIDE LIQUID ROTENONE PYRETHRINS SPRAY CONCENTRATE	2 t.	
	s-bioallethrin + deltamethrin BONIDE SPIDER KILLER	---	
<b>DOGWOOD</b>			
<i>See North Carolina State University Web publication <a href="#">"Insects and Related Pests of Shrub,"</a> for more information on the biology and control of several dogwood insect pests.</i>			
<b>Dogwood Borers</b>			
	permethrin HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCL. TURF, GARDENS, AND ORN.	1.5-3 fl.oz.	
<b>Dogwood Club Gall</b>			
	See Comments.	---	No pesticides are recommended. Pruning and removal of galls may be helpful.
<b>Oystershell Scale</b>			
	acetamiprid ACETAMIPRID RTU INSECTICIDE ORTHO FLOWER, FRUIT, AND VEGETABLE INSECT KILLER CONCENTRATE	---	
	carbaryl LIQUID CARBARYL GARDEN SPRAY	---	For use on crawlers only.
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	Concentrate, 1 T.	For use on crawlers only. Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray cyfluthrin into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>DOGWOOD (cont.)</b>			
<b>Oystershell Scale (cont.)</b>			
	horticultural oil DORMANT SPRAY AND SUMMER OIL SPRAY ORTHO VOLCK OIL SPRAY	3 T.  Dormant Season: 5.3 fl.oz. Growing Season: 2.5 fl.oz.	Apply oil as a dormant spray and/or when crawlers are active. Dormant Spray and Summer Oil Spray may be mixed with other insecticides and with fungicides other than those containing sulfur. It may cause plant injury in house plants such as ivy, ferns, and the like. <b>DO NOT</b> spray plants in mid-day with this spray if temperature is above 85°F. Spray in late afternoon or early morning. <b>DO NOT</b> spray when temperatures are below 40°F.
	lime sulfur HI-YIELD IMPROVED LIME SULFUR SPRAY	4 fl.oz. + 1.25 fl.oz. dormant oil	For use as a dormant spray on deciduous trees and shrubs only.
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	
<b>ELM</b>			
<b>Elm Bark Beetles</b>			
	permethrin HI-YIELD PERMETHIN CONCENTRATE LAWN, GARDEN, PET, AND LIVESTOCK INSECT SPRAY	1.5-3 fl.oz.	Keep trees growing vigorously to prevent infestations. Remove broken branches and weak trees, which may serve as breeding sites. Apply in early May and again in early June.
	See Comments.		Pesticides for this pest are for use by professionals only. Keep trees growing vigorously to prevent infestations. Re-move broken branches and weak trees which may serve as breeding sites.
<b>Elm Leaf Beetles</b>			
<i>See the Ohio State University Extension Web publication "<a href="#">Elm Leaf Beetle</a>" for more information on the biology and control of the elm leaf beetle.</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>ELM (cont.)</b>			
<b>Elm Leaf Beetles (cont.)</b>			
	carbaryl ELIMINATOR 5% SEVIN DUST ELIMINATOR 10% SEVIN DUST	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	Concentrate, 1 T.	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray the product into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until the spray has dried.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	deltamethrin BUGGSLAYER INSECTICIDE	3 T.	
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	8 T. per foot of plant height	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL BONIDE SYSTEMIC GRANULES INSECT CONTROL	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	lambda-cyhalothrin BONIDE BEETLE KILLER	0.5 T.	
	permethrin BONIDE BUG BEATER II YARD AND GARDEN CONTROL CONCENTRATE	1 fl.oz.	
	pyrethrins + piperonyl butoxide SPECTRACIDE BUG STOP FOR GARDENS CONCENTRATE	4 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>MAGNOLIA</b>			
<b>Magnolia Scale</b>			
<i>For more information on the biology and control of magnolia scale, see Ohio State University Extension Web publication "<a href="#">Magnolia Scale and Its Control.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	Concentrate, 1 T.	Use for crawlers only. Apply to flowering plants in early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until the spray has dried.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	horticultural oil FERTI-LOME DORMANT SPRAY AND SUMMER OIL SPRAY	3 T.	
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	Treat when crawlers are active, normally in August and September.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>PINE</b>			
<b>Aphids</b>			
<i>For more information on controlling aphids, see Alabama Cooperative Extension System publication ANR-189, "<a href="#">Controlling Aphids on Ornamentals.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	2 T.	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	Concentrate, 1 T.	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until the spray has dried.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	horticultural oil FERTI-LOME DORMANT SPRAY AND SUMMER OIL SPRAY	3 T.	
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	
	neem BIONEEM	6 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>PINE (cont.)</b>			
<b>Black Turpentine Beetles</b>			
	permethrin HI-YIELD PERMETHIN CONCENTRATE LAWN, GARDEN, PET, AND LIVESTOCK INSECT SPRAY	1.5 t.	
	HI-YIELD KILL-A-BUG II	2 T.	
	HI-YIELD KILL-A-BUG II RTU		
	SPECTRACIDE ANT SHIELD	See label.	
	HOME BARRIER INSECT KILLER		
	pyrethrins EVERGREEN PYRETHRUM DUST	---	
<b>Ips Engraver Beetles And Southern Pine Beetles</b>			
<i>For more information, see Alabama Cooperative Extension System publication <a href="#">"Management of Pine Bark Beetles Attacking Landscape Pines."</a></i>			
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	permethrin HI-YIELD PERMETHIN CONCENTRATE LAWN, GARDEN, PET, AND LIVESTOCK INSECT SPRAY	1.5 t.	Trees should be kept vigorous and free of injury. Spray trees adjacent to infested trees to prevent attack. Rapid tree removal is the only control once beetles have attacked. Spraying should be done by commercial applicators. In forest stands, promptly salvage all merchantable infested trees. Cut remaining trees and destroy the infestation in these and all remaining slash by burning.
	HI-YIELD KILL-A-BUG II	2 T.	
	HI-YIELD KILL-A-BUG II RTU		
	SPECTRACIDE ANT SHIELD	See label.	
	HOME BARRIER INSECT KILLER		
	pyrethrins EVERGREEN PYRETHRUM DUST	---	
<b>Sawflies</b>			
<i>For more information on the biology and control of pine sawflies, see University of Kentucky Web publication <a href="#">"Pine Sawflies."</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	Treat when larvae are first detected.
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	azadirachtin SAFER® BRAND BIONEEM® INSECTICIDE AND REPELLENT MULTI-PURPOSE CONCENTRATE	3 fl.oz.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>PINE (cont.)</b>			
<b>Sawflies (cont.)</b>			
	carbaryl ELIMINATOR BUG KILLER CONCENTRATE SEVIN BRAND CARBARYL INSECTICIDE	3 T.	
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO- USE	2 T.	
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	lambda-cyhalothrin BONIDE CATERPILLAR KILLER	0.5 T.	Apply when bagworm larvae begin to hatch.
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	
	permethrin HI-YIELD 38 PLUS TURF, TERMITE AND ORN. INSECT CTRL. CONCENTRATE 38% PERMETHRIN	0.125-0.25 t.	
	thiamethoxam + lambda- cyhalothrin MAXIDE DUAL ACTION INSECT KILLER CONCENTRATE	3 fl.oz.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>PINE (cont.)</b>			
<b>Scale</b>			
<i>For more information on the biology and control of scale insects on ornamental plants, see Texas Agricultural Extension Service Web publication <a href="#">"Scale Insects on Ornamental Plants."</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	For treatment of crawlers, only.
	carbaryl LIQUID CARBARYL GARDEN SPRAY	4 t.	For treatment of crawlers, only.
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	Concentrate, 1 T.	For treatment of crawlers, only. Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	
	horticultural oil DORMANT SPRAY AND SUMMER OIL SPRAY ORTHO VOLCK OIL SPRAY CONCERN PESTICIDAL SPRAY OIL	3 T.  Dormant Season: 5.3 fl.oz. Growing Season: 2.5 fl.oz. 5 T.	Apply oil as a dormant spray and/or when crawlers are active. Dormant Spray and Summer Oil Spray may be mixed with other insecticides and with fungicides except those containing sulphur. It may cause plant injury in house plants such as ivy, ferns, and the like. <b>DO NOT</b> spray plants in midday with Dormant Spray if temperature is above 90°F. Spray in late afternoon or early morning. Spray when temperature is above 40°F. and there is no danger of freezing. <b>DO NOT</b> apply as a foliar spray when temperature is above 80°F.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>PINE (cont.)</b>			
<b>Scale (cont.)</b>			
	malathion HI-YIELD 5% MALATHION DUST	See label.	
	HI-YIELD 55% MALATHION SPRAY		
	SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE		
	ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE		
	permethrin ORTHO BUG B GON GARDEN AND LANDSCAPE READY SPRAY INSECT KILLER	See label.	
	pyrethrins and soap YARD AND GARDEN INSECT KILLER	10 T.	
<b>Tip Moths (Including Nantucket Pine Tip Moths)</b>			
<i>For more information on the biology and control of pine tip moths, see the USDA Forest Disease and Insect Web publication "<a href="#">Nantucket Pine Tip Moth.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL CONCENTRATE FERTILOME TREE AND SHRUB SYSTEMIC INSECT GRANULES	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
<b>SYCAMORE</b>			
<b>Sycamore Lace Bugs</b>			
<i>For more information on the biology and control of lace bugs on ornamentals, see Alabama Cooperative Extension System Web publication ANR-191, "<a href="#">Controlling Lace Bugs on Ornamentals.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	capsaicin ANIMAL REPELLENT CONTAINING NATURAL HOT PEPPER WAX	See label.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>SYCAMORE (cont.)</b>			
<b>Sycamore Lace Bugs (cont.)</b>			
	carbaryl EASY GARDEN LIQUID CONCENTRATE	4 t.	When insects are noticed, treat by covering foliage well with spray.
	GARDEN TECH SEVIN CONCENTRATE BUG KILLER	3 T.	
	GARDEN TECH SEVIN READY-TO-USE	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	Concentrate, 1 T.	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE HI-YIELD SYSTEMIC INSECT GRANULES	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	Treat foliage thoroughly, especially the undersides of leaves, when insects are detected. Repeat if needed.
	permethrin ELIMINATOR HOME PEST INSECT CONTROL	See label.	
	pyrethrins EVERGREEN PYRETHRUM DUST	---	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS</b>			
<b>Aphids</b>			
<i>See Alabama Cooperative Extension System publication ANR-189, "<a href="#">Controlling Aphids on Ornamentals.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	2 T.	Treat infested trees and shrubs in the fall to reduce the number producing over-wintering eggs. Begin treatment when aphids appear in the spring.
	acetamiprid ACETAMIPRID RTU INSECTICIDE	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	See label for rates and directions for use.	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO-USE	2 T.	
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and directions for use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	insecticidal soap SAFER® BRAND INSECT KILLING SOAP	5 T.	Examine plants after two days and spray again if needed. For other pests, spray weekly as needed up to three times.
	SAFER® BRAND ROSE AND FLOWER INSECT KILLER	---	<b>DO NOT</b> make repeated sprays at short intervals without thoroughly rinsing foliage.
	insecticidal soap + pyrethrins SAFER® BRAND YARD AND GARDEN INSECT KILLER	---	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Aphids (cont.)</b>			
	malathion HI-YIELD 5% MALATHION DUST	See label.	Treat foliage thoroughly, especially the undersides of leaves, when insects are detected. Repeat if needed.
	HI-YIELD 55% MALATHION SPRAY		
	SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE		
	ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE		
	neem BIONEEM	6 T.	
	insecticidal oil FERTILOME SCALECIDE	5-7.6 T.	Fertilome Scalecide is a dormant spray oil. Spray when the temperature is above 40°F. and there is no danger of freezing. <b>DO NOT</b> apply as a foliar spray when temperature is above 80°F.
	pyrethrum GORDON GARDEN GUARD	See label.	
<b>Ambrosia Beetles</b>			
	See Comments.		Cut and destroy infested plants. No chemical control is recommended.
<b>Azalea Caterpillars</b>			
<i>Caterpillars may be removed by hand. Treat when larvae are small. For more information about the biology and control of the azalea caterpillar, see North Carolina State University Web publication "<a href="#">Azalea Caterpillar.</a>"</i>			
	acetamiprid ACETAMIPRID RTU INSECTICIDE	---	
	<i>Bacillus thuringiensis</i> AMERICAN BRAND THURI- CIDE CONCENTRATE	4 t.	Use according to label directions.
	SAFER® BRAND CATER- PILLAR KILLER	1-4 t.	
	FERTILOME DIPEL DUST	---	
	bifenthrin ORTHO BUG B GON MAX LAWN AND GARDEN INSECT KILLER CONCENTRATE	---	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Azalea Caterpillars (cont.)</b>			
	carbaryl		
	EASY GARDEN LIQUID CONCENTRATE	4 t.	When insects are noticed, treat by covering foliage well with spray.
	FERTILOME LIQUID CAR- BARYL GARDEN SPRAY	4 t.	
	GARDEN TECH SEVIN READY-TO-USE		
	GARDEN TECH SEVIN CONCENTRATE BUG KILLER	3 T.	
	HI-YIELD 10% CARBARYL GARDEN AND PET DUST	See label.	
	HI-YIELD 5% CARBARYL GARDEN AND PET DUST	---	
	cyfluthrin		
	BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	Concentrate, 1 T.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid		
	BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	esfenvalerate		
	ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO- USE	2 T.	
	gamma-cyhalothrin		
	SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	
	neem		
	BIONEEM	6 T.	
<b>Bagworms</b>			
<i>Insecticides may be used to control bagworms if applied early in the spring (May) or summer (June) while caterpillars are small. Also, homeowners may want to handpick the bags. See the Alabama Cooperative Extension System Web publication ANR-191, "<a href="#">Controlling Bagworms on Ornamentals.</a>"</i>			
	acephate		
	BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide		
	ORTHO SYSTEMIC INSECT KILLER	3 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Bagworm (cont.)</b>			
	<i>Bacillus thuringiensis</i>		Use according to label directions.
	AMERICAN BRAND THURICIDE CONCENTRATE	4 t.	
	SAFER® BRAND CATERPILLAR KILLER	1-4 t.	
	bifenthrin		
	ORTHO BUG B GON MAX LAWN AND GARDEN INSECT KILLER CONCENTRATE	---	
	carbaryl		
	FERTILOME LIQUID CARBARYL GARDEN SPRAY	4 t.	
	HI-YIELD 5% CARBARYL GARDEN AND PET DUST	---	
	HI-YIELD 10% CARBARYL GARDEN AND PET DUST	See label.	
	cyfluthrin		
	BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	Concentrate, 1 T.	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	cyfluthrin + imidacloprid		
	BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply to flowering plants during early morning or late evening when bees are not present. <b>DO NOT</b> spray into fish pools, ponds, streams, or lakes. Remove animal feeding dishes before applying. Keep children and pets away from treated areas until spray has dried.
	gamma-cyhalothrin		
	SPECTRACIDE TRIAZIDE ONCE AND DONE! INSECT KILLER CONCENTRATE	0.5 T.	For bagworm control, apply when bagworm larvae begin to hatch.
	malathion		
	HI-YIELD 5% MALATHION DUST	See label.	Treat in spring or early summer when bagworms are small. Repeat in 10 to 14 days if necessary.
	HI-YIELD 55% MALATHION SPRAY		
	SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE		
	ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE		

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Bagworms (cont.)</b>			
	neem BIONEEM	6 T.	
	permethrin HI-YIELD KILL-A-BUG II HI-YIELD KILL-A-BUG II RTU	2 T.	
	HI-YIELD PERMETHRIN CONCENTRATE LAWN, GARDEN, PET AND LIVE- STOCK INSECT SPRAY	1.5 t.	
	SPECTRACIDE ANT SHIELD HOME BARRIER INSECT KILLER	See label.	
	pyrethrins CONCERN MULTI-PURPOSE INSECT KILLER	See label.	
	spinosad FERTI-LOME BORER, BAG- WORM, LEAFMINER AND TENT CATERPILLAR SPRAY	4 T.	
	CONSERVE NATURALYTE INSECT CONTROL	4 T.	
<b>Black Twig Borers</b>			
	See Comments.		Prune and burn infested twigs. No chemical control is recommended.
<b>Cutworms</b>			
<i>See Web-based information in the Alternative Control Outreach Research Network (ACORN) for pesticide alternatives for <a href="#">cutworm control</a>.</i>			
	<i>Bacillus thuringiensis,</i> var. kurstaki DIPEL DUST	0.5-1 oz./ 50 sq.ft.	
	carbaryl CUTWORM AND CRICKET BAIT	---	For turf applications, use one pound per 1,000 square feet.
	FERTILOME LIQUID CARBARYL GARDEN SPRAY	4 t.	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	Concentrate, 1 T.	Apply in early morning or late evening when bees are not present.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Cutworms (cont.)</b>			
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO- USE	2 T.	
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	
	neem BIONEEM	6 T.	
	permethrin HI-YIELD KILL-A-BUG II HI-YIELD KILL-A-BUG II RTU HI-YIELD PERMETHRIN CONCENTRATE LAWN, GARDEN, PET AND LIVE- STOCK INSECT SPRAY SPECTRACIDE ANT SHIELD HOME BARRIER INSECT KILLER	2 T.  1.5 t.  See label.	
<b>Eastern Tent Caterpillars</b>			
<i>Eastern tent caterpillars are found mostly on flowering fruit trees. Their webs are constructed in forks and crotches of trees in spring. Treat when small webs are noticed. Treat late in the day when larvae are in webs. Direct spray into webs and thoroughly saturate. As an alternative to spraying, webs and larvae can be removed and destroyed by hand late in the day. For more information on the biology and control of the eastern tent caterpillar, see University of Kentucky Web publication "<a href="#">Eastern Tent Caterpillar.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	<i>Bacillus thuringiensis,</i> var. kurstaki DIPEL DUST SAFER® BRAND CATER- PILLAR KILLER AMERICAN BRAND THURICIDE CONCENTRATE	1-2 t.  4 t.	Thoroughly dust plant surfaces.
	carbaryl CARBARYL 10% DUST FERTILOME LIQUID CARBARYL GARDEN SPRAY	4 t.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Eastern Tent Caterpillars (cont.)</b>			
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO- USE	---	
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	
	neem BIONEEM	6 T.	
	permethrin HI-YIELD KILL-A-BUG II HI-YIELD KILL-A-BUG II RTU	2 T.	
	HI-YIELD PERMETHRIN CONCENTRATE LAWN, GARDEN, PET AND LIVE- STOCK INSECT SPRAY	1.5 t.	
	SPECTRACIDE ANT SHIELD HOME BARRIER INSECT KILLER	See label.	
	spinosad FERTI-LOME BORER, BAG- WORM, LEAFMINER AND TENT CATERPILLAR SPRAY	4 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Fall Webworms</b>			
<i>Fall webworms attack many deciduous trees and shrubs. Their webs enclose the foliage on ends of branches and are seen in summer and fall. Treat when webs are noticed and saturate thoroughly. As an alternative to spraying, webs and larvae can be removed and destroyed by hand late in the day. For more information on the biology and control of the fall webworm, see Ohio State University Extension Fact Sheet <a href="#">"Fall Webworm Management."</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	<i>Bacillus thuringiensis</i> , var. kurstaki DIPEL DUST		Thoroughly dust plant surfaces.
	SAFER® BRAND CATERPILLAR KILLER	1-2 t.	
	AMERICAN BRAND THURICIDE CONCENTRATE	4 t.	
	carbaryl FERTILOME LIQUID CARBARYL GARDEN SPRAY	4 t.	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO-USE	2 T.	
	gamma-cyhalothrin SPECTRACIDE TRIAZICIDE ONCE AND DONE! INSECT KILLER	0.5 T.	
	neem BIONEEM	6 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Fall Webworms (cont.)</b>			
	permethrin		
	HI-YIELD KILL-A-BUG II	2 T.	
	HI-YIELD KILL-A-BUG II RTU		
	HI-YIELD PERMETHRIN	1.5 t.	
	CONCENTRATE LAWN, GARDEN, PET AND LIVE-STOCK INSECT SPRAY		
	SPECTRACIDE ANT SHIELD	See label.	
	HOME BARRIER INSECT KILLER		
	spinosad		
	FERTI-LOME BORER, BAG-WORM, LEAFMINER AND TENT CATERPILLAR SPRAY	4 T.	
<b>Gall Insects</b>			
	See Comments.		Galls generally do not occur in large numbers. Pruning and removal may be helpful. No insecticide is recommended.
<b>Holly Leafminers</b>			
<i>For more information, see University of Tennessee Extension publication "<a href="#">Holly Leafminers.</a>"</i>			
	acetamiprid		
	ACETAMIPRID RTU INSECTICIDE	---	
	cyfluthrin + imidacloprid		
	BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	disulfoton		
	2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	FERTI-LOME SYSTEMIC INSECTICIDE GRANULES		
	imidacloprid		
	BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	neem		
	BIONEEM	6 T.	
	spinosad		
	FERTI-LOME BORER, BAG-WORM, LEAFMINER AND CATERPILLAR SPRAY	4 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Japanese Beetles</b>			
<i>Japanese beetles are not yet found statewide in Alabama but are a problem where they are found. Japanese beetles feed on more than 400 species of plants and can be a pest of turf. For more information on Japanese beetles see <a href="#">"Japanese Beetles in Alabama."</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	carbaryl FERTILOME LIQUID CARBARYL GARDEN SPRAY	4 t.	
	HI-YIELD 10% CARBARYL GARDEN AND PET DUST	See label.	
	HI-YIELD 5% CARBARYL GARDEN AND PET DUST		
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	malathion HI-YIELD 5% MALATHION DUST	See label.	
	HI-YIELD 55% MALATHION SPRAY		
	ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE		
	neem BIONEEM	6 T.	Products containing neem are repellant to Japanese beetles.
	pyrethrins JAPANESE BEETLE KILLER	---	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Lace Bugs</b>			
<i>Treat when nymphs appear in the spring. Repeat if needed. Be sure to direct spray to the undersides of leaves. For more information on the biology and control of this pest, see Alabama Cooperative Extension System Web publication <a href="#">"Controlling Lace Bugs on Ornamentals."</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	carbaryl HI-YIELD 5% CARBARYL GARDEN AND PET DUST HI-YIELD 10% CARBARYL GARDEN AND PET DUST	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND READY-TO-USE GRANULES	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	insecticidal soap CONCERN INSECT KILLING SOAP	See label.	
	insecticidal soap and pyrethrins YARD AND GARDEN INSECT KILLER	10 T.	
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Lace Bugs (cont.)</b>			
	neem BIONEEM	6 T.	
	rotenone-pyrethrin BONIDE ROTENONE-PYRETHRIN SPRAY	4 t.	
<b>Leafhoppers</b>			
<i>For more information on controlling leafhoppers in the garden and landscape, see Utah State University Extension publication <a href="#">"Leafhoppers in the Home Garden."</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acetamiprid ACETAMIPRID RTU INSECTICIDE	---	
	capsaicin, related compounds BONIDE HOT PEPPER WAX CONCENTRATE	8-16 T.	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES BONIDE SYSTEMIC GRANULES	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	esfenvalerate ORTHO BUG B GON MAX GARDEN AND LANDSCAPE INSECT KILLER READY-TO-USE	4 T.	
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	neem BIONEEM	6 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Leafhoppers (cont.)</b>			
	permethrin		
	HI-YIELD KILL-A-BUG II	2 T.	
	HI-YIELD KILL-A-BUG II RTU		
	HI-YIELD PERMETHRIN	1.5 t.	
	CONCENTRATE LAWN, GARDEN, PET AND LIVE-STOCK INSECT SPRAY		
	SPECTRACIDE ANT SHIELD	See label.	
	HOME BARRIER INSECT KILLER		
	pyrethrins and soap		
	YARD AND GARDEN INSECT KILLER	10 T.	
	rotenone and pyrethrins		
	ROTENONE-PYRETHRINS SPRAY	1 t.	
<b>Leafminers</b>			
<i>For more information on controlling the boxwood leafminer, see North Carolina Extension Service publication "<a href="#">Boxwood Leafminer.</a>"</i>			
	acephate		
	BONIDE SYSTEMIC INSECT CONTROL	3 T.	Make treatments in spring when mines appear. Boxwood leafminer treatments should be applied in late March. A spreader-sticker may be used for plants with slick, waxy leaves.
	acephate + fenbutatin oxide		
	ORTHO SYSTEMIC INSECT KILLER	3 T.	
	acetamiprid		
	ACETAMIPRID RTU INSECTICIDE	---	
	carbaryl		
	LIQUID CARBARYL GARDEN SPRAY	4 t.	
	cyfluthrin + imidacloprid		
	BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	imidacloprid		
	BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	malathion		
	HI-YIELD 55% MALATHION SPRAY	2 t.	
	neem		
	BIONEEM	6 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Leafrollers</b>			
<i>For more information on the biology and control of the canna leafroller, see the University of Florida Web publications "<a href="#">Larger Canna Leafroller</a>" and "<a href="#">Lesser Canna Leafroller</a>."</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	<i>Bacillus thuringiensis,</i> var. kurstaki SAFER® BRAND CATER- PILLAR KILLER	1-2 t.	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	neem BIONEEM	6 T.	
	permethrin HI-YIELD KILL-A-BUG II HI-YIELD KILL-A-BUG II RTU HI-YIELD PERMETHRIN CONCENTRATE LAWN, GARDEN, PET AND LIVE- STOCK INSECT SPRAY SPECTRACIDE ANT SHIELD HOME BARRIER INSECT KILLER	2 T. 1.5 t. See label.	
	rotenone and pyrethrins ROTENONE-PYRETHRINS SPRAY	1 t.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Leafrollers (cont.)</b>			
	spinosad FERTI-LOME BORER, BAG- WORM, LEAFMINER AND TENT CATERPILLAR SPRAY	4 T.	
<b>Mealybugs</b>			
<i>For more information on the biology and control of mealybugs, see North Carolina State University Cooperative Extension's Web publication "<a href="#">Mealybugs.</a>"</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	acetamiprid ACETAMIPRID RTU INSECTICIDE	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI- INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	insecticidal oil CONCERN PESTICIDAL SPRAY OIL	5 T.	
	insecticidal soap CONCERN INSECT KILLING SOAP	5 T.	
	malathion HI-YIELD 5% MALATHION DUST HI-YIELD 55% MALATHION SPRAY SPECTRACIDE MALATHION INSECT SPRAY CONCENTRATE ORTHO MALATHION PLUS INSECT SPRAY CONCENTRATE	See label.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Mealybugs (cont.)</b>			
	neem BIONEEM	6 T.	
	rotenone and pyrethrins YARD AND GARDEN INSECT KILLER	RTU	
	permethrin PERMETROL LAWN INSECTICIDE	See label.	
<b>Mimosa Webworms</b> —See pesticide recommendations for Fall Webworms.			
<b>Nantucket Pine Tip Moth</b> —See pesticide recommendations for Pine Tip Moths in the Pines section.			
<b>Other Foliage-Feeding Caterpillars</b> —See pesticide recommendations for other caterpillars in this section, such as Fall Webworm, Leaf Rollers, or Eastern Tent Caterpillar.			
<b>Pine Bark Beetles</b> —See Ips Engraver Beetles and Southern Pine Beetles in the Pines section.			
<b>Sawflies</b>			
<i>For more information on sawflies, see the University of Minnesota’s Web publication <a href="#">“Sawflies of Trees and Shrubs.”</a></i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	imidacloprid BAYER ADVANCED GAR- DEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and directions for use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the treated plant.
	neem BIONEEM	6 T.	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Scale Insects</b> —See pesticide recommendations for scale insects in the Pines section.			
<b>Slugs and Snails</b>			
<i>For more information on biology and control of slugs and snails, see the University of California's <a href="#">Pest Management Guidelines for Slugs and Snails</a>.</i>			
	iron phosphate SLUGGO GARDEN SAFE SLUG AND SNAIL BAIT	See label.	
	metaldehyde ELIMINATOR SNAIL AND SLUG BAIT	---	
	HI-YIELD IMPROVED SLUG AND SNAIL BAIT	---	
	ORTHO'S BUG-GETA SNAIL AND SLUG KILLER	---	
	metaldehyde + carbaryl SLUG, SNAIL, SOWBUG BAIT ORTHO BUG GETA PLUS DEADLINE BUG BAIT	---	
<b>Spidermites</b>			
<i>For more information on spidermites, see the Colorado State University Web publication "<a href="#">Spider Mites</a>."</i>			
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	Inspect plants in April for presence of two-spotted spidermites. Use insecticidal soaps and/or oils early in the season in order to preserve beneficials. See label for restrictions for using oils on evergreens and for temperature requirements. Spruce spidermites are active on evergreens in the spring and, sometimes, in the fall. Shake foliage over white paper to verify presence of spidermites. Alternate miticides once or twice during the season if mites continue to be a problem. Spray twice at 10-day intervals or as label specifies.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES BONIDE SYSTEMIC GRANULES 2%	---	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Spidermites (cont.)</b>			
	oil		
	CONCERN PESTICIDAL SPRAY OIL	5 T.	
	insecticidal soap		
	CONCERN INSECT KILLING SOAP	5 T.	
<b>Spittlebugs</b> —See pesticide recommendations for Leafhoppers.			
<i>For information on the two-lined spittlebug, see North Carolina State University's Web publication <a href="#">"Managing the Two-Lined Spittlebug in the Home Landscape."</a></i>			
<b>Thrips</b>			
<i>For more information on controlling thrips in the landscape, see the university of Florida's Web publication <a href="#">"Thrips on Ornamental Plants."</a></i>			
	acephate		Destroy old blossoms. Treat weekly during the growing season.
	BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide		
	ORTHO SYSTEMIC INSECT KILLER	3 T.	
	acetamiprid		
	ACETAMIPRID RTU INSECTICIDE	---	
	capsaicin and related compounds		
	BONIDE HOT PEPPER WAX CONCENTRATE	8-16 T.	
	cyfluthrin		Apply in early morning or late evening when bees are not present.
	BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	See label.	
	cyfluthrin + imidacloprid		Apply in early morning or late evening when bees are not present.
	BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	
	disulfoton		<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	---	

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Thrips (cont.)</b>			
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	neem BIONEEM	6 T.	
	spinosad FERTI-LOME BORER, BAGWORM, LEAFMINER AND TENT CATERPILLAR SPRAY	4 T.	
<b>Whiteflies</b>			
<i>For more information on whitefly biology and control, see the University of California <a href="#">Pest Management Guidelines for Whiteflies</a>.</i>			
	acephate BONIDE SYSTEMIC INSECT CONTROL	3 T.	
	acephate + fenbutatin oxide ORTHO SYSTEMIC INSECT KILLER	3 T.	
	acetamiprid ACETAMIPRID RTU INSECTICIDE	---	
	cyfluthrin BAYER ADVANCED POWERFORCE® MULTI-INSECT KILLER	See label.	Apply in early morning or late evening when bees are not present.
	cyfluthrin + imidacloprid BAYER ADVANCED GARDEN ROSE AND FLOWER INSECT KILLER RTS	---	Apply in early morning or late evening when bees are not present.
	disulfoton 2-IN-1 SYSTEMIC ROSE AND FLOWER CARE READY-TO-USE GRANULES	See label.	<b>DO NOT</b> apply near fish pools, ponds, streams, or lakes.
	horticultural oil, such as CONCERN PESTICIDAL SPRAY OIL	5 T.	Spray as for scale control.

Insect	Insecticide and Formulation	Amount per Gallon	Comments
<b>VARIOUS TREES AND SHRUBS (cont.)</b>			
<b>Whiteflies (cont.)</b>			
	imidacloprid BAYER ADVANCED GARDEN TREE AND SHRUB INSECT CONTROL CONCENTRATE	See label for rates and use.	When applied to soil, insecticide is absorbed by plants. Once absorbed, it moves throughout the plant.
	neem BIONEEM	6 T.	
	permethrin BONIDE EIGHT INSECT CONTROL VEGETABLE, FRUIT AND FLOWER CONCENTRATE	4 T.	
	soap INSECTICIDAL SOAP	5 T.	

**Note:** See labels for use information, including phytotoxicity to certain plants.

**Note:** Some insecticides are broken down in alkaline conditions (high pH). Therefore, it is best to apply spray mixtures soon after mixing and prepare only the needed amount of spray material. For more information on this topic, see the Mississippi State University Web publication, [“Water pH Affects Fungicide and Insecticide Performance.”](#)

**Note:** Pesticide recommendations are based on the best information currently available. No endorsement of named products is intended nor is criticism implied of similar products which are not mentioned

**For disease and nematode control, see recommendations for Commercial Ornamentals.**

#### 2011 IPM-1304

**For more information**, contact your county Extension office. Visit <http://www.aces.edu/counties> or look in your telephone directory under your county's name to find contact information.

---

Use pesticides **only** according to the directions on the label. Follow all directions, precautions, and restrictions that are listed. Do not use pesticides on plants that are not listed on the label.

---

The pesticide rates in this publication are recommended **only** if they are registered with the Environmental Protection Agency or the Alabama Department of Agriculture and Industries. If a registration is changed or canceled, the rate listed here is no longer recommended. Before you apply **any** pesticide, check with your county Extension agent for the latest information.

---

Trade names are used **only** to give specific information. The Alabama Cooperative Extension System does not endorse or guarantee any product and does not recommend one product instead of another that might be similar.

---

Published by the Alabama Cooperative Extension System (Alabama A&M University and Auburn University), an equal opportunity educator and employer.

