

Controlling Rats and Mice Around Your Home

Many types of rodents may cause problems in home settings. Squirrels, flying squirrels, and chipmunks may become a nuisance from time to time, but the most troublesome and potentially harmful are rats and mice.

Norway rats, roof rats, and house mice are “commensal” rodents; that is, they live in close association with humans. Rats and mice may cause harm by transmitting disease, destroying property, and biting. While rats and mice may be an economic threat in agricultural settings, this publication focuses on controlling these rodents in and around buildings where people live.


Rats

Two species of rats are the most frequent offenders. Neither the Norway rat (*Rattus norvegicus*) nor the roof rat (*Rattus rattus*) are native to North America. They both arrived on boats from Europe along with the early settlers. While these introductions were not intentional, Norway rats and roof rats have established themselves as permanent residents.

An average adult Norway rat weighs about 1 pound. Its long, hairless tail is shorter than the head and body. Norway rats will nest under buildings and concrete slabs, along stream banks, and in garbage dumps. Often, they will burrow under the foundations of buildings and develop an elaborate burrow system with an entrance and several escape holes.

Although Norway rats can climb, they tend to inhabit the lower floors of multi-story buildings. In general, Norway rats are troublesome close to the ground, while roof rats inhabit the rafters.

Roof rats are sometimes called black rats. As their name implies, roof rats are more aerial than Norway rats. They may be found in trees, vine-covered fences, or attics. They often enter buildings from the roof or from utility lines. Roof rats differ in appearance from Norway rats in that they are sleek, graceful, and generally not as large, and they have a pointed muzzle, larger ears, and a tail that is longer than the body.




Rats are usually nocturnal; however, as population densities increase they may be seen during daylight hours. They have an excellent sense of smell, taste, and touch.


Rat populations can expand rapidly. If unchecked, one pair can produce 6 to 12 young in 21 days. Sexual maturity is reached at 3 months. Therefore, assuming adequate food, water, and shelter, a single pair of rats may multiply into more than 640 rats in 1 year.

Mice

Like Norway rats and roof rats, the house mouse (*Mus musculus*) is not native to North America; it, too, arrived with the European settlers. House mice usually weigh less than 1 ounce and are grayish-brown in color. They are masters at gaining access to buildings and will chew, gnaw, climb, or swim to reach their destination.




Young Rat



House Mouse


Mus musculus

	Feet	Head
Young Rat	Large	Large
House Mouse <i>Mus musculus</i>	Small	Small



Roof Rat

Rattus rattus



Norway Rat

Rattus norvegicus

	Tail	Body	Ear	Eye	Nose
Roof Rat <i>Rattus rattus</i>	Longer than head + body.	Light and slender	Large	Large	Pointed
Norway Rat <i>Rattus norvegicus</i>	Shorter than head + body.	Heavy and thick	Small	Small	Blunt

Characteristics of commensal rodents.

House mice are primarily nocturnal. However, unlike Norway rats and roof rats, seeing mice during daylight hours does not necessarily indicate a high population.

House mice will eat just about anything, but they prefer grains and seeds. However, they eagerly consume foods high in fat, protein, or sugar. One characteristic that distinguishes them from Norway rats and roof rats is their ability to survive with little or no free water. They get the water they need from the foods they eat.

House mice shred paper, cloth, and other materials to make a ball-shaped nest. A litter of five to six young are born about 20 days after mating. These young are weaned in 3 weeks, and they are sexually mature at 6 to 10 weeks. A female may have five to ten litters per year. Thus in one year, a single pair of mice may multiply into several thousand.

Rat And Mouse Control

Biological Control

Many types of wild and domestic animals may prey on rats and mice. Hawks, owls, and snakes will help keep rodent populations in check out-of-doors. While it is true that house cats may help reduce rat and mouse populations around the home, free-ranging house cats can have tremendous negative impacts on native wildlife such as songbirds. Biological control measures vary greatly in their effectiveness and are not reliable as a primary technique.

Rodent Proofing

Effective rodent control involves exclusion, sanitation, and chemical control. Wherever possible, buildings must be made rodent-proof. Rats and mice can climb wires, cables, vines, and trees to enter a house. They can climb the exterior siding of structures made of rough wood, concrete, or rough sheet metal. Rats may crawl along pipes up to 3 inches in diameter. The front teeth of rodents are not designed for gnawing hard, flat surfaces. However, if the surface is rough or has an edge to it, they can gnaw through most materials.

Rats have been known to gain entry into a structure through holes as small as 1/2 inch. Mice can enter through openings of 1/4 inch. Blocking possible entrance holes with secured sheet metal can help control rodent access to a building. Sheet metal cones or guards on pipes and cables may help prevent rats and mice from crawling or climbing to gain entry. Rat and mouse control is best achieved by combining exclusion techniques with a strong program of sanitation and poisoning.

Sanitation

Cleaning up around buildings can help control rat and mouse populations. Poor sanitation is one of the basic reasons for high populations of commensal rodents in urban and suburban areas. Pick up all garbage and litter, and store it in a rodent-proof container. In addition to removing possible sources of food, remove piles of trash or debris that could serve as cover.

Rodenticides

If proper exclusion techniques and a sound sanitation program are used, the need for rodenticides will be reduced. However, rodenticides may be useful in bringing a rodent problem under control or as a maintenance program to prevent rodent populations from becoming established.

Before beginning a poisoning program, estimate the number of rats or mice present. The following criteria are useful in estimating the number of rats in an area.

No rats or a few present: No sign. If rats are present, there are only a few, and they may have only recently invaded.

Medium numbers present: Old droppings and gnawing common; one or more rats seen by flashlight at night; no rats seen in daytime.

Large numbers present: Fresh droppings, tracks, and gnawing. Three or more rats seen at night and rats seen in daytime.

Mice do not wander as far from the nest as rats. Thus a relative measure of mouse density may be obtained by sprinkling talc or flour at 20 to 30 foot intervals. The number of mouse tracks observed in these patches is a good indicator of how widespread the mouse problem may be.

There are two broad categories of rodenticides available, single-dose and multiple-dose. The single-dose, or acute, rodenticides are preferred where rat or mouse numbers are large and a quick reduction in population is desired. They also are useful where it may be difficult to get rats or mice to consume bait several days in succession.

Single-dose rodenticides are usually more effective with prebaiting. Rats and mice will get "bait shy" if they eat poison and get sick but do not die. Prebaiting, training rats or mice to feed on non-toxic baits at a bait station, before putting out poisons will allow the rats to get used to the new food source without associating it with sickness or death.

Multiple-dose, or chronic, rodenticides are mostly anticoagulants with a cumulative toxic effect. Death occurs about 4 or 5 days after the ro-

dent first eats the bait. These do not produce bait-shyness and are among the safest rodenticides. Because they act slowly, anticoagulant baits may serve as their own prebait. Anticoagulants kill by inhibiting the clotting ability of the blood, so the animal dies of internal bleeding. Mice are generally less susceptible to anticoagulant baits than are rats.

Many different types and preparations of rodenticides are effective. Your specific situation dictates which one is best to use.

Rats and mice have poor eyesight and are essentially colorblind. Therefore, poison baits can be dyed for safety. Poison grains and pellets which have been dyed a bright color can easily be avoided by humans.


Bait Stations

Place rodenticides in bait stations to increase their effectiveness and safety. Bait stations protect bait from moisture, dust, and weather; provide a protected place for rodents to feed, allowing them to feel more secure; keep nontarget animals away from bait; and help prevent accidental spilling of bait. In addition, baitstations allow easy inspection of bait to determine use.


Bait stations do not have to be elaborate structures. They can be made by nailing flat boards at an angle to the wall. This provides a protected area between the board and the wall.

An 18-inch length of pipe (2 to 3 inch diameter) may be used as a bait station. A more elaborate design is a wooden box with a 2½-inch hole in each end. The top of the box is hinged to allow inspection of the bait. Some of the rodenticides may be purchased in a commercially manufactured bait-station.

Bait stations should be placed in areas of rodent activity. Placing the stations between the ro-



A bait station made by nailing a flat board to a wall.



A bait station made from a length of pipe.



dent's shelter and food supply will increase exposure and use. Bait stations should be placed next to walls, as rats and mice normally use these areas as travel lanes.

Traps

Traps are the best method of rodent control where poisons are unwanted or inadvisable. Common snap traps, glue boards, and live traps may all be effective if rat numbers are low. For roof rats, the traps should be placed along beams, rafters, or other travel ways. For Norway rats and house mice, traps should be placed under cabinets or next to a wall. Traps may be baited with a variety of food items; peanut butter makes a good bait that adheres to the trap trigger.

Summary

Controlling rats and mice in buildings involves a combination of exclusion, sanitation, and wise use of rodenticides. Wherever possible, rodents should be excluded by blocking or sealing all areas where they may enter the building. A judicious program of sanitation will help control rat populations before they increase. Rodenticides may be used to bring problem rat populations into check. However, poisons alone will not provide long-term relief from rat infestations. Rodenticides must be used along with exclusion and sanitation efforts. Trapping may be effective where rat or mouse populations are low.



Placement of traps.



ANR-688

James B. Armstrong, *Extension Wildlife Scientist*, Associate Professor, Forestry and Wildlife Science, Auburn University

Portions of this publication adapted from: *Prevention and Control of Wildlife Damage*, R.M. Timm, editor. Nebraska Cooperative Extension Service, University of Nebraska–Lincoln.

For more information, call your county Extension office. Look in your telephone directory under your county's name to find the number.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, Acts of May 8 and June 30, 1914, and other related acts, in cooperation with the U.S. Department of Agriculture. The Alabama Cooperative Extension System (Alabama A&M University and Auburn University) offers educational programs, materials, and equal opportunity employment to all people without regard to race, color, national origin, religion, sex, age, veteran status, or disability. 6M, Reprinted Nov 2003, ANR-688