


IPM



Homeowner Lawns

Chemical Weed Control Recommendations for 2012

There is nothing simple, automatic, or easy about controlling weeds in home lawns. Each lawn or home situation is different. There are different lawn grasses, different soils, different ornamental plants, and different weed problems. Also, different degrees of management will be used in maintaining the lawn. All of these factors must be considered when attempting to control weeds.

The first step in any weed control program is to identify the weed or weeds that are a problem. Often, a homeowner cannot identify the different weeds. Personnel in the county Extension office, local nurseries, and lawn and garden centers can provide the necessary identification. Once problem weeds have been identified, it is then important to select the appropriate herbicide product for the situation and target weeds. No single product will control all the weeds, and no single product can be used in all lawn grasses or in all situations.

Herbicide products available to homeowners are packaged in small containers—typically in pints and quarts. These products are manufactured in diluted forms for ease of application and for safety as well as to reduce the overall cost. Each product contains one or more different herbicides. Although several products sold by different companies contain the same herbicides, they may differ in concentration of herbicides, ratios, and/or chemical form. Selecting the herbicide product to use is made more difficult by the fact that lawn grasses behave or respond differently to each herbicide. Some herbicide products will have different rates for different lawn grasses. The applicator should abide by these rates when making the application.

The product label is the best source of information as to which lawn grasses can be treated with the particular product. Read the label carefully before purchase and use the information to ensure safety to the applicator as well as the home lawn setting.

There are several terms that are used throughout this section on weed control which should be defined.

Preemergence herbicide: A herbicide that is applied to the lawn grass surface before problem weeds germinate and emerge from the ground. These herbicides must be applied before a weed problem develops. Usually, preemergence herbicides are ineffective when applied to emerged weeds.

Postemergence herbicide: A herbicide that is applied after weeds have emerged, but while they are small and actively

growing. This type of herbicide is applied to the leaf and stem tissue of the problem weeds. Generally, postemergence herbicides will not control weeds that germinate and emerge after the application.

Non-selective herbicide: A herbicide that kills all vegetation treated. Typically, these products are applied to the foliage of the weeds. This type herbicide will also severely injure or kill desirable plants. The non-selective herbicide products listed in this publication do not have any soil activity; that is, they are not soil sterilants.

RTU: An abbreviation that stands for “Ready To Use.” It is used to designate the products that are manufactured in a ready-to-use state. In other words, the product requires no additional water or handling and is typically available in a spray- or applicator-type container.

“Green-up” period (transition period): A short period of time in the spring and/or fall when desirable lawn grass is emerging from its dormant state. It is during this period that grasses are most sensitive to herbicides and, in many cases, the herbicide labels prohibit their use.

Granular herbicide forms can be applied with a drop-type fertilizer spreader. Herbicide products available in a liquid form must be mixed with water prior to use and applied in some type of applicator which will provide uniform application. Homeowners typically use devices such as hose-end sprayers, pressure- or pump-type sprayers, or the more sophisticated backpack sprayers. Generally, equipment used for applying herbicides should no longer be used for treating desirable plants. It is difficult to remove all traces of herbicides from spray applicators. It is possible that herbicide contamination could cause problems on desirable shrubs, flowers, or trees.

The volume of water or spray solution to use when making a herbicide application is suggested on the herbicide container label. Generally, when pump-up sprayers are to be used, most herbicide labels suggest that 2 to 4 gallons of herbicide spray mix be applied per 1000 square feet of lawn surface. However, when hose-end sprayers are used, many herbicide labels suggest that 4 to 6 gallons of herbicide spray mix be applied per 1000 square feet of lawn surface. Read the label carefully to determine the spray volume as well as the other necessary application instructions before using the herbicide. Following is a list of herbicides currently recommended for use on home lawns.

Table 1. Herbicides Recommended for Controlling Weeds on Home Lawns

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.	Lawn Grass Type	Comments
			Preemergence
atrazine			
IMAGE HERBICIDE FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS WITH ATRAZINE (4%)	8.6 fl.oz.	Centipedegrass, St. Augustinegrass	Provides preemergence and postemergence control of broadleaf weeds and annual bluegrass. Control of summer annual grasses is weak. Atrazine may be applied anytime to St. Augustine and centipede grasses except during "green up." Best results occur when application is made before weeds emerge or when still small. Two applications of atrazine during a single year may be needed to control some hardy weeds. See label for appropriate rate and times of application. DO NOT apply to cool-season grasses such as fescue, ryegrass, or bluegrass. DO NOT reseed for 4 months before or 6 months after herbicide application. Rain or irrigation is needed within 7 days after application for consistent control. See label for additional application instructions.
SPECTRACIDE WEEDSTOP FOR LAWNS (4%)	8.6 fl.oz.	Centipedegrass, St. Augustinegrass	
HI-YIELD ATRAZINE WEED KILLER	8.6 fl.oz.	Centipedegrass, St. Augustinegrass	
SA-50 BRAND ATRAZINE ST. AUGUSTINE WEED KILLER (4%)	8.6 fl.oz.	Centipedegrass, St. Augustinegrass	
benfen + trifluralin			
HI-YIELD CRABGRASS CONTROL 2G	2.25-3.5 lb.	Bermudagrass, Centipedegrass, Kentucky bluegrass, Ryegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	Provides preemergence control of annual grasses and several broadleaf weeds. Irrigate treated area after application to activate herbicide. DO NOT apply to newly sprigged areas. DO NOT reseed treated areas.
dithiopyr			
HI-YIELD T/O WEED AND GRASS STOPPER (0.125%)	See label.	Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	Provides preemergence control of annual grasses such as crabgrass and foxtails and some small-seeded broadleaf weeds such as spurge. Will provide postemergence control of small crabgrass less than 4 weeks old.
STA-GREEN CRAB-EX CRABGRASS PREVENTER	3.4 lb.	Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
GREENLIGHT CRABGRASS PREVENTER 2	2 lb.	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.	Lawn Grass Type	Comments
Preemergence (cont.)			
isoxaben			
GREEN LIGHT PORTRAIT BROADLEAF WEED PREVENTER (0.38%G)	4.76 lb.	Bermudagrass, Centipede grass, Kentucky bluegrass, St. Augustine grass, Tall fescue, Zoysiagrass	Provides preemergence control of winter annual broadleaf weeds when applied in early fall. Rainfall or irrigation needed for activation. DO NOT reseed for 4 months.
FERTI-LOME BROADLEAF WEED CONTROL WITH GALLERY (0.38%G)	4.6-5.7 lb.	Bermudagrass, Centipede grass, Kentucky bluegrass, St. Augustine grass, Tall fescue, Zoysiagrass	
pendimethalin			
SCOTT'S HALTS CRABGRASS PREVENTER (1.71%G)	2 lb.	Bermudagrass, Centipede grass, Kentucky bluegrass, St. Augustine grass, Tall fescue, Zoysiagrass	Provides preemergence control of summer grasses. Application should be made in early spring before weed seeds germinate. For goosegrass control in the spring, a second application or treatment should be made 8 weeks after the first application. Can be applied in late August or early September for annual bluegrass and chickweed control. DO NOT apply within 5 months after sprigging grass in lawn; DO NOT reseed area for 4 months after application. DO NOT use on newly seeded areas until after the fourth mowing.
Postemergence			
2,4-D			
HI-YIELD 2,4-D AMINE NO. 4	0.75-1.5 fl.oz.	Bermudagrass, Zoysiagrass	Provides postemergence control of many broadleaf weeds, wild garlic, and wild onion. Best results are obtained when weeds are small and actively growing. A second treatment in 10 to 14 days may be needed to control hardy weeds. Spray only when air is calm to avoid drift to sensitive plants. DO NOT water lawn for 24 hours after applying. DO NOT apply herbicide within the dripline of trees, shrubs, and flowers. Use half the recommended rate on St. Augustine grass or on newly seeded lawns. DO NOT use when temperatures are above 85°F. For wild onion or garlic control, treat in November or December and again in February. Repeat program for 3 consecutive years. Treat ONLY if day temperature is above 60°F for several days before treatment.
SA BRAND 2,4-D AMINE WEED KILLER	1-1.5 fl.oz.	Bermudagrass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.		Lawn Grass Type	Comments
	Postemergence (cont.)			
2,4-D + dicamba + quinelorac + sulfentrazone				
SPECTRACIDE WEED STOP FOR LAWNS PLUS CRABGRASS KILLER READY-TO-SPRAY	6.4 fl.oz.		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides the broadest spectrum of postemergence broadleaf and grass weed control. Apply to actively growing weeds of correct size. When treating bermudagrass, do not apply when day temperatures exceed 85°F.
SPECTRACIDE WEED STOP FOR LAWNS PLUS CRABGRASS KILLER CONCENTRATE	6.4 fl.oz.		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
SPECTRACIDE WEED STOP FOR LAWNS PLUS CRABGRASS KILLER RTU	RTU		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
2,4-D + MCPP + dicamba				
ACE LAWN WEED KILLER	4 fl.oz.		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides the broadest spectrum of broadleaf weed control. Best results are obtained when weeds are small and actively growing. To ensure actively growing weeds, water 2 to 3 days prior to spraying. This combination works faster when applied at temperatures above 70°F. Use the low rate on St. Augustine and centipede grasses and the high rates on zoysia, bluegrass, and fescue lawns. Some of the products listed permit only spot treatment on St. Augustine and centipede lawns to reduce overall injury. DO NOT water for 24 hours after application. DO NOT apply when air temperature is expected to exceed 85°F. DO NOT spray within the dripline or root zone of desirable trees, shrubs, flowers, or vegetables. DO NOT reseed treated areas for 3 to 4 weeks after use. Make application in the spring when weeds are small and actively growing. Hard-to-control weeds may require a second treatment 3 to 4 weeks later. Apply when air is calm to avoid spray drift to desirable ornamental plants. DO NOT apply when air temperature is above 85°F. DO NOT water for 48 hours after use. Water regularly thereafter to maintain a quality lawn. On newly established lawns, DO NOT use until grass has been mowed three or four times. DO NOT reseed sod or sprig treated areas for 3 weeks after use.
BAYER WEED KILLER FOR LAWNS	RTU		Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
BAYER WEED KILLER FOR LAWNS CONCENTRATE	2-4 fl.oz.		Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
BONIDE WEED BEATER LAWN WEED KILLER CONCENTRATE	2-4 fl.oz.		Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
BONIDE WEED BEATER LAWN WEED KILLER READY-TO-SPRAY	See label. Hose-end		Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
ENFORCER WEED SHOT LAWN WEED KILLER	RTU		Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
ENFORCER WEED SHOT LAWN WEED KILLER CONCENTRATE	4-8 fl.oz. Hose-end		Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.	Lawn Grass Type	Comments
Postemergence (cont.)			
2,4-D + MCPP + dicamba (cont.)			Provides the spectrum of broadleaf weed control. Best results are obtained when weeds are small and actively growing. To ensure actively growing weeds, water 2 to 3 days prior to spraying. This combination works faster when applied at temperatures above 70°F. Use the low rate on St. Augustine and centipede grasses and the high rates on zoysia, bluegrass, and fescue lawns. Some of the products listed permit only spot treatment on St. Augustine and centipede lawns to reduce overall injury. DO NOT water for 24 hours after application. DO NOT apply when air temperature is expected to exceed 85°F. DO NOT spray within the dripline or root zone of desirable trees, shrubs, flowers, or vegetables. DO NOT reseed treated areas for 3 to 4 weeks after use. Make application in the spring when weeds are small and actively growing. Hard-to-control weeds may require a second treatment 3 to 4 weeks later. Apply when air is calm to avoid spray drift to desirable ornamental plants. DO NOT apply when air temperature is above 85°F. DO NOT water for 48 hours after use. Water regularly thereafter to maintain a quality lawn. On newly established lawns, DO NOT use until grass has been mowed three or four times. DO NOT reseed sod or sprig treated areas for 3 weeks after use.
FERTI-LOME WEED-OUT LAWN WEED KILLER CONCENTRATE	2-4 fl.oz.	Bermudagrass, Centipede grass, Kentucky bluegrass, St. Augustine grass, Tall fescue, Zoysiagrass	
GORDON'S TRIMEC LAWN WEED KILLER	2-4 fl.oz.	Bermudagrass, Centipede grass, Ryegrass, St. Augustine grass, Tall fescue, Zoysiagrass	
GORDON'S TRIMEC READY SPRAY LAWN WEED KILLER	4-6 fl.oz. RTS	Bermudagrass, Centipede grass, Ryegrass, St. Augustine grass, Tall fescue, Zoysiagrass	
ORTHO WEED-B-GON MAX WEED KILLER FOR LAWNS READY TO USE	RTU	Bermudagrass, Centipede grass, Tall fescue, Zoysiagrass	
PREEN LAWN BROADLEAF WEED CONTROL	2 lb.	Bermudagrass, Centipede grass, Fescue, Kentucky bluegrass, Zoysiagrass	
SPECTRACIDE WEED STOP FOR LAWNS	RTU	Bermudagrass, Centipede grass, St. Augustine grass, Zoysiagrass	
SPECTRACIDE WEED STOP FOR LAWNS READY TO SPRAY	RTS	Bermudagrass, Centipede grass, St. Augustine grass, Zoysiagrass	
SPECTRACIDE WEED STOP FOR LAWNS CONCENTRATE	2-4 fl.oz.	Bermudagrass, Centipede grass, Kentucky bluegrass, St. Augustine grass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.	Lawn Grass Type	Comments
Postemergence (cont.)			
2,4-D + MCPP + dicamba + carfentrazone			Provides broadest spectrum of postemergence broadleaf weed control. Apply to small, actively growing weeds. Do not apply during "spring transition." Treatment is effective in controlling a number of winter weeds. DO NOT apply when day temperature is above 90°F. Use low rate on bermudagrass and zoysia lawns. RTU products are available.
FERTI-LOME WEED-FREE ZONE	0.75-2 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
GORDON'S SPEED-ZONE LAWN WEED KILLER	1.1-1.4 fl.oz.	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
ORTHO WEED-B-GON MAX FOR SOUTHERN LAWNS CONCENTRATE	3-6 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
ORTHO WEED-B-GON MAX FOR SOUTHERN LAWNS READY-TO-SPRAY	3-6 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
ORTHO WEED-B-GON MAX FOR SOUTHERN LAWNS READY-TO-USE	RTU	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
2,4-D + MCPP + dicamba + dithiopyr			Apply spring or fall to small actively growing grasses and broadleaf weeds. In spring, apply when first dandelions flower. Make second application 6 to 8 weeks later for best weed and crabgrass control. In fall, apply when day temperatures cool and weeds are small and actively growing (around Labor Day).
SPECTRACIDE WEED STOP FOR LAWNS CRABGRASS PREVENTER AND BROADLEAF WEED KILLER (1% GRANULE)	3.6 lb.	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
2,4-D + MCPP + dicamba + isoxaben			Provides broadest spectrum of broadleaf weed control. Use low rate on centipedegrass and St. Augustinegrass. Use high rate on bermudagrass and zoysia lawns. Water 1 to 2 days before application and do not mow for 1 to 2 days after application. Apply to small, actively growing weeds when day temperatures are between 50 and 90° F. This product will provide residual control of many broadleaf weeds.
BAYER SOUTHERN SEASON LONG WEED CONTROL FOR LAWNS CONCENTRATE	3.2-6.4 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
BAYER SOUTHERN SEASON LONG CONTROL FOR LAWNS READY-TO-SPRAY	3.2-6.4 fl.oz. RTS	Bermudagrass, Centipedegrass, Kentucky bluegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.	Lawn Grass Type	Comments
Postemergence (cont.)			
2,4-D + quinelorac + dicamba GREENLIGHT WIPE-OUT CRABGRASS KILLER PLUS CONCENTRATE	5 fl.oz.	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides broad spectrum control of broadleaf weeds and crabgrass. Apply in temperature range of 45 to 90°F. DO NOT apply to bermudagrass when day temperatures are above 85°F.
2,4-D + triclopyr + dicamba ORTHO WEED-B-GON MAX CONCENTRATE	2-4 fl.oz.	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides broad spectrum weed control. Use low rate on bermudagrass and zoysia and high rate on fescue and bluegrass. DO NOT apply to centipede or St. Augustinegrass.
ORTHO WEED-B-GON MAX READY TO SPRAY	2-4 fl.oz.	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
bentazon BASAGRAN T/O	0.75 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, Ryegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	Apply when yellow nutsedge is actively growing under good moisture conditions. Apply at 10- to 14-day intervals until nutsedge is controlled. DO NOT make more than three applications in a single growing season. Avoid spraying adjacent trees, shrubs, and flowers. DO NOT mow 3 to 5 days before or after application. Purple nutsedge will not be controlled by this herbicide. See label for recommendations about adding crop oil concentrate for control of some weeds, including yellow nutsedge.
HI-YIELD BASAGRAN	0.75 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, Ryegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
SA BRAND BASAGRAN SEDGE CONTROL	0.75 fl.oz.	Bermudagrass, Centipedegrass, Kentucky bluegrass, Ryegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	
fenoxyprop BAYER ADVANCED BERMUDAGRASS CONTROL FOR LAWNS READY-TO-SPRAY	6.4 fl.oz.	Kentucky bluegrass, Ryegrass, Tall fescue	Provides control of bermudagrass in select cool-season grasses. See label for other grasses controlled. DO NOT apply to warm-season turfgrasses.
halosulfuron GOWAN SEDGEHAMMER 75DF	0.9 gram	Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	Controls yellow and purple nutsedge and suppresses green kyllinga. Follow label directions for mixing and use. Apply to nutsedge with three to eight leaves. A second application may be needed 6 to 10 weeks after the first. DO NOT mow for 2 days before or after application. DO NOT apply in vegetable or flower gardens. Add a non-ionic surfactant at 2 teaspoons per gallon of spray mix.
HI-YIELD NUTSEDGE CONTROL 75WDG	0.9 gram	Bermudagrass, Centipedegrass, St. Augustinegrass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate:		Comments
	Product Amount per 1000 Sq. Ft.	Lawn Grass Type	
Postemergence (cont.)			
imazaquin			
IMAGE HERBICIDE CONSUMER CONCENTRATE (3.3%) KILLS NUTSEDGE	3.75 fl.oz.	Bermudagrass, Centipedegrass, St. Augustinegrass, Zoysiagrass	To control summer weeds, including nutsedge, apply after turf "green-up" in spring when weeds are small and actively growing. Irrigate treated area 1 to 7 days after applying. To control winter weeds, including wild garlic, treat in the fall after first frost when weeds are small and actively growing. Mowing 1 to 2 weeks after treatment will improve appearance and control. DO NOT apply on newly seeded, sprigged, or sodded turfgrass. DO NOT apply to St. Augustinegrass in the fall for winter weed control or on any labeled grass that does not go dormant.
quinclorac + 2,4-D + dicamba			
ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL READY-TO-SPRAY	6.4 fl.oz.	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	Provides broad spectrum control of postemergence broadleaf and grass weeds. Do not apply when temperature is above 85°F. See label for appropriate timing and weed size. Make only two applications per year.
ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL CONCENTRATE	6.25 fl.oz	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	
BAYER ALL-IN-ONE LAWN WEED AND CRABGRASS KILLER READY-TO-SPRAY	6.25 fl.oz	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	
BAYER ALL-IN-ONE LAWN WEED AND CRABGRASS KILLER	RTU	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	
BAYER ALL-IN-ONE LAWN WEED AND CRABGRASS KILLER CONCENTRATE	6.4 fl.oz.	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	
BONIDE WEED BEATER PLUS CRABGRASS AND BROADLEAF WEED KILLER	See label.	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.	Lawn Grass Type	Comments
Postemergence (cont.)			
quinclorac + 2,4-D + dicamba (cont.)			
BONIDE WEED BEATER PLUS CRABGRASS KILLER READY-TO-SPRAY	4.8 fl.oz. RTS	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides broad spectrum control of postemergence broadleaf and grass weeds. Do not apply when temperature is above 85°F. See label for appropriate timing and weed size. Make only two applications per year.
FERTI-LOME WEEDOUT WITH Q	6.4 fl.oz.	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	
GORDON'S TRIMEC CRABGRASS PLUS LAWN WEED KILLER CONCENTRATE	6.4 fl.oz.	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	
GORDON'S TRIMEC CRABGRASS PLUS LAWN WEED KILLER READY-TO-SPRAY	6.4 fl.oz. RTS	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
GORDON'S TRIMEC CRABGRASS PLUS LAWN WEED KILLER READY-TO-USE	RTU	Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
quinclorac + MCPP + 2,4-D + dicamba			
ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL	RTU	Bermudagrass, Kentucky bluegrass, Ryegrass, Tall fescue, Zoysiagrass	Provides broad spectrum control of postemergence broadleaf and grass weeds. Apply within temperature range of 45 to 90°F. Apply to actively growing weeds at correct time and size.
quinclorac + sulfentrazone			
IMAGE HERBICIDE KILLS CRABGRASS II	24 gm/3 gal./1500 sq.ft.	Bermudagrass, Centipedegrass Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides very broad spectrum control of broadleaf, grasses, and sedges. Apply to small actively growing crabgrass and other weeds. Apply only one time per year.

Herbicide and Trade Name Examples	Application Rate: Product Amount per 1000 Sq. Ft.		Lawn Grass Type	Comments
	Postemergence (cont.)			
sethoxydim				
BONIDE GRASS BEATER CONCENTRATE	0.8 fl.oz.		Centipedegrass	Controls annual grasses and suppresses bahiagrass growth. DO NOT make application within 3 weeks after “green up.” The low rate can be used on newly planted centipede when 3 inches of stolon growth has occurred. Two applications can be made to established centipedegrass for bahiagrass suppression. Time the second application 10 to 14 days after the first application. DO NOT mow the centipede lawn for 7 days before or after application. DO NOT mix Vantage or Poast with any other herbicide or pesticide product. Will not control nutgrass or any broadleaf weeds. Add 1.5 tablespoons of crop oil concentrate per 1000 square feet when applying Poast or Bonide.
POAST	0.375-0.5 fl.oz.		Centipedegrass	
SA BRAND VANTAGE	0.5-1.4 fl.oz.		Centipedegrass	
SEGMENT	0.5-0.8 fl.oz.		Centipedegrass	
VANTAGE	0.5-0.75 fl.oz.		Centipedegrass	
MCPA + MCPP+ dicamba + carfentrazone				
BONIDE WEED BEATER ULTRA CONCENTRATE	0.75-1.8 fl.oz.		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	Controls a number of winter broadleaf weeds when weeds are small and actively growing. Consistent control under cooler temperature conditions. Apply when temperature is above 45°F but less than 85°F. Use low to medium rate on bermudagrass and zoysiagrass. See label directions.
BONIDE WEED BEATER ULTRA RTS	0.5-1 fl.oz.		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
BONIDE WEED BEATER ULTRA RTU	RTU		Bermudagrass, Kentucky bluegrass, Tall fescue, Zoysiagrass	
triclopyr				
ORTHO WEED-B-GON CHICKWEED, CLOVER, AND OXALIS KILLER (8%)	5 fl.oz.		Kentucky bluegrass, Tall fescue, Zoysiagrass	Provides control of difficult-to-control broadleaf weeds. Apply to small actively growing weeds, but before day temperatures exceed 90°F. Retreatment can be made 2 to 3 weeks after first application if necessary.
MCPA + triclopyr + dicamba				
BONIDE CHICKWEED, CLOVER, AND OXALIS KILLER CONCENTRATE	2-4 fl.oz.		Bermudagrass, Kentucky bluegrass, Tall fescue, Ryegrass, Zoysiagrass	Provides control of a number of broadleaf weeds in cool-season turfgrasses and several warm-season turfgrasses. Use low rate on bermudagrass and zoysia lawns. Do not apply when temperature is above 90°F. Keep away from desirable trees and shrubs.
BONIDE CHICKWEED, CLOVER, AND OXALIS KILLER READY-TO-USE	RTU		Bermudagrass, Kentucky bluegrass, Tall fescue, Ryegrass, Zoysiagrass	
MCPP + 2,4-D + dicamba				
SOUTHERN AG LAWN WEED KILLER WITH TRIMEC	2.5-5 fl.oz.		Bermudagrass, Centipedegrass, Kentucky bluegrass, Tall fescue, St. Augustinegrass	Provides broadleaf weed control with less injury in 2,4-D-sensitive grasses. Apply to small, actively growing weeds not growing under stress. Apply before day temperatures exceed 90°F.

Table 2. Effectiveness of Selected Herbicides on Common Lawngrass Weeds

Weeds	Preemergence				Postemergence	
	atrazine	bensulide	dithiopyr	pendimethalin	bentazon	2,4-D
Annual bluegrass	X	X	X	X	--	--
Annual lespedeza	X	--	--	--	--	X
Bahiagrass	--	--	--	--	--	--
Bittercress	--	--	--	--	--	X
Common chickweed	X	--	X	X	X	X
Common greenbrier	--	--	--	--	--	--
Common woodsorrel	X	--	X	--	--	X
Cudweed	--	--	--	--	--	X
Dandelion	X	--	--	--	--	X
Dichondra	X	--	--	--	--	X
False dandelion	X	--	--	--	--	X
Florida betony	X	--	--	--	--	X
Goosegrass	--	X	X	X	--	--
Gripeweed	X	--	X	--	--	--
Henbit	X	X	X	X	--	X
Large crabgrass	X	X	X	X	--	--
Lawn burweed	X	--	--	--	X	X
Pennywort	X	--	--	--	--	X
Prostrate spurge	X	--	X	X	--	X
Purple nutsedge	--	--	--	--	--	--
Virginia buttonweed	--	--	--	--	--	X*
Wild garlic	--	--	--	--	--	X*
Wild violets	--	--	--	--	--	X*
Yellow nutsedge	--	--	--	--	X	--

Weeds	Postemergence					
	2,4-D + MCPP + dicamba	2,4-D + quinclorac + dicamba	imazaquin	halosulfuron	MCPP + MCPA + dicamba + carentrazone	sethoxydim
Annual bluegrass	--	--	--	--	--	--
Annual lespedeza	X	X	--	--	X	--
Bahiagrass	--	--	--	--	--	X*
Bittercress	X	X	--	--	X	--
Common chickweed	X	X	X	--	X	--
Common greenbrier	--	--	--	--	--	--
Common woodsorrel	X	X	--	--	X	--
Cudweed	X	X	--	--	X	--
Dandelion	X	X	--	--	X	--
Dichondra	X	X	--	--	X	--
False dandelion	X	X	--	--	X	--
Florida betony	X	X	--	--	--	--
Goosegrass	--	--	--	--	--	X
Gripeweed	X	--	--	--	--	--
Henbit	X	X	--	--	X	--
Large crabgrass	--	X	--	--	--	X
Lawn burweed	X	X	X	--	X	--
Pennywort	X	X	--	--	X	--
Prostrate spurge	--	X	--	--	X	--
Purple nutsedge	--	--	X	X	--	--
Virginia buttonweed	--	--	--	--	X	--
Wild garlic	X	X	X	--	X	--
Wild violets	X	X	--	--	--	--
Yellow nutsedge	--	--	X	X	--	--

X = Weed controlled by herbicide. * = Suppression only. -- = No control.

Table 3. Landscape Maintenance Herbicides for Home Lawns

CHEMICAL EDGERS AND TOTAL VEGETATION CONTROL PRODUCTS		
Herbicide Trade Name Examples	Application Rate	Site
IMAGE YEAR LONG VEGETATION KILLER (dichlobenil)	4 lb./400 sq.ft.	Non-selective
GORDON'S LIQUID EDGER 2 (glyphosate 1%)	1 gal/750 sq.ft.	
GORDON'S PRONTO FAST ACTING VEGETATION KILLER (imazapyr + glyphosate)	18 fl.oz/3-5 gal water/1000 sq.ft.	
ORTHO GROUND CLEAR VEGETATION KILLER CONCENTRATE (glyphosate + imazapyr)	1gal./300 sq.ft.	Non-selective
ORTHO GROUND CLEAR VEGETATION KILLER (RTU) (glyphosate + imazapyr)	1.25 gal./75 sq.ft.	Non-selective
NON-SELECTIVE LANDSCAPE MAINTENANCE		
Herbicide Trade Name Examples	Application Rate	
2,4-D + 2,4-DP + dicamba		
GORDONS ALL SEASON BRUSH NO-MORE	3-5 fl.oz./gal. water	
2,4-D + MCPP + dicamba		
ENFORCER POISON IVY AND OAK KILLER	RTU	
diquat		
ACE GRASS AND WEED KILLER	See label.	
ENFORCER NEXT DAY GRASS AND WEED KILLER CONCENTRATE (2.38%)	See label.	
diquat + fluazifop + dicamba		
SPECTRACIDE WEED AND GRASS KILLER CONCENTRATE 2	7 fl.oz./gal.	
SPECTRACIDE WEED AND GRASS KILLER	RTU	
diquat + fluazifop + dicamba + oxyfluorfen		
SPECTRACIDE WEED AND GRASS KILLER WITH EXTENDED CONTROL	RTU	
fluazifop		
FERTI-LOME OVER-THE-TOP GRASS KILLER (1.7%)	10 fl.oz./gal./1000 sq.ft.	
ORTHO GRASS-B-GON (0.48%)	RTU	
glyphosate		
BONIDE KLEENUP SYSTEMIC WEED AND GRASS CONCENTRATE (41%)	1.5-2.5 fl.oz./gal. water	

continued

NON-SELECTIVE LANDSCAPE MAINTENANCE (cont.)

Herbicide Trade Name Examples	Application Rate
glyphosate (cont.)	
BONIDE KLEENUP WEED & GRASS KILLER (1.92%)	RTU
ENFORCER ROOTS AND ALL LAWN WEED KILLER (1.92%)	RTU
ENFORCER ROOTS AND ALL CONCENTRATE (27%)	See label.
ENFORCER ROOTS AND ALL SUPER CONCENTRATE (41%)	1.5-2.5 fl.oz./gal. water
ENFORCER BRUSH KILLER CONCENTRATE (27%)	See label.
GORDON'S PRONTO WEED AND GRASS KILLER (2%)	RTU
GORDON'S PRONTO WEED AND GRASS KILLER CONCENTRATE (25%)	4-5 fl.oz./gal. water
GORDON'S PRONTO FAST-ACTING BRUSH KILLER (32.3%)	3-9 fl.oz./gal. water/300 sq.ft.
GREEN LIGHT COM-PLEET (41%) SYSTEMIC GRASS AND WEED KILLER 2	5 fl.oz./gal. water
GREEN LIGHT COM-PLEET SYSTEMIC GRASS AND WEED KILLER (1.92%)	RTU
HI-YIELD KILLZALL WEED AND GRASS KILLER	RTU
HI-YIELD SUPER CONCENTRATE KILLZALL (41%)	2.5-5 oz./gal. water
ROUNDUP POISON IVY AND TOUGH BRUSH KILLER PUMP AND GO SPRAYER (glyphosate 1% and triclopyr 0.1%)	RTU
ROUNDUP POISON IVY AND TOUGH BRUSH KILLER (glyphosate 1% and triclopyr 0.1%)	RTU
ROUNDUP CONCENTRATE POISON IVY PLUS TOUGH BRUSH KILLER (glyphosate 18% and triclopyr 2%)	6 fl.oz./gal. water
ROUNDUP PRO (41%)	2.5 fl.oz./gal. water
ROUNDUP READY-TO-USE WEED AND GRASS KILLER (2%) + pelargonic acid (2%)	RTU
ROUNDUP EXTENDED CONTROL WEED AND GRASS KILLER PLUS WEED PREVENTER II (glyphosate + pelargonic acid + imazapic)	RTU

continued

NON-SELECTIVE LANDSCAPE MAINTENANCE (cont.)

Herbicide Trade Name Examples	Application Rate
glyphosate (cont.)	
ROUNDUP EXTENDED CONTROL WEED AND GRASS KILLER PLUS WEED PREVENTER (CONCENTRATE) (glyphosate 18%, imazapic 0.3%, diquat 0.74%)	21 fl.oz./3 gal. water/1000 sq.ft.
SOUTHERN AG WEEDGUN (0.96%)	RTU
SOUTHERN AG WEEDPRO (41%)	1.5-2.5 fl.oz./gal. water
glyphosate + oxyfluorfen	
ORTHO SEASON LONG MAX READY TO USE	RTU
glyphosate + oxyfluorfen + diquat	
ORTHO SEASON LONG MAX CONCENTRATE	See label.
MCPA + triclopyr + dicamba	
BONIDE BRUSH KILLER, BK 32	3.2 fl.oz./gal. water
BONIDE BRUSH KILLER, BK 32	RTU
prometon	
BONIDE TOTAL VEGETATION KILLER (2.2%)	1 gal./500 sq.ft.
ENFORCER SCORCHER TOTAL VEGETATION KILLER (2%)	1 gal./500 sq.ft.
triclopyr	
BONIDE STUMP OUT AND VINE KILLER	RTU
FERTI-LOME BRUSH KILLER/STUMP KILLER (8.8%)	4-8 fl.oz./gal. water
GREEN LIGHT CUT VINE AND STUMP KILLER (0.8%)	RTU
IMAGE BRUSH AND VINE KILLER (8.8%)	4 fl.oz./gal. water/1000 sq.ft.
IMAGE BRUSH AND VINE KILLER	RTU
ORTHO MAX POISON IVY AND TOUGH BRUSH KILLER CONCENTRATE (8%)	4 fl.oz./gal. water
ORTHO MAX POISON IVY AND TOUGH BRUSH KILLER (0.7%)	RTU
SA BRAND BRUSH KILLER (8.8%)	4-8 fl.oz./gal. water
VINE-X VINE AND BRUSH CONTROL (13.6%)	RTU

Table 4. Relative Tolerance of Selected Lawn Grasses to Herbicides

Herbicide	Bermudagrass	Centipedegrass	St. Augustinegrass	Tall fescue	Zoysiagrass
2,4-D amine	Yes	Yes*	Yes*	Yes	Yes
2,4-DP	Yes	Yes*	Yes*	No	Yes
atrazine	Yes*	Yes	Yes	No	Yes*
benefin	Yes	Yes	Yes	No*	Yes
bensulide	Yes	Yes	Yes	Yes	Yes
bentazon	Yes	Yes	Yes	Yes	Yes
dicamba	Yes	Yes*	Yes*	Yes	Yes
dithiopyr	Yes	Yes	Yes	Yes	Yes
glyphosate	No	No	No	No	No
halosulfuron	Yes	Yes	Yes	Yes	Yes
imazaquin	Yes	Yes	Yes	No	Yes
isoxaben	Yes	Yes	Yes	Yes	Yes
MCP	Yes	Yes	Yes	Yes	Yes
pendimethalin	Yes	Yes	Yes	No*	Yes
prodiamine	Yes	Yes	Yes	Yes	Yes
sethoxydim	No	Yes	No	No	No

* See label for specific application instructions, rates, and/or timing.

Turf Herbicide Manufacturers and Suppliers

Bayer Advanced LLC
2 TW Alexander Dr.
Research Triangle Park, NC 27709
1-887-229-3724
www.advancedlawn.com/

Lebanon Seaboard Corporation
1600 East Cumberland Street
Labanon, PA 17042
1-800-233-1067
www.preen.com/

Scotts Consumer Service
14111 Scottslawn Road
Marysville, OH 93041
1-888-270-3714
www.scotts.com/

Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424
1-800-536-8231 Ext. 228
www.bonideproducts.com/

Monsanto Company
Lawn & Garden Products
P.O. Box 418
Marysville, OH 43041
1-800-246-7219
www.roundup.com/

Southern Ag Insecticides, Inc.
P.O. Box 218
Palmetto, FL 34221
1-941-722-3285
www.southernag.com/

Enforcer Products Inc.
P.O. Box 1060
Cartersville, GA 30120
1-770-386-0801
www.enforcer.com/

The Ortho Group
P.O. Box 190
Marysville, OH 43040
1-800-225-2883
www.ortho.com/

Spectrum Group
Div. of United Industries Corp.
P.O. Box 142642
St. Louis, MO 63114-0642
1-800-917-5438
www.spectracide.com/
(Spectracide, Sta-Green)

Green Light Company
P.O. Box 17985
San Antonio, TX 78217-0985
1-210-494-3481
www.greenlightco.com/

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, MO 64101
1-800-821-7925
www.pbigordon.com/

VPG, Inc.
P.O. Box 460
Bonham, TX 75418-5000
1-888-583-5296
www.v-p-g.com/
(Fertilome, Hi-Yield)

Gowan Company LLC
370 Main Street
Yuma, AR 72694
1-800-883-1844
www.gowanco.com/

Homeowner Lawns: Chemical Weed Control prepared by John W. Everest, Professor Emeritus, Agronomy and Soils, Auburn University; and Michael G. Patterson, *Extension Weed Scientist*, Professor, Agronomy and Soils, Auburn University.

2012 IPM-0590

For more information, contact your county Extension office. Visit <http://www.aces.edu/counties> or look in your telephone directory under your county's name to find contact information.

Use pesticides **only** according to the directions on the label. Follow all directions, precautions, and restrictions that are listed. Do not use pesticides on plants that are not listed on the label.

The pesticide rates in this publication are recommended **only** if they are registered with the Environmental Protection Agency or the Alabama Department of Agriculture and Industries. If a registration is changed or canceled, the rate listed here is no longer recommended. Before you apply **any** pesticide, check with your county Extension agent for the latest information.

Trade names are used **only** to give specific information. The Alabama Cooperative Extension System does not endorse or guarantee any product and does not recommend one product instead of another that might be similar.

Published by the Alabama Cooperative Extension System (Alabama A&M University and Auburn University), an equal opportunity educator and employer.

