

TIMELY INFORMATION

Agriculture & Natural Resources

September 18, 2007

PP-638

AUGUST PLANT PROBLEM REPORT FROM THE AUBURN PLANT DIAGNOSTIC LAB

AUGUST PLANT PROBLEM REPORT FROM THE BIRMINGHAM PLANT DIAGNOSTIC LAB

AUGUST INSECT REPORT FROM THE AUBURN PLANT DIAGNOSTIC LAB

DISEASE POSSIBILITIES FOR SEPTEMBER

Jackie Mullen

Extension Plant Pathology Specialist-Auburn

Jim Jacobi

Extension Plant Pathology Specialist-Birmingham

Charles Ray

Research Fellow IV-Auburn

Auburn Plant Disease Report-August (J. Mullen)

August was dry in most areas, but especially in the northern half of the state. Temperatures were abnormally high during the first half of the month with 100°F or higher in some parts of the state. We received 85 plant samples in August.

We saw a dieback of white clover, and culture work showed that both *Rhizoctonia solani* and *Sclerotium rolfsii* were associated with the damaged area. See A. Hagan if you have questions.

We saw an unusually severe case of tomato spotted wilt virus (TSWV) on pepper. We have forwarded some of the sample to J. Murphy in the Entomology & Plant Pathology Department at Auburn and to a University of Florida virologist for further tests to be sure no other viruses are present besides TSWV.

We received a few reports of blackened zoysia grass. The black coloration on dying/dead zoysia grass was caused by masses of *Curvularia* spores on the leaf surfaces. This fungus is usually considered to be secondary, developing only on previously damaged plants. It is reported to cause a leaf blight in high temperature situations or after high temperature stress. *Curvularia* is controlled by methods used to control *Bipolaris* (*Helminthosporium*).

During August, Asian soybean rust was identified on soybean in Escambia, Geneva, Marengo, Mobile, Monroe, & Washington Counties. ASBR was found on kudzu and wild legumes in Covington County. For more information on ASBR, contact Ed Sikora and/or Mary Delaney.

Table 1. Plant Diseases Seen In The Auburn Plant Diagnostic Lab in August.

<u>Plant</u>	<u>Disease</u>	<u>County</u>
Apple	Bitter Rot (<i>Colletotrichum gloeosporioides</i>)	Hale
Bean, Green	Anthrachnose (<i>Colletotrichum</i>)	Elmore
Begonia	Pythium Crown Rot	Lee
Bermuda	Bipolaris Crown Rot & Leaf Blight (<i>B. sorokiniana</i>)	Jefferson
Blueberry	Anthrachnose (<i>Colletotrichum</i>)	Dallas
Boxwood	Phytophthora Root Decay	Colbert
Cantaloupe	Fusarium Wilt	Russell
Centipede	Bipolaris Lower Leaf Blight & Suspect Heat Stress	Jefferson
	Take-All Patch & Stress	Elmore
Clover, White	<i>Rhizoctonia solani</i> Blight	Bullock
	Southern Blight (<i>Sclerotium rolfsii</i>)	Bullock
Corn	Northern Corn Leaf Blight (<i>Bipolaris turcicum</i>)	Monroe
	Southern Rust (<i>Puccinia maydis</i>)	Lee, Monroe
Cucumber	Powdery Mildew	Etowah
Gardenia	Phytophthora Root Rot	Baldwin
Grain Sorghum	Root Knot Nematode (<i>Meloidogyne</i>)	Baldwin
	Rough Leaf Spot (<i>Ascochyta sorghina</i>)	Lee

<u>Plant</u>	<u>Disease</u>	<u>County</u>
Holly	Phytophthora Crown & Root Rot	Bullock
Holly, Soft Touch	Phytophthora Root Rot	Autauga
Peach	Brown Rot (<i>Monilinia fructicola</i>)	Macon
	Phytophthora Root Decay	Baldwin
	Root Knot Nematode (<i>Meloidogyne</i> sp.)	Baldwin
Pepper	Tomato Spotted Wilt Virus	Lee
Plum	Plum Leaf Scald (<i>Xylella fastidiosa</i>)	Chilton
Rose	Phytophthora Root Rot	Lee
Soybean	Alternaria Leaf Spot	Morgan
	Anthracnose (<i>Colletotrichum</i>)	Greene
	Charcoal Rot (<i>Macrophomina phaseolicola</i>)	Greene, Tuscaloosa
	Southern Blight (<i>Sclerotium rolfsii</i>)	Geneva
St. Augustine	Gray Leaf Spot (<i>Piricularia grisea</i>)	Mobile
	Take-All Patch (<i>Gaeumannomyces graminis</i> Var. <i>graminis</i>)	Autauga(2), Elmore, Etowah, Jefferson(2), Mobile
Tomato	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Geneva, Russell
	Bacterial Wilt (<i>Ralstonia solanacearum</i>)	Geneva, Macon
Vinca, Annual	Phytophthora Stem Blight (<i>P. nicotiana</i>)	Montgomery
Zoysia, Emerald	Suspect Ring Nematode Problem	Lee
	Take-All Patch	Jefferson
Zoysia, Meyer	Curvularia Leaf Spot/Blight & Heat Stress	Lee, Macon
	Take-All Patch & Stress	Elmore

*Counties are not reported for greenhouse, nursery, and golf course samples.

Monthly Plant Problem Report From The Birmingham Lab (J. Jacobi)

We received 105 samples for the month of August. As expected, many of the samples showed symptoms of environmental stress from ongoing drought. The heat wave during August was especially hard on trees and shrubs that had been transplanted in the last year. For a complete description of the heat wave and the records set during August see the following commentary from the National Weather Service (http://www.srh.noaa.gov/bmx/significant_events/2007/augcli.php).

Some of the other problems seen last month were Phytophthora root rot on aucuba and English boxwood, Phomopsis gall on chestnut oak, Fusarium crown rot on hosta, chinch bugs on St. Augustinegrass, potyvirus on pumpkin, and stem canker on soybean. For more information on hosta diseases and pest here's an excellent reference (<http://www.extension.iastate.edu/Publications/SUL14.pdf>).

Table 2. 2007 August Problems Seen In The Birmingham Plant Diagnostic Lab.

<u>Plant</u>	<u>Problems</u>	<u>County</u>
Azalea	Azalea Lacebug	Jefferson(2)
	Cold Injury	Limestone
	Phytophthora Root Rot	Limestone
Bentgrass	Ectotrophic Root Infecting Fungi	*
	Pythium Root Rot	*
	Ring Nematodes	*
Boxwood, Common	Macrophoma Twig Blight	Jefferson
Boxwood, English	Phytophthora Root Rot	Jefferson
Cleyera Cypress, Leyland	Botryosphaeria Dieback	Jefferson
	Tip Blight (<i>Botryosphaeria</i>)	Jefferson
Euonymus	Euonymus Scale	Jefferson

<u>Plant</u>	<u>Problems</u>	<u>County</u>
Holly, Lusterleaf	Pythium Root Rot	Jefferson
Hosta	Fusarium Crown Rot	Madison
Hydrangea, Bigleaf	Corynespora Leaf Spot	Jefferson
Juniper	Minute Cypress Scale	Jefferson
Maple, Hybrid Cultivars	Maple Spider Mite	Shelby
	Phyllosticta Leaf Spot	Shelby
Mondograss	Anthracnose	Jefferson
	Pythium Root Rot	Jefferson
Oak, Chestnut	Phomopsis Gall	Shelby
Oak, Post	Anthracnose (<i>Discula</i>)	Shelby
Oak, Southern Red	Ganoderma Rot	Jefferson
Oak, Willow	European Fruit Lecanium Scale	Jefferson
	Orangestriped Oakworm	Shelby
	Powdery Mildew	Jefferson
Pear, Asian	Fire Blight	Jefferson
Pecan	Pecan Phylloxera	Jefferson
Periwinkle, Madagascar	Phytophthora Blight	Marengo
Pumpkin	Potyvirus Group	Madison, Shelby
Rhododendron	Phytophthora Root Rot	Jefferson
Rhododendron	Phytophthora Root Rot	Jefferson
Rose	Crown Gall	Shelby

<u>Plant</u>	<u>Problems</u>	<u>County</u>
Rosemary	Pythium Root Rot	*
St. Augustinegrass	Chinch Bugs	Jefferson (3)
Soybean	Stem Canker	Marshall
Sweet Flag	Asochyta Leaf Spot	Shelby
Tobacco, Flowering	Blister Beetle	Shelby
Tomato	Magnesium Deficiency	Jefferson
	Tomato Spotted Wilt Virus	Jefferson

*Counties are not reported for greenhouse, nursery, or golf course samples.

Auburn Entomology Report-August (C. Ray)

County	Host	Category	Identification	Scientific Name
Montgomery	Holly	Ornamental	Forbes Scale	<i>Diaspidiotus forbesi</i>
Lee	Coleus	Ornamental	Eastern Subterranean Termite	<i>Reticulotermes flavipes</i>
Jefferson	Ornamentals	Ornamental	A Blister Beetle	Meloidae
Tuscaloosa	Home	Household-Miscellaneous	White Fringed Beetle	<i>Naupactus</i> sp.
Tuscaloosa	Home	Household-Miscellaneous	Vegetable Weevil	<i>Listriдерes</i> sp.
Houston	Ornamentals	Ornamental	White Fringed Beetles	<i>Naupactus</i> sp.
Houston	Bird Feeder	Household-Miscellaneous	Nymphal Leaf Footed Bugs	Coreidae

County	Host	Category	Identification	Scientific Name
Jefferson	Pantry	Household-Stored Products	Indian Meal Moth	<i>Plodia interpunctella</i>
Covington	Building	Household-Medical	Brown Widow Spider	<i>Latrodectes geometricus</i>
Etowah	Not Available		Nymphal False Chinch Bugs	<i>Nysius raphanus</i>
Jefferson	St. Augustine Sod	Turfgrass	Chinch Bugs	<i>Blissus</i> sp.
Dale	Human	Medical	Nymph Brown Dog Tick	<i>Rhipicephalus sanguineus</i>
Etowah	Home	Household-Miscellaneous	Dust Mite	<i>Dermatophagoides</i> sp.
Mobile (4)	Building	Household-Structural	Powder Post Beetle	<i>Lyctus planicollis</i>
Mobile	Building	Household-Miscellaneous	Juvenile Cockroach	<i>Periplaneta</i> sp. Nymph
Marion	Hay	Forage Crops	Ground Beetle Larvae	Carabidae
Calhoun	Pool	Household-Miscellaneous	White Fringed Beetles	<i>Naupactus</i> sp.
Cullman		Row Crops	False Chinch Bug	<i>Nysius raphanus</i>
Chilton	Home	Household-Stored Product	Foreign Grain Beetle	<i>Ahasverus advena</i>
Chilton	Home	Household-Stored Product	A Rove Beetle	Staphylinidae
Clay	Home	Household-Stored Product	Cigarette Beetle	<i>Lasioderma serricorne</i>
Clay	Home	Household-Stored Product	Small Hive Beetle Larvae	<i>Aethina tumida</i>

County	Host	Category	Identification	Scientific Name
Elmore	Home	Household-Miscellaneous	Velvet Ant	Mutillidae
Jefferson	Cryptomeria	Ornamental	Box Elder Bug Nymphs	<i>Boisea trivittata</i>
Chilton (2)	Lawn	Household-Miscellaneous	A Spider Wasp	Pompilidae
Chilton	Lawn	Household-Miscellaneous	A Spider Wasp	<i>Entypus unifasciatus</i>
Marion	Building	Household-Miscellaneous	False Chinch Bug	<i>Nysius raphanus</i>
Shelby	Home	Household-Miscellaneous	Giant Resin Bee	<i>Megachile sculpturalis</i>
Limestone		Miscellaneous	Ailanthus Webworm	<i>Attiva punctella</i>
Jackson	Lawn	Household-Miscellaneous	Trap Door Spider	<i>Ummidia</i> sp.

Disease Possibilities For September

Seasonably cooler conditions are more favorable for powdery mildew and downy mildew. Both of these diseases cause yellow blotches on dicot leaves. With powdery mildew, blotches may be more diffuse and a white dusty layer may be visible on the upper and/or lower leaf surfaces. With downy mildew, yellow spots may begin as more definitive angular yellow spots. These spots may merge resulting in large yellow areas. On lower leaf surfaces when weather is wet, humid and temperatures are 60-80°F, a brown-gray-colored webbing may be present on lower leaf surfaces. These diseases are often confirmed in the lab by microscopic observation of characteristic spores.

Evidence of bacterial scorch disease may occur in September. Scorch disease, caused by the bacteria *Xylella*, causes leaf edge scorch and dieback of elm, oaks (red and black oaks including northern red, pin, scarlet, southern red, laurel, shingle, and water oaks), sycamore, mulberry, and red maple. Initial symptoms of scorch may first occur in mid-late June, but disease is often not noticed until late summer or early fall when symptoms are more pronounced. Generally, leaf symptoms progress from older to younger leaves, with leaves at branch tips often showing no symptoms. Scorched leaves curl upward and remain attached. Infected trees develop a

progressive dieback and general (usually slow, over many years) decline. Scorch can be confirmed with an ELISA test. Disease symptoms may be confused with drought or root problems. In August of 2002, this disease was confirmed in a sycamore sample from Barbour County and in a plum sample from Mobile County. Bacterial scorch was diagnosed last year on sycamore in Montgomery County. It was recently found in plum in Chilton County and oak (Red Group) in Marengo County (ELISA tests were positive.)

Many fungal leaf spot diseases will develop on pre-senescent shade tree foliage in September. Generally these spots are of no concern. It is, however, always a good idea to remove fallen spotted foliage from the area later this fall or winter. Stressed trees are more susceptible to these leaf spots.

So far in September, Asian soybean rust has been confirmed in Sumter County on soybean. For more information, check with Ed Sikora or Mary Delaney.

Table 3. Brief Disease Descriptions and Control Recommendations for Diseases Often Seen in August and Late Summer.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Abelia	Cercospora Leaf Spot	Small-large brown, circular leaf spots.	Sanitation.
Alfalfa	Summer Black Stem and Leaf Spot (<i>Cercospora</i>)	Small brown spots become larger (2-6 mm diam.) and reddish brown or smoky brown. During humid conditions, spots become ashy-gray with spores. Lesions occur on stems; small stems and petioles may die from girdling lesions.	Maintain appropriate fertility; harvest frequently.
Ajuga	Cercospora Leaf Spot	Medium brown, circular-irregularly shaped leaf spots of varying sizes.	Sanitation. Cleary's 3336 or Halt may be used.
Almond, Dwarf Flowering	Powdery Mildew	White dusting of fungus on leaf surfaces; later necrosis.	Sanitation; Cleary's 3336, Halt, or OHP 6672.
Althea	Rust (<i>Puccinia</i> or <i>Kuehneola</i>)	Orange, powdery specks on small yellow leaf spots appear.	Sanitation.
Apple	Bitter Rot (<i>Colletotrichum</i>)	Initially small gray or brown spots appear on the fruit. These spots enlarge into medium brown circular lesions. Orange	Sanitation. See the Spray Guide for Fruit Crops.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		spores often develop in concentric rings.	
	Black Rot (<i>Botryosphaeria</i>)	On young fruit, tiny red flecks appear. As fruit matures lesions become large black and irregular sometimes with a red halo. Sometimes alternating rings of brown & black develop. Limb cankers are red-brown, slightly sunken, cracked. Leaf spots are brown with a purple border (4-5 mm diam.)	Sanitation. See the Spray Guide for Fruit Crops.
	Fly Speck (<i>Schizothyrium</i>)	Tiny black dots occur in patches (usually) on the surface (only) of apple fruit.	Sanitation. See the Fruit Spray Guide.
	Sooty Blotch (<i>Gloeodes</i>)	Medium gray spots which resemble sooty smudges appear on surface layer of apple skin only.	Sanitation. See the Fruit Spray Guide.
Arbor-vitae	Armillaria Root Rot	Tree dieback; roots are rotted; check for honey-colored mushrooms, thin, white fungal mats under the bark, and black, thread-like rhizomorphs at crown-soil line area.	Sanitation – tree removal. See ANR-907
Aucuba	Lasiodiplodia Canker	Black sunken spots or sunken areas on aucuba stems. Dieback of foliage results.	Cleary's 3336, or Domain protective spray labeled for ornamentals; sanitation.
Azalea	Armillaria Root Rot	Dieback. At or near the soil level, a white fungal mat seen under the bark. Brown, honey-colored mushrooms may develop at trunk area.	Sanitation of the whole tree.
	Botryosphaeria Canker	Cracked, sunken lesions on branches.	Pruning.
	Cercospora Leaf Spot	Small dark brown-black, usually circular spots (1-2 mm) scattered over leaf	Sanitation. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		surface.	
	Colletotrichum Leaf Spot	Small, round, brown leaf spots.	Sanitation. Protective sprays of Daconil or Cleary's 3336 or Halt could be used.
	Phomopsis Canker/Dieback	Elongated, sunken, elliptical cankers on twigs and branches with resulting dieback.	Sanitation; protective sprays of Cleary's 3336 may help.
	<i>Phytophthora</i> Crown and Root Rot	Lower stem near soil and roots become brown and water-soaked.	Sanitation and proper soil or potting mix drainage are important. See Alabama Pest Management Handbook and ANR-571.
	<i>Rhizoctonia</i> Aerial Blight	Brown, irregular spots and lesions begin on lower leaves. Whole leaves may become blighted; leaf drop occurs.	Sanitation; See AL Pest Management Handbook.
Azalea, Native	Cercospora Leaf Spot	Small, brown, circular spots develop.	Sanitation. Cleary's or Halt may be used.
Bahia Grass	Dollar Spot (<i>Sclerotinia</i>)	Pale, cream-colored silver dollar-sized spots appear in grass area. Individual grass blades show cream colored spots with brown-black borders.	Sanitation. See the AL Pest Management Handbook.
	Helminthosporium-type Leaf Spot	Tiny (2-3 mm), elongated brown spots may cover leaf and stem, stolon surfaces. Severe spotting may cause foliage death.	Maintain appropriate fertility; harvest as frequently as possible.
Bald, Cypress	Cercosporidium Blight	Lower foliage becomes brown. Microscopic study usually shows small spore bodies of Cercosporidium.	Pruning. Cleary's 3336 will provide protective disease control.
Basil	Rhizoctonia Stem & Root Rot	Wilt, dieback, brown, dry lesion on lower stems, crown, and roots.	Sanitations; crop rotation.
Beans, Garden	Anthracoese (<i>Colletotrichum</i>)	Black, sunken cankers on pods which become red-orange when spores are produced. Similar spots are found on stems.	See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		Foliage symptoms involve black, dead portions of veins on the underside of the leaf. Infections of older plants cause damage primarily to pods.	
	Pythium Lower Stem Rot	Lower stems near soil-line show brown, wet rot.	See AL Pest Management Handbook.
	Rhizoctonia Aerial Blight	Leaf blight develops. Leaves become brown and tattered.	Sanitation.
	Rhizoctonia Lower Stem Rot	Lower stems near soil-line have dried brown lesions.	See AL Pest Management Handbook.
	Root-Knot Nematode (<i>Meloidogyne</i>)	Roots develop round-irregularly shaped galls; plants become yellowed; wilt during dry periods.	Rotate to dwarf French marigolds for 1 year or grasses (such as bahia or centipede) for 3-4 years or solarization.
	Uromyces Rust	Reddish-brown powdery pustules on all above ground plant parts, especially lower leaf surfaces.	Protective fungicide sprays. See the AL Pest Management Handbook.
Begonia	Phytophthora Crown Rot	Crown tissues are dark and wet-rotted.	Sanitation. Reduce watering.
	Phytophthora & Pythium Root Rot	Roots become brown and water-soaked, decayed.	Sanitation. Reduce watering. See the AL Pest Management Handbook.
	Rhizoctonia Root Rot	Brown, dry, decayed roots.	Sanitation. Banrot protective drenches.
	Root-knot Nematode (<i>Meloidogyne</i>)	Galls on roots; plants stunted and wilted.	Solarization.
	Tomato Spotted Wilt Virus	Yellow mosaic and ring spots present. Sometimes brown spots also present.	Thrips control. Sanitation.
Bentgrass	Anthracnose (<i>Colletotrichum</i>)	Brown leaf spots, dieback; yellowing.	Sanitation; collect clippings; Cleary's 3336. Check with A. Hagan.
	Bipolaris Leaf Spot	Tiny brown leaf spots that will coalesce to cause large areas of leaves to be	See ANR-621 or the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		blighted.	
	Fairy Ring	Dead rings or half ring patterns in lawn; mushrooms.	See ANR-372.
	Nematode Damage from Ring (<i>Criconeoides</i>) and Sting (<i>Belonolaimus</i>) Nematodes	Thinned, blighted, yellowed turf areas.	See ANR-523.
	Pythium Root Rot	Roots become light brown, wet, and rotted; foliage dies.	See ANR-594 and the AL Pest Management Handbook.
Bermuda, Coastal	<i>Bipolaris</i> (<i>Helminthosporium</i>) Leaf Spot	Tiny (2-3 mm), elongated brown spots may cover leaf and stem, stolon surfaces. Severe spotting may cause foliage death.	Maintain appropriate fertility; harvest as frequently as possible.
Bermudagrass	<i>Bipolaris</i> Leaf/Stem Spot & Blight	See Bermuda, Coastal.	Collect grass clippings; See AL Pest Management Handbook.
	Dollar Spot (<i>Sclerotinia</i>)	Pale, cream-colored, silver dollar-sized spots appear in grass area. Individual grass blades show cream colored spots with brown-black borders.	Sanitation. See the AL Pest Management Handbook.
	Fairy Ring	Large rings or half rings of dead grass with an outer border of dark green turf. When conditions are wet, mushrooms will form in the dark green ring area.	See ANR-372. Also see AL Pest Management Handbook.
	Nematode, Ring (<i>Criconeoides</i> sp.)	Irregular patches become thin, yellow and dieback.	See ANR-523.
	Nematode, Sting (<i>Belonolaimus</i> sp.)	Irregular patches become thin, yellow and dieback.	See ANR-523.
	<i>Rhizoctonia</i> Brown Patch	Symptoms may vary slightly depending upon the situation. Typically, light-medium brown, large, circular patches occur on lawns. Individual grass blades develop small to large brown lesions. Lesions	Collect grass clippings; See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		may involve the whole leaf blade or whole plants.	
	<i>Rhizoctonia</i> Leaf & Sheath Blight (<i>R. zea</i>)	Brown leaf spots; dieback; blight.	Sanitation; collect grass clippings; see ANR-492; See the AL Pest Management Handbook.
	Take-All (<i>Gaeumannomyces</i>)	Spots or areas in turf become yellowed and thinned. Roots become decayed in spots.	See ANR-823. Also, see AL Pest Management Handbook.
Bermuda, Tifdwarf	Bermudagrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	Areas yellow and die out.	See take-all patch recommendations, ANR-823.
Bermuda, Tifeagle	Curvularia Blight	Large leaf areas become blighted. This fungus usually develops as a secondary event.	See AL Pest Management Handbook for brown patch recommended fungicides.
Blackberry	Rust (<i>Gymnoconia</i>)	Lower leaf surfaces are covered with orange powdery masses of spores; witches brooms may develop. Plants stunted; a systemic disease.	Remove infected plants.
	Septoria Leaf Spot	Reddish-brown angular leaf spots.	See the AL Pest Management Handbook.
Blueberry	Phomopsis Canker	Branch dieback to cankers; cankers are elliptical, dry, sunken lesions.	Pruning. Make cuts 3 inches beyond the damage area.
Boxwood	Phytophthora Root Rot	Brown, water-soaked roots.	See AL Pest Management Handbook.
	Volutella Blight	Dieback; canker, small-orange specks that are the fruiting bodies of the fungus.	Sanitation. See the AL Pest Management Handbook.
Boxwood, American	Macrophoma Leaf Spot	Yellow-brown blotches or areas, sometimes with small black specks that are the fungus fruiting bodies.	Correct stress conditions. Cleary's 3336 or Halt may be used.
Cabbage	<i>Alternaria brassicicola</i> Stem Blight	Dark brown-black, oval stem lesions; some lesions have a zonate pattern; dieback.	Sanitation; See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Black Rot (<i>Xanthomonas</i>)	Yellow v-shaped lesions at leaf edges. V-shaped lesions become black; eventually, main stem/stalk become black and soft rotted.	Sanitation; Crop rotation 2 years, see AL Pest Management Handbook.
Cactus, Christmas	Pythium Root Rot	Plant dieback; soft, wet, brown rot at base of stem at soil level.	Remove plant.
Camellia, Sasanqua	Glomerella Canker	Branch dieback; cankers are brown, sunken, sometimes cracked at edges.	Pruning. Make cuts 3 inches beyond the damage.
Celosia	Phytophthora, Pythium, Fusarium Lower Stem Rot & Root Rot	Roots brown and decayed.	Sanitation. Reduce watering. Improve soil drainage.
Centipede	Brown Patch (<i>Rhizoctonia</i>) Dollar Spot (<i>Sclerotinia</i>)	See bermudagrass. Pale, cream-colored, silver dollar-sized spots appear in grass area. Individual grass blades show cream colored spots with brown-black borders.	See Bermudagrass. Sanitation. See the AL Pest Management Handbook.
	Ring & Spiral Nematode Damage (<i>Criconemoides</i> and <i>Rotylenchus</i>)	Patches or areas become yellowed and dieback.	See ANR-523.
Cherry	Septoria Leaf Spot	Small (0.5 cm or less in diam.), angular brown spots.	Sanitation in the fall.
Cherry, Kwanzan	Anthracoise (<i>Colletotrichum</i>)	Brown spots and blotches along leaf veins and along leaf edges.	Sanitation. Cleary's 3336 or Halt may be applied.
Cherry Laurel	Phytophthora Root Rot/Overwatering	Roots become brown, wet, decayed.	Sanitation; correct excess water problem.
Chrysanthemum	Ascochyta Stem Blight	Brown, irregular, sunken, stem cankers; dieback; yellowing.	Sanitation; See the AL Pest Management Handbook under Ascochyta ray blight.
	Fusarium Wilt	Plants develop yellowing & wilt that begins with lower foliage and progresses upward; vascular browning in lower stems.	Remove plants. Crop rotation for several years.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Phytophthora Blossom Blight	Brown, wet blotches and blight on flowers.	Sanitation. Avoid over-head irrigation.
	Phytophthora Root Rot	Roots become brown, wet, decayed.	Sanitation; correct excess water problem. See AL Pest Management Handbook.
	Pythium Root Rot	Wilt, dieback; yellowing of lower leaves; roots become brown and wet rotted.	Sanitation. See the AL Pest Management Handbook.
Clematis	Botrytis Blight	Necrotic blotches. Gray mold may be present when conditions are web.	Sanitation. See the AL Pest Management Handbook.
	Phytophthora Crown & Root Rot	Dieback. Root and lower stem develop wet decay.	Sanitation. Reduce water levels in the area.
Cleyera	Phytophthora Root Rot	See Boxwood.	Sanitation. Improve soil drainage.
Collards	<i>Alternaria</i> Leaf Spot	Gray-black, sooty spots with ring patterns on older leaves; disease of seedlings is severe.	Sanitation. See the AL Pest Management Handbook.
	Black Rot (<i>Xanthomonas</i>)	Dark V-shaped lesion at leaf edge; blackening of leaf veins; black vascular ring if stem is cut cross-wise.	Rotation for 2-3 years; solarization may help.
	Rhizoctonia Wire Stem	The lower stem and major root becomes thin and discolored brown with a dry rot. Plants decline and die.	Remove damaged plants; see the AL Pest Management Handbook.
Coral Bells	Cylindrocladium Root Rot	Brown-black root decay.	Sanitation. Cleary's 3336 or Halt drenches for protective action.
	Pythium Root Rot	Roots become light brown, decayed, and water-soaked.	Sanitation. Reduce water levels in the area.
Coneflower	Phytophthora Root Rot	Dieback. Roots become decayed with a brown, wet rot.	Sanitation. Reduce water levels.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Corn	Charcoal Rot (<i>Macrophomina</i>)	Gray lesions develop on lower stems and roots. Under the epidermis, numerous resting structures and fruiting bodies are scattered throughout the stem and root tissues which become dried and separate easily. These tissues appear as though they were sprinkled with finely-ground pieces of charcoal.	Maintain healthy plants with proper fertilization and watering if possible, crop rotation.
Cotoneaster	Phyllosticta Leaf Spot	Round, cream-colored leaf spots with dark borders. When severe, leaf drop.	Sanitation; See AL Pest Management Handbook.
	Phytophthora Root Rot	Dieback; lower foliage dieback and yellowing; roots become brown and wet rotted.	Sanitation – remove damaged plants; remove root associated soil; reduce irrigation; improve drainage.
	Rhizoctonia Aerial Blight	Lower and inner foliage becomes blighted/brown.	Sanitation of damaged foliage. Use protective sprays of Cleary's 3336.
Cotton	Alternaria Leaf Spot	Irregular, or circular, slightly zonate brown leaf spots/blotches.	See Bill Gazaway.
	Ascochyta Leaf Spot	Light brown, mostly circular, small to larger spots.	See Bill Gazaway.
	Botryodiplodia Pod Decay	Pods become black and decayed.	Sanitation.
	Cercospora Leaf Spot	Irregularly shaped brown leaf spots develop.	See Bill Gazaway.
	Nematode, Reniform (<i>Rotylenchulus</i>)	Irregularly shaped brown leaf spots develop.	See Bill Gazaway.
	Nematode, Root-knot (<i>Meloidogyne</i>)	Plants grow poorly; roots have galls.	Crop rotation. See Bill Gazaway.
Crape Myrtle	Cercospora Leaf Spot	Irregular shaped brown leaf spots develop.	Sanitation. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Cucumber	Cercospora Leaf Spot	Irregular, light brown spots.	Sanitation. See the AL Pest Management Handbook.
	Downy Mildew	Yellow, irregular spots that become dark. Gray mold on lower leaf surface may develop.	Sanitation. See the AL Pest Management Handbook.
Cypress, Leyland	Botryosphaeria Canker.	Elongated, sunken, cracked lesions on branches.	Pruning.
	<i>Cercosporidium</i> Blight (formerly <i>Cercospora</i>)	Needle and twig blight that usually begins on lower foliage.	Sanitation. Cleary's 3336 protective sprays.
	Phytophthora Crown Rot	Brown, wet lower trunk decay.	Sanitation. See AL Pest Management Handbook.
	Pythium Feeder Root Rot	Light brown feeder root decay.	Sanitation. See AL Pest Management Handbook.
	Seiridium Canker	Elongated, sunken lesions with oozing sap.	Sanitation; pruning; protective sprays of Cleary's 3336.
Daylily	Anthracnose (<i>Colletotrichum</i>)	Circular, brown leaf spots.	See the AL Pest Management Handbook.
	Daylily Rust (<i>Puccinia hemerocallidis</i>)	Leaves develop small orange, powdery spots and affected leaf areas become yellow, then brown.	Sanitation. Protective fungicide treatments of Banner Maxx or Heritage are available for commercial situations. Spectracide Immunox or Fertiloam Sysem Fungicide may be used in homeowner situations.
	Leaf Streak (<i>Kabatiella</i> sp.)	Elongated brown/yellow streaks on leaves.	Sanitation. See the AL Pest Management Handbook.
	Phytophthora Root Rot	Wilt, yellowing of lower foliage; roots become brown and wet rotted.	Sanitation. Remove damaged plants and root associated soil. Reduce irrigation or improve soil drainage. Crop rotate.
	Pythium Root Rot	Wilt, yellowing of lower foliage; roots become light brown & wet rotted.	Sanitation. Remove damaged plants. Reduce irrigation. Improve soil

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			drainage.
Dianthus	Phytophthora Stem Blight	Brown, water-soaked cankers.	Sanitation. Reduce water levels.
	Pythium Root Rot	Slightly brown, water-soaked root decay.	Sanitation. Reduce water levels. Subdue may be used.
Dogwood	Botryosphaeria Canker	Slightly sunken lesion, sometimes with cracks along the margin.	Sanitation.
	Cercospora Leaf Spot	Angular-irregular tan-brown lesions (2-6 mm diam.) sometimes with a thin yellow halo.	Usually sanitation is the only control measure needed.
	Powdery Mildew (<i>Oidium</i> ; <i>Microsphaera</i> or <i>Phyllactinia</i>)	White, powdery patches on leaves; affected areas become blighted.	See AL Pest Management Handbook. Sanitation.
	Septoria Leaf Spot	Angular, brown spots, about 1 cm or less in diam; may be confused with Cercospora leaf spot.	Collect and remove fallen leaves this fall.
Dusty Miller	Alternaria Leaf Spot	Dark, angular spots	Sanitation; Cleary's 3336.
Eleagnus	Phytophthora Root Rot	Brown, wet root decay.	Sanitation. Improve soil drainage.
Euonymus	Anthracoese (<i>Colletotrichum</i>)	Small brown spots (about 5 mm or smaller) on foliage.	Sanitation; See the AL Pest Management Handbook for protective fungicide recommendations.
Euonymus, Creeping	Rhizoctonia Root Rot	Plants dieback; yellowing of foliage; roots become dry rotted.	Remove plants and root associated soil.
Fatsia	Colletotrichum Blotch	Brown leaf spots & blotches.	Sanitation of fallen leaves. Cleary's 3336 or Halt would provide protective disease control.
Fern	Anthracoese (<i>Colletotrichum</i>)	Gray-brown irregular blotches on fronds. Orange spore masses may be present in humid weather.	Sanitation; See the AL Pest Management Handbook under leaf spot.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Rhizoctonia Aerial Blight	Gray or brown irregular blotches on fronds; some 'shot-hole'.	Sanitation; See the AL Pest Management Handbook.
Fescue	Brown Patch (<i>Rhizoctonia</i>)	See bermudagrass.	See bermudagrass.
	Helminthosporium Leaf Spot	Tiny, elongate brown leaf spots that may be numerous, coalesce and cause blight of entire leaf blade.	See ANR-621 or AL Pest Management Handbook.
Fig	Cercospora Leaf Spot	Gray-brown irregular spots, blotches.	Sanitation of leaves in the fall.
Forsythia	Phytophthora Root Rot	Dieback. Lower foliage yellowing and dieback first. Roots become brown discolored and wet rotted.	Sanitation. Reduce water levels. Protective drenches of Subdue where appropriate.
Gardenia	Phytophthora Crown Rot	Lower stem/trunk at the soil line develops wet decay.	Sanitation. (See the AL Pest Management Handbook under Root Rot for protective treatment.)
Grancy Gray Beard	Algal Leaf Spot (<i>Cephaleuros</i>)	Green-reddish, slightly raised spots with wavy edges.	Sanitation.
Grape	Cercospora Leaf Spot	Angular medium brown leaf spots.	Recommendations for anthracnose should help.
Hickory	Scab (<i>Cladosporium</i>)	Small, dark brown, slightly raised leaf spots.	Sanitation in the fall.
Holly, Blue Maid	Botryosphaeria Canker	Brown or black sunken, cracked lesions (cankers) on branches.	Sanitation. Protective sprays of Cleary's 3336, Domain or a WP benomyl labelled for ornamentals.
	Phytophthora Root Rot	Feeder roots become water-soaked, decayed.	See the AL Pest Management Handbook.
Holly, Helli	Rhizoctonia Aerial Blight	Lower foliage becomes blighted.	Sanitation. See the AL Pest Management Handbook.
Holly, Japanese	Black Root Rot (<i>Thielaviopsis</i>)	Dieback; lower foliage often shows yellowing & dieback first. Roots develop black lesions and black tips.	Sanitation. Protective drenches of Cleary's 3336. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Pythium Root Rot	Foliage becomes yellowed on lower branches. Roots become slightly discolored and rotted.	Sanitation. Reduce soil/media water levels. See the AL Pest Management Handbook.
Honeylocust	Crcospora Leaf Spot	Round, brown leaf spots.	Sanitation.
Hosta	Anthracnose	Brown, circular-oval leaf spots; may show some zonation.	Sanitation; Cleary's 3336 will provide some control.
	Southern Blight (<i>Sclerotium rolfsii</i>)	Lower stems become rotted. A white mold may develop. Small, brown or black mustard seed sized overwintering bodies may appear.	Sanitation. (Removal of soil in root zone, if a small area.)
Hydrangea	Armillaria Root Rot	Sudden dieback; white, thin fungal layer may be present under bark; black thread-like structures may be present over or under bark near roots; honey-colored mushrooms may be present near base of shrub.	Sanitation.
	Cercospora Leaf Spot	Relatively large (0.5-1.0 cm) dark brown circular spots with reddish borders.	Sanitation. See the AL Pest Management Handbook.
	Corynespora Leaf Spot	Brown, somewhat circular spots; sometimes spots have zonate pattern.	Sanitation.
	Phytophthora & Pythium Root Rot	Roots become brown and water-soaked.	Sanitation. See the AL Pest Management Handbook.
	Powdery Mildew	White dusting on leaves. Necrosis follows.	See the AL Pest Management Handbook.
Hypericum	Rust (<i>Uromyces</i>)	Yellow leaf spots; brown powdery pustules on lower leaf surfaces.	Sanitation of infected plant parts.
Impatiens	Alternaria Leaf Spot	Small circular or angular dark brown spots.	Sanitation; a mancozeb product such as Duosan or Zyban.
Indian Hawthorn	Cercospora Leaf Spot	Angular, vein-bound brown-red spots.	Sanitation. Cleary's 3336 may be used.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Entomosporium Leaf Spot	Red-black, circular-irregular spots.	Sanitation. See the AL Pest Management Handbook.
Iris	Bacterial Soft Rot	Soft, wet, watery rot of rhizome.	Sanitation. Control insect problems.
	Fusarium Rhizome Rot	Areas of the rhizome exhibit a dry, brown rot.	Sanitation. See the AL Pest Management Handbook.
Irish Potato	Rhizoctonia Black Scurf	Small, black, crusty bodies on tuber surface.	See the AL Pest Management Handbook.
Ivy, English	Alternaria Leaf Spot	Round or oval, brown leaf spots.	Sanitation. Protect T/O or other labeled mancozeb product.
	Anthracnose (<i>Colletotrichum</i>)	Circular or irregularly-shaped brown leaf spots develop.	Sanitation. See the AL Pest Management Handbook.
	Cercospora Leaf Spot	Brown, irregular leaf spot.	Sanitation. Cleary's 3336, Halt, or OHP 6672 may be used.
	Phytophthora Crown & Root Rot	Tissues dark and water-soaked.	Sanitation; reduce irregular or improve drainage.
	Phytophthora Leaf & Stem Rot	Dark, water-soaked, irregular lesions that become dry.	Sanitation; avoid overhead irrigation; Heritage or Protect T/O.
	Pythium Root Rot	Light brown rotted roots.	See Phytophthora Root Rot.
Juniper	Cercospora (formerly Asperisporium and Cercospora) Blight	Blight of needles beginning with lower foliage.	Sanitation. Cleary's 3336 protective sprays.
	Pestalotia Needle Blight	Lower foliage needle blight associated with plant stress.	Sanitation.
	Phomopsis Dieback	Juniper branch tips become brown. Cankers develop on twigs and dieback continues down the twig.	See the AL Pest Management Handbook.
	Phytophthora Root Rot	See Holly.	See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Kudzu	Bacterial Leaf Spot (<i>Pseudomonas</i> sp.)	Circular, dark spots with a yellow halo.	---
	Cercospora Leaf Spot	Brown, irregular leaf spots.	---
Leucothoe 'Drooping Rainbow'	Cercospora Leaf Spot	Brown circular to irregular spots.	Sanitation; Cleary's 3336 or Halt.
Ligustrum	Cercospora Leaf Spot	Brown irregular spots (about 1 cm diam.) on foliage; when leaf spot is severe, defoliation may result.	Sanitation; See the AL Pest Management Handbook.
Lilac	Phyllosticta Leaf Spot	Brown or light brown leaf spots with dark margins.	Sanitation. Cleary's 3336 or Halt would provide protective disease control.
Liriope	Anthracnose (<i>Colletotrichum</i>)	Brown irregular blotches on leaf blades; often, leaf tip areas are involved.	Sanitation; See the AL Pest Management Handbook.
London Plane Tree	Bacterial Leaf Scorch	Inner leaves first develop a leaf scorch. Gradually all foliage become scorched, dieback follows.	Remove infected trees.
Loquat	Botryodiplodia Canker	Sunken, dried lesions with cracked edges.	Sanitation. Make cuts 3 inches beyond the damage.
Loripetalum	Pythium Root Rot	Light brown, water-soaked, rotted roots.	Sanitation. Reduce irrigation or improve soil drainage.
Magnolia	Black Mildew	Black mold develops on lower leaf surfaces.	Reduce high humidity levels by increasing air circulation; pruning suggested.
Magnolia, Saucer	Phytophthora Leaf Spots	Brown, water-soaked spots develop.	Sanitation; for small trees, do not irrigate over-head.
Maple	Anthracnose (<i>Kabatiella</i>)	Brown spots and blotches on foliage; enlarged spots may involve more than half of individual leaves.	Sanitation. See the AL Pest Management Handbook.
	Phyllosticta Leaf Spot	Gray circular spots (¼ inch diam. Approx.) with dark brown or reddish brown borders.	Sanitation. See the AL Pest Management Handbook.
Maple, Red	Botryosphaeria Canker	Elongated, sunken, often	Sanitation. Remove stress

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		cracked lesions.	factors.
Marigold	Alternaria Leaf Spot	Small (0.2-0.3 cm diam.) dark brown-black spots. Numerous spots cause death of plants.	Sanitation. See the AL Pest Management Handbook.
	Phytophthora Crown Rot	Crowns become brown, decayed, water-soaked.	Sanitation. See the AL Pest Management Handbook.
	Pythium Crown Rot	Crowns become brown, decayed, water-soaked.	Sanitation. See the AL Pest Management Handbook.
	Rhizoctonia Crown Rot	Tissues become brown and dry rotted.	Sanitation. Banrot may be used as a protective treatment.
Millet	<i>Piricularia</i> Leaf Spot	Irregular, 3-6 mm diameter gray-brown leaf spots.	Crop rotation.
Mondograss	Anthracnose	Light brown blotches, leaf spots on leaves; often spots have dark brown borders.	Cut foliage back. Cleary's 3336 will help.
Muscadine	Anthracnose (<i>Colletotrichum</i>)	Circular or angular, brown lesions on leaves or stems, which may coalesce. Centers of lesions may become gray-white. Lesion borders are dark purple-brown-black. Cracking may occur. Similar-looking lesions may occur on fruit. Fruit lesions extend into fruit pulp.	Sanitation. See AL Pest Management Handbook.
	Black Rot (<i>Guignardia</i>)	Brown circular spots with dark brown borders on foliage and fruit. Spots may coalesce to involve large area of tissue.	See the AL Pest Management Handbook.
Mustard	Cercospora Leaf Spot	Irregularly-shaped brown leaf spots.	Sanitation.
Myrtle, Wax	Botryosphaeria Canker	Sunken, cracked lesions on stems.	Pruning 3-4 inches from edge of decay.
Nandina	Cucumber Mosaic Virus	Plants show stunted new growth; some mosaic, leaf distortion, mottle and/or	Sanitation; aphid control may help a small amount.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		curling/puckering may be present.	
Nectarine	Brown Rot	Brown, soft rot sometimes with gray spore masses.	Sanitation. See AL Pest Management Handbook under 'peach'.
	Phomopsis Canker	Brown, sunken, dry decay lesions on twigs and branches.	Sanitation. See AL Pest Management Handbook under 'peach'.
Oak	Bacterial Scorch (<i>Xylella</i>)	Inner leaves first develop a leaf edge scorch. Gradually all foliage becomes scorched; dieback follows.	Removal of infected trees.
	Hypoxylon Canker	Bark cracking and sloughing off; gray or black hard stroma layer develops under bark layer.	Sanitation.
	Oak Leaf Blister (<i>Taphrina</i>)	Brown puckered spots.	Sanitation.
	Powdery Mildew (<i>Microsphaera</i>)	White dusty coating on upper leaf surfaces. Some distortion of new leaves.	Collect and remove fallen leaves this fall.
Oak, Black	Hypoxylon Canker	Dark brown or gray, hard, flat, fungal bodies form under the bark; bark cracks and fall off.	Sanitation – pruning.
Oak, Pin	Xylella Scorch	Lower and oldest leaves show leaf edge scorch; problem progresses upward through the tree canopy. Dieback develops; eventual tree death.	Remove dying trees.
Oak, Red	Tubakia Leaf Spot	Round or irregular-round spots or blotches.	Sanitation of fallen leaves in the fall.
Oak, Sawtooth	Tubakia Leaf Spot (Formerly <i>Actinopelte</i>)	Circular, brown leaf spots may have tiny black specks scattered on leaf spot surface.	Sanitation of fallen leaves in fall.
Oak, Shumard	Bacterial Leaf Scorch (<i>Xylella</i>)	Inner leaves develop leaf scorch; gradually all foliage becomes scorched; dieback follows.	Remove infected trees.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Hypoxyylon Canker	Dark brown or gray, hard, flat, fungal bodies form under the bark; bark cracks and fall off.	Sanitation – pruning.
Okra	Fusarium Wilt	Plant wilts, beginning with lower foliage. Vascular system at lower stems is brown or brown streaked.	Sanitation. Do not plant okra for 10 or more years.
	Nematode, Root Knot (<i>Meloidogyne</i> sp.)	Poor growth; plants are stunted; roots have small-large galls.	Sanitation; See ANR-523.
Pansy	Anthrachnose (<i>Colletotrichum</i>)	Small, round, light brown, cream-colored spots.	Sanitation. Cleary’s 3336.
	Myrothecium Crown Rot	Dieback; decayed crowns.	Sanitation; See A. Hagan.
	<i>Phyllosticta</i> Leaf Spot	Relatively small (2-3 mm diam.) medium brown, roughly circular spots. Spot centers may become gray.	Sanitation. Protective sprays of Cleary’s 3336, Domain, or a benomyl WP labeled for ornamentals.
	Phytophthora & Pythium Root Rot	Brown, wet-rotted roots.	Sanitation. See AL Pest Management Handbook.
	Phytophthora Root Rot	Roots become brown and water-soaked.	Sanitation. See the AL Pest Management Handbook.
	Pythium Root Rot	Roots become brown and water-soaked.	Sanitation; See the AL Pest Management Handbook.
	Rhizoctonia Brown Rot	Lower stem develops a brown, dry rot.	Sanitation. Cleary’s 3336 or chlorothalonil products.
	Thielaviopsis Root Rot	Roots become covered with black spots/lesions.	Sanitation; Cleary’s, or Domain protective drenches.
Peace Lily	Pythium Root Rot	Lower foliage turns yellow first; plants wilt; roots become light brown and wet rotted.	Remove plants. Remove root associated soil; reduce irrigation; improve soil water drainage.
Peanut	Cylindrocladium Black Rot	Lower stems develop a black rot decay.	See A. Hagan.
	Diplodia Collard Rot	Wilt; plant collapse and death; brown lesions with dark brown margins at	See A. Hagan.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		lower stems/collar; roots become gray-black and shred.	
	Early Leaf Spot (<i>Cercospora</i>)	Brown spots with halos develop on lower leaf surfaces; spore production usually on upper leaf surface.	Protective fungicide sprays. See Peanut Spray Guide and Timely Information PP-350 on Folicur.
	Late Leaf Spot (<i>Cercosporidium</i>)	Brown to black spots, sometimes difficult to distinguish from early leaf spot unless spores are observed microscopically; spore production usually on lower leaf surface.	Protective fungicide sprays. See Peanut Spray Guide and PP-350.
	Lesion Nematode Pod Damage (<i>Pratylenchus</i>)	Pods shriveled, sunken lesions.	See A. Hagan.
	Pepper Spot (<i>Leptosphaerullina</i>)	Tiny, black spots scattered on upper leaf surfaces; another symptom is a brown wedge-shaped lesion at upper surface leaf tip area; a yellow halo is usually present.	See A. Hagan.
	Rust (<i>Puccinia</i>)	Orange pustules on foliage.	See A. Hagan.
	Peanut Mottle Virus	Plants stunted with regular green, light green color pattern.	See A. Hagan.
	<i>Rhizoctonia</i> Limb Rot	Oval to elliptical, brown spots develop on stems. Young spots may have a target ring pattern. Whole limbs or stems become blighted.	See AL Pest Management Handbook and PP-350; Careful water management.
	<i>Rhizoctonia</i> Pod Rot	Dull, light or dark brown, sunken lesions. Dark brown fungal threads (hyphae) may be seen on seed surface and inside wall of the shell.	See <i>Rhizoctonia</i> Limb Rot.
	Root-Knot Nematode (<i>Meloidogyne</i>)	Plants are stunted and grow poorly; galls develop on roots.	Crop rotation. See Timely Information, Nematode Suppressive Crops, PP-

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			341.
	Southern Blight (White Mold)	Soft decay spots on stems near the soil usually become covered with white mold that sometimes contains tiny black spherical fungal bodies.	See the AL Pest Management Handbook.
	Tomato Spotted Wilt Virus	Stunted plants; leaves show ring spot and mosaic patterns; new leaves small with abnormally shortened internodes.	Control thrips and weeds; Sanitation.
Pear, Apple	Bitter Rot (<i>Colletotrichum</i>)	Brown, circular spots develop on the fruit surface. Orange dots of spore masses in a circular pattern develop on the surface of the discolored fruit skin. A brown rot extends through the fruit in a v-shaped area.	Sanitation. See AL Pest Management Handbook for fungicide sprays. Follow recommendations for Black Rot.
	Black Rot (<i>Botryosphaeria obtuse</i>)	Brown, circular lesions (4-5 mm diam.) with purple borders on leaves; leaf yellowing and drop may occur; surface lesions on fruit show black-brown concentric rings which remain firm. Branch cankers may be small, long (5 m), sunken and/or cracked, superficial or deep.	Sanitation. See AL Pest Management Handbook for fungicide sprays.
Pear, Bradford	Alternaria Leaf Spot	Oval or round, brown leaf spots.	Sanitation of leaves in the fall. Protect T/O will help provide protective disease control.
	Fabraea Leaf Spot	Black circular spots (about 0.2-0.4 cm diam.)	Sanitation of fallen leaves in the fall. Regular spray schedule may help. See AL Pest Management Handbook.
Peas, Southern	<i>Fusarium</i> Wilt	Vascular tissues of lower stem are discolored reddish-brown.	Rotate peas out of the area for 10+ years.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Mosaic Virus	Leaves show a yellow-green mosaic color pattern; sometimes green bands occur along the veins; plants are stunted.	Sanitation; control insects. Use resistant varieties such as Corona, Pinkeye Purplehull-BVR, Texas Pinkeye, Genegreen, Grant Blackeye or Royal Blackeye.
Pecan	Scab (<i>Cladosporium</i>)	Small, circular, olive-green or black, slightly raised spots develop on leaves, petioles, and nut shuck tissue. Lesions may coalesce causing terminals to die.	See the Pecan Spray Guide; Sanitation.
	Zonate Leaf Spot (<i>Cristulariella</i>)	On upper leaf surfaces, gray-brown concentric-ring spots (up to 2 cm diam.) appear. On lower leaf surfaces, spots are paler brown with dark brown borders.	Sanitation; protective fungicide sprays. See the Pecan Spray Guide.
Peony	Botrytis Leaf Spot	Irregularly-shaped brown-gray blotches.	Sanitation. Pruning to increase air circulation. Cleary's 3336, Halt.
Pepper	Bacterial Blight	Leaves develop dark brown angular spots which are often water-soaked along the edges. Leaf yellowing and drop often occurs on spotted leaves.	See AL Pest Management Handbook.
	Fusarium Wilt (<i>Fusarium oxysporum</i>)	Plant wilts, beginning with the lower foliage. Vascular browning present in the lower stem.	Rotate away from pepper and other solanaceous plants.
Pepper & Other Vegetables	Southern Blight (<i>Sclerotium rolfsii</i>)	Initially a dark brown lesion forms on the stem just below the soil surface. Plants wilt and turn yellow. The lower stem rot may also become a root rot. Coarse white fungal threads develop at the soil line around the stem. Eventually small tan, spherical fungal mustard seeds develop around and in bodies (resembling the	Use Terraclor 75WP on pepper. See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		coarse white fungal threads).	
Petunia	Myrothecium Crown Rot	Crowns become decayed, brown and soft.	Sanitation. Protective sprays of Daconil may be used.
	Phytophthora Aerial Blight	Stems & leaves develop brown, water-soaked decay.	Sanitation. Reducing water levels. See the AL Pest Management Handbook.
	Phytophthora Crown & Root Rot	Crowns & roots become decayed and water-soaked.	Sanitation. See the AL Pest Management Handbook.
	Pythium Crown Rot	Crown water-soaked and decayed.	Sanitation. Reduce water in the area. See the AL Pest Management Handbook.
Photinia	<i>Armillaria</i> Root Rot	Plant may decline slowly or suddenly; lower trunk under the bark and roots may be covered with closely appressed white fungal mat with black, threadlike structures.	Sanitation. See ANR-907.
	<i>Phytophthora</i> Root Rot	See Azalea.	--
Pine, Virginia	Lophodermium (<i>Ploioderma</i>) Needle Cast	Older needles turn brown and drop; very small (1-2 mm or 1/32 inch) football shaped, black fruiting bodies develop on brown needles.	Protective fungicides spray. See AL Pest Management Handbook.
	Rhizosphaeria Needle Cast	Needles become gray-brown. Twig blight may develop.	Sanitation. See the AL Pest Management Handbook.
Pittosporum	Southern Blight (<i>Sclerotium rolfsii</i>)	Lower trunk becomes rotted and softened. White mycelial mats and tiny black spherical bodies (sclerotia) may be present on trunk at soil surface.	Sanitation.
Plum	Black Knot (<i>Plowrightia morbosum</i>)	Branches exhibit elongated black-surfaced, irregular galls that may involve a long (10 cm or more) distance of the branch.	Sanitation of galls. See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Poinsettia	Bacterial (<i>Erwinia</i>) Stem Rot	Black, water-soaked spots or lesions on stems. Lesions may girdle stems.	Sanitation; pot-level irrigation; See AL Pest Management Handbook.
	Pythium Root Rot	Roots become medium brown, soft, water-soaked and rotted.	See AL Pest Management Handbook.
	<i>Rhizoctonia</i> Stem Rot & Root Rot	Lower stems develop dry medium-dark brown surface lesions; roots may become brown and dried.	See AL Pest Management Handbook.
	<i>Rhizopus</i> Stem Rot	Stem sections become glassy and water-soaked; a delicate black mass of fungal threads and small black spherical structures may develop over the lesions.	Sanitation.
Pumpkin	Downy Mildew (<i>Pseudoperonospora</i>)	Yellow spots/blotches develop on upper leaf surfaces; gray spots appear on corresponding areas of lower leaf areas. When temperature are cool-moderate and humid, a gray mycelium/spore layer will develop on lower leaf surface spots.	See the AL Pest Management Handbook.
	Mosaic Virus	Leaves develop a yellow mosaic pattern on dark green background. New growth is stunted.	Sanitation. Control insects. Do not save seed.
	Plectosporium Blight	Cream-colored, raised, scabby lesions on stems, leaves, fruit, and peduncles.	Sanitation. Check with Ed Sikora.
	Root Knot Nematode (<i>Meloidogyne</i>)	Plants become stunted. Roots are galled.	Sanitation. Solarization. See ANR-713.
Raspberry	Septoria Leaf Spot	See Blackberry, under leaf spot.	See Blackberry, under leaf spot.
Red Cedar	Phomopsis Tip Blight	Tips of twigs become yellowed and browned. Dieback may spread down the twig-branch. Lower foliage is affected first.	Sanitation. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Rhododendron	Cercospora Leaf Spot	Brown, round leaf spots.	Sanitation. Cleary's 3336 or Halt may be used for protective control.
	Phytophthora Root Rot	Dieback; roots become brown, water-soaked initially then dried, decayed.	Sanitation; correct excess water problem; see AL Pest Management Handbook under azalea.
	Rust (<i>Puccinastrum</i>)	Golden brown spore pustules on lower surfaces of leaves is followed by leaf blight.	Remove hemlock from the area. Remove infected rhododendron plant parts. See the AL Pest Management Handbook.
Rose	Armillaria Root Rot	Roots become dry rotted. Honey-colored mushrooms may develop. A thin white mold may develop under the bark.	Sanitation. See ANR-907.
	Phytophthora Root Rot	Dieback; active infections are wet-rotted; old infections are dried.	Sanitation; reduce watering.
Rosemary	Phytophthora Root Rot	Roots become brown, decayed, water-soaked.	Sanitation. Reduce watering.
	Rhizoctonia Web Blight	Lower foliage becomes blighted.	Sanitation. Reduce watering.
Salvia	Pythium Root Rot	Foliage turns yellow, yellowing on lower foliage first; roots become light brown and rotted.	Sanitation. Reduce soil water levels.
	Rhizoctonia Crown Rot	Crowns become brown & dry-rotted.	Sanitation. Cleary's 3336 or Halt may be used.
Satsuma	Greasy Spot (<i>Mycosphaerella-Cercospora</i>)	A yellow mottle develops on upper leaf surface with a yellow-brown blister on the lower leaf surface. Later spots become brown-black and greasy looking. Leaf drop follows.	Sanitation; copper protective sprays.
Schip Laurel	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Angular brown spots with water-soaked margins; shot-holes develop.	Sanitation.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Scuppernong	Black Rot (<i>Guignardia</i>)	Vines & fruit develop black decay/rot. Leaves develop brown, irregular spots with circles of small black specks.	Sanitation. See AL Pest Management Handbook under grapes.
Sequoia, Giant	Cercosporidium Needle Blight	Lower foliage become brown.	Pruning off dying branch areas. Cleary's 3336 or Halt may be applied.
Snapdragon	Cercospora Leaf Spot	Pale brown angular leaf spots of variable size.	Cleary's 3336, Domain, or a benomyl labeled for ornamentals.
Sorghum, Grain	Anthracnose {Red Rot} (<i>Colletotrichum</i>)	Red spots and lesions on leaves and stalks. Heads may become infected, reddish and rotted.	Sanitation.
	<i>Fusarium</i> Head Blight	Entire seed head may rot and become covered by cream-pink fungal spore masses.	Sanitation.
	Gloeocercospora Blight (Zonate Leaf Spot)	Zonate spots on leaves; red-purple bands alternate with yellow bands.	Sanitation.
Soybean	Aerial Blight (<i>Rhizoctonia</i>)	Small spots or large areas of leaves, stems, or pods may become brown and blighted; leaves may become tattered; leaf drop; disease favored by high humidity.	See Soybean Spray Guide.
	Anthracnose (<i>Colletotrichum</i>)	Large, irregular, brown areas on stems, pods, petioles.	Deep plow.
	Asian Soybean Rust (<i>Phakopsora pachyrhizi</i>)	Leaf spots, leaf yellowing, leaf drop, reduced yield.	See Ed Sikora.
	Bean Pod Mottle Virus	Most noticeable as abnormally green stems at harvest time; leaf mottle, reduced yield and reduced seed quality possible.	See Ed Sikora.
	Brown Spot (<i>Septoria</i>)	Irregular dark brown spots (1-4 mm diam.) on upper and lower leaf surfaces.	See Soybean Spray Guide.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Cercospora Leaf Spot	Usually spots begin on lower leaves first. Leaf spots are circular, light brown with dark brown edges often called frog eye leaf spot.	---
	Charcoal Rot (<i>Macrophomina</i>)	A light grey discoloration of tap root and lower stem. Inner stem tissues appear shredded and gray as if sprinkled with finely powdered charcoal. A disease of hot, dry conditions.	Rotation; proper fertilization.
	Pod and Stem Blight (<i>Diaporthe, Phomopsis</i>)	Stems, petioles, pods, seeds become just slightly discolored. During wet, warm conditions, linearly arranged black dots (fruiting bodies) appear on infected tissues.	See Soybean Spray Guide.
	Root-Knot Nematode (<i>Meloidogyne</i>)	Irregularly-shaped galls appear on roots. Plants grow poorly.	Rotation; Fumigation; See Soybean Spray Guide.
	Stem Canker (<i>Diaporthe</i>)	Reddish brown, slightly sunken cankers that girdle stems and kill plants.	See the Soybean Control Recommendations ANR-413.
	Sudden Death Syndrome (<i>Fusarium solani</i>)	Leaves become yellowed and then browned in interveinal areas; browned leaves fall from plants; the tap roots and often lateral roots become browned and decayed. Rotting may extend to the crown area at the soil line.	Sanitation and deep plowing of plant residue; rotation; call Ed Sikora if more information is needed.
	Target Leaf Spot (<i>Corynespora cassiicola</i>)	Oval or circular, slightly zonate brown leaf spots.	See Ed Sikora.
Squash, Summer	Cercospora Leaf Spot	Irregular medium-brown spots (0.2-0.6 cm diam., usually).	See the AL Pest Management Handbook.
	Cucumber Mosaic Virus	Plants develop mosaic, stunting and abnormal shoestring leaves.	Sanitation. Aphid control may help a little.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Papaya Ringspot Virus	Plants develop mosaic and stunting.	Sanitation. Aphid control may help a little.
	Watermelon Mosaic Virus II	Pronounced mosaics of green and yellow.	Sanitation. Aphid control may help a little. Virus may be transmitted mechanically. Control weeds. WMV is not seed transmitted.
	Zucchini Yellow Mosaic Virus	Foliage and fruit develops mosaic and may be stunted.	Sanitation. Aphid control may help a little.
St. Augustine	Brown Patch (<i>Rhizoctonia</i>)	See Bermudagrass.	--
	Grey Leaf Spot (<i>Piricularia</i>)	Small spots usually develop into large 4-8 mm diam.), brown or gray lesions with purple or brown borders. A yellow halo or general chlorosis may develop around spots. When severe entire foliage may turn gray-brown.	Collect clippings; See AL Pest Management Handbook.
	Take-All Patch (<i>Gaeumannomyces</i>)	Spots/areas of turf become thinned and yellowed. Eventually plants die and the problem area becomes larger.	See Timely Information PP-312.
St. John's Wart	Pythium Root Decay	Roots are slightly brown, water-soaked, rotted; tissues pull apart easily.	Sanitation; correct excess water problem.
Strawberry	Anthrachnose (<i>Colletotrichum</i>)	Lesions on stolons are brown-black, longitudinal and sunken. During high humidity cream-pink spore masses may form on lesions. Crown rot appears as reddish-brown firm rot or streaks of rot; plants wilt and die when crown rot is severe.	Protective fungicide sprays; Sanitation.
	Cylindrocladium Petiole & Crown Rot	Petioles & lower stems become brown & dry rotted; lower foliage become yellowed and yellowing/dieback spread upward.	--

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Phomopsis Leaf Blight	Brown spots/blotches often associated with leaf edges.	See the AL Pest Management Handbook.
	Phytophthora Crown Rot	Lower foliage yellowing first, then whole plant collapse. Lower stem at soil line develops internal red-brown wet, decay.	Sanitation. See the AL Pest Management Handbook.
	Powdery Mildew	White, powdery patches on foliage which eventually become irregular, dead areas.	Sanitation. See the AL Pest Management Handbook for commercial crop or ANR-50 for homeowner.
Sunflower	Alternaria Leaf Blight	Brown-gray oval spots and cankers.	Sanitation.
Sweet Potato	Fusarium Surface Rot on Roots	Lesions are circular, light-dark brown, firm, and dry; rot does not extend beyond vascular ring. In storage lesions eventually become shrunken, cracked; roots dry out.	Avoid wounding.
	Scurf (<i>Monilochaetes</i>)	A brown patchy discoloration of root which usually begins as small, brown specks or spots. The discoloration is entirely superficial, but cracks will cause roots to dry out.	See AL Pest Management Handbook.
Sycamore	Bacterial Scorch (<i>Xylella fastidiosa</i>)	Older leaves on branches develop leaf edge scorch. Gradually branches will dieback.	Remove the damaged tree.
Thrift	Rhizoctonia Blight	Stem and leaf browning.	Sanitation; Cleary's 3336.
Tomato	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Dark, water-soaked, irregular and somewhat circular leaf spots (usually 1-3 mm diameter).	Sanitation. See the AL Pest Management Handbook.
	Bacterial (Pith Necrosis) Canker (<i>Pseudomonas</i>)	Sunken, dark-colored, dried or water-soaked cankers appear on (usually) lower stems. When the stem is split	Sanitation. See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		lengthwise, the hollow stems show a network of cross tissue 'threads'. Adventitious root initials may develop on surface areas of cankers.	
	Bacterial Wilt (<i>Ralstonia</i>)	Plant wilt rapidly. Lower stem vascular system and surrounding tissues may be brown. Ooze is visible from cut lower stem suspended in water.	Sanitation. Crop rotation away from susceptible crops. See ANR-797.
	Cladosporium Leaf Mold	A dark gray surface mold develops on leaves in blotches areas.	Sanitation. See the AL Pest Management Handbook.
	Cucumber Mosaic Virus	Plants stunted with mosaic, leaf distortions.	Sanitation. Aphid control may help a small amount.
	Double Virus Streak (Combination of Tobacco Mosaic Virus and Potato Virus X)	Leaves first show a light green mottle followed by numerous small (2 mm) gray-brown spots; severely damaged leaves may die. Later, leaves become dwarfed, curled, yellowed with small brown spots. Narrow, dark brown streaks develop on petioles and stems. Shoot tips may die. Plants are stunted. Fruit set reduced; fruits develop greasy, brown lesions.	Sanitation.
	Early Blight (<i>Alternaria solani</i>)	Oval or circular spots on leaves and stems.	Sanitation. See the AL Pest Management Handbook.
	Late Blight (<i>Phytophthora infestans</i>)	Dark brown-black blotches/spots on leaves/stem.	See the AL Pest Management Handbook.
	Phytophthora (Buckeye) Fruit Rot	Medium-brown, slightly sunken, zonate patterned with concentric rings, small to large spots develop on fruit. Brown discoloration may extend to fruit center.	Keep fruit away from the ground. See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Potato Virus Y	Plants stunted with some mosaic patterns, vein clearing.	Sanitation. Aphid control may help a small amount.
	Root Knot Nematode (<i>Meloidogyne</i>)	Plants are stunted. Roots are galled.	Sanitation. Resistant varieties or solarization.
	Septoria Leaf Spot	Small gray circular-angular spots.	See the AL Pest Management Handbook.
	Tobacco Etch Virus	Plants stunted with some mosaic patterns, vein clearing, yellowing, sometimes necrotic rings.	Sanitation. Aphid control may help a small amount.
	Tobacco Mosaic	Plants grow poorly. Leaves develop a mottle or mosaic of green and light green.	Remove damaged plants. Wash hands well before handling healthy plants.
	Tomato Spotted Wilt Virus	New growth stunted, leaves spotted and/or with ring spots; whole plant stunting and wilting; ringspots on fruit.	Sanitation. Thrips control will help in some situations.
Tumeric	Anthracnose	Circular-irregular light brown spots.	Sanitation. For commercial samples, see the AL Pest Management Handbook.
Turnip	Anthracnose	Irregular, light brown spots.	Sanitation.
	Black Rot (<i>Xanthomonas</i>)	See Collards.	Sanitation; crop rotation; see AL Pest Management Handbook.
	<i>Cercospora</i> Leaf Spot	Irregular gray-brown leaf spots with whitish centers and brown margins.	See AL Pest Management Handbook.
Verbena	Anthracnose (<i>Colletotrichum</i>)	Circular black spots on foliage.	Sanitation; Cleary's 3336.
	Pythium Lower Stem Rot	Brown water-soaked lesions.	Sanitation; reduce irrigation.
Veronica	Fusarium Crown Rot	Lower stems become dry rotted.	Sanitation Cleary's 3336, Halt, or OHP 6672.
	Pythium Crown & Root Rot	Wilt & dieback. Roots become slightly brown and wet rotted.	Sanitation. Protective drenches of Subdue may be used.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Viburnum	Anthracnose (<i>Colletotrichum</i> sp.)	Circular-irregularly shaped, brown leaf spots.	Sanitation. See the AL Pest Management.
	Cercospora Leaf Spot	Irregular, gray-brown leaf spots.	Sanitation. Protective sprays of Cleary's 3336 may be used.
Vinca (Annual Periwinkle)	Anthracnose (<i>Colletotrichum</i>)	Brown irregular areas, blotches develop on leaves and stems.	Sanitation. Protective sprays of Cleary's or Domain or a WP benomyl labeled on ornamentals may help.
	Phytophthora Stem Rot and/or Crown/Root Rot	Stems and/or lower stems near soil line and roots become browned and water-soaked.	Sanitation; improve soil drainage.
	Pythium Root Rot	Roots become brown decayed and water-soaked.	Sanitation. Reduce watering schedule. Rotate to different crop.
	Rhizoctonia Aerial Blight	Lower leaves become blighted; a thin mycelial webbing may develop.	Sanitation; Cleary's, Domain or a WP benomyl labeled on ornamentals may help.
Watercress	Cercospora Leaf Spot	Brown, circular-irregular leaf spots.	Sanitation.
	Pythium Stem & Root Rot	Stems and roots develop brown, water-soaked lesions.	Sanitation.
Watermelon	Fusarium Wilt	Plants show wilt & leaf yellowing beginning at base of plant.	Sanitation. Resistant varieties.
	Gummy Stem (<i>Mycosphaerella</i>)	Leaves develop black spots at leaf edges. Stem develop oozing, gummy cankers.	See AL Pest Management Handbook.
	Papaya Ringspot Virus	Leaves develop a mosaic (yellow-green); stunting.	Sanitation. Control of aphids may help a small amount.
Wax Myrtle	Anthracnose (<i>Colletotrichum</i>)	Brown, irregular spots, blotches develop on leaves.	Sanitation. If disease is severe, protective sprays of Cleary's 3336 or Domain or a WP benomyl may help.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Gummy Stem Blight (<i>Mycosphaerella</i>)	Black lesions/spots at leaf edges; elongate cracking on stem with amber-colored ooze.	See AL Pest Management Handbook.
Willow	Cercospora Leaf Spot	Irregularly shaped brown spots.	Sanitation of leaves in the fall.
Yaupon	Volutella Blight	Cankered, sunken stem areas, dieback, sometimes orange spore masses give the sunken areas and orange color.	Sanitation. Cleary's 336 or Halt protective sprays.
Zinnia	Alternaria Leaf Spot	Circular-oval, gray-brown spots. Zonate pattern sometimes seen.	Sanitation of diseased foliage. See the AL Pest Management Handbook.
Zoysia	Brown Patch (<i>Rhizoctonia</i>)	See Bermudagrass.	See the AL Pest Management Handbook.
	Dollar Spot (<i>Sclerotinia</i>)	Small spots in lawn (silver-dollar sized) become a white-gray colored. Individual grass blades develop dark water-soaked irregular spots which become whitish-gray with dark borders.	See the AL Pest Management Handbook or ANR-493.
	Rust (<i>Puccinia</i>)	Small (1-3 mm), yellow-orange-red flecks on grass blades; yellow-orange-red powder will wipe off on fingers. When severe, leaf blades will yellow and eventually die.	This is usually only a problem in shaded areas; fungicides may be applied when disease is severe; See the AL Pest Management Handbook or ANR-621.
	Take-All Patch (<i>Gaeumannomyces</i>)	Individual plants yellow, wither, die; black lesions on roots and stolons.	Keep soil pH at 5.5-6.0; use only ammonium-based fertilizer; avoid frequent irrigation.