

TIMELY INFORMATION

Agriculture & Natural Resources

May 18, 2009

PP-675

APRIL PLANT DISEASES FROM THE AUBURN PLANT DIAGNOSTIC LAB

APRIL PLANT DISEASES FROM THE BIRMINGHAM PLANT DIAGNOSTIC LAB

APRIL INSECT SAMPLES AT THE AUBURN PLANT DIAGNOSTIC LAB

DISEASE POSSIBILITIES FOR MAY

LAB NOTES

Jackie Mullen
Extension Plant Pathology Specialist

Jim Jacobi
Extension Plant Pathology Specialist

Charles Ray
Research Fellow IV-Auburn

Auburn Plant Disease Report-April (J. Mullen)

We received 89 plant samples for diagnoses in April. Ample rainfall and seasonal moderate temperatures have been conducive for a number of diseases.

Fire blight on apple and pear; Exobasidium galls on azalea and camellia; oak leaf blister; brown patch on turf grasses; anthracnose leaf spots on camellia, rhododendron, and other plants; rust leaf spots on oats; Helminthosporium-type leaf spots on triticale and wheat and powdery mildews have been common this spring.

Table 1. 2009 April Plant Diseases Seen In The Plant Diagnostic Lab at Auburn.

<u>Plant</u>	<u>Disease</u>	<u>County</u>
Camellia	Anthrachnose (<i>Colletotrichum</i>)	Elmore
Cantaloupe	Pythium Stem Rot	Geneva
Centipede	Brown Patch (<i>Rhizoctonia solani</i>)	Bullock, Jefferson, Tuscaloosa
Gardenia	Phytophthora Root Rot	*
Hydrangea, Oak Leaf	Armillaria Root Rot	Montgomery
<i>Hydrangea macrophylla</i>	Bacterial Leaf Spot	*
	Phytophthora Root Rot	*
Kudzu	Bacterial Leaf Spot (<i>Pseudomonas syringae</i> cv phaseolicola)	Mobile
Lupine	Sclerotinia Blight (<i>S. major</i>)	Lee
Oak	Oak Leaf Blister (<i>Taphrina deformans</i>)	Hale
Oats	Anthrachnose (<i>Colletotrichum graminicola</i>)	Escambia
	Crown Rust (<i>Puccinia coronata</i>)	Baldwin, Escambia
	Loose Smut (<i>Ustilago nigra</i>)	Baldwin
Rhododendron	Anthrachnose (<i>Colletotrichum</i>)	*
	Phytophthora Leaf Spot	*
Rose	Black Spot (<i>Diplocarpon rosa</i>)	Escambia
St. Augustine	Brown Patch (<i>Rhizoctonia solani</i>)	Mobile, Tuscaloosa
Triticale	Net Blotch (<i>Drechslera</i> , possibly <i>dictyoides</i>)	Baldwin

<u>Plant</u>	<u>Disease</u>	<u>County</u>
	Take-All Patch (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	Autauga, Mobile(2), Montgomery, Tuscaloosa
Veronica	Phytophthora Root Rot	*
	Septoria Leaf Spot	*
Wheat	Powdery Mildew (<i>Erysiphe graminis</i>)	Escambia
	Septoria Glume Blotch	Baldwin
	Tan Spot (<i>Drechslera tritici-repentens</i>)	Dallas

*Locations are not reported for nursery, greenhouse, and golf course samples.

Birmingham Plant Disease Report-April (J. Jacobi)

The lab received a total of 58 samples during the month of April. Towards the end of April the number of samples increased with typical spring diseases such as leaf gall and anthracnose on maples and other trees being widespread.

One of the more unusual samples was rust on mayapple, which is a plant we don't get in the lab very often. The symptoms of disease include bright yellow spots on the tops of the leaves and orange spots or pustules on the lower leaf surface. It is not considered a major problem and no control is needed. For pictures of the rust see the following web site (<http://www.herbarium.iastate.edu/fungi/fungispecies.php?sp=Puccinia+podophylli+Schwein>).

Table 2. 2009 April Problems Seen In The Birmingham Plant Diagnostic Lab.

<u>Plant</u>	<u>Problem</u>	<u>County</u>
Ash	Anthracnose	Jefferson
Aucuba	False Oleander Scale	Chilton
Azalea	Lacebugs	Jefferson
	Leaf Gall (<i>Exobasidium</i>)	Jefferson(2)
Centipedegrass	Large Patch (<i>Rhizoctonia</i>)	Jefferson
	Take-All Root Rot	Jefferson

<u>Plant</u>	<u>Problem</u>	<u>County</u>
Cleyera	Cercospora Leaf Spot	Shelby
Cypress, Bald	Lecanium Scale (<i>Parthenolecanium</i>)	Tuscaloosa
Holly 'Nellie R. Stevens'	Cottony Camellia Scale	Jefferson
Maple, Red	Anthraxnose	Jefferson
	Eyespot Gall Midge	Jefferson
Mayapple	Rust (<i>Puccinia</i>)	Blount
Oak, Willow	Lecanium Scale (<i>Parthenolecanium</i>)	Tuscaloosa
Pieris, Japanese	Lacebug	Jefferson
Magnolia, Saucer	Bacterial Blight (<i>Pseudomonas</i> suspected)	*
Magnolia, Southern	False Oleander Scale	Tuscaloosa
Rhododendron	Botryosphaeria Canker	Jefferson
Rose	Powdery Mildew	Jefferson
	Vole Damage	Jefferson
Wandering Jew	Mosaic Virus (Potyvirus)	*
Zoysiagrass	Fairy Ring	Jefferson

*Locations are not reported for nursery, greenhouse, and golf course samples.

Auburn Entomology Report-April (C. Ray)

County	Host	Category	Identification	Scientific Name
Colbert	Home	Household-Miscellaneous	Carpenter Ant	<i>Camponotus</i> sp.
Lee		Miscellaneous	An Andrenid Bee	<i>Andrena</i> sp.
Colbert	Home	Household-Stored	Indian Meal	<i>Plodia</i>

County	Host	Category	Identification	Scientific Name
		Products	Moth	<i>interpunctella</i>
Morgan	Tree	Miscellaneous	Luna Moth Cocoon	<i>Actias luna</i>
Montgomery	Viburnum	Ornamentals	Greenhouse Thrips	<i>Heliothrips hemarrhoidalis</i>
Russell	Garden	Row Crops	Weevil Larvae and Pupae	Curculionidae
Montgomery	Wax Myrtle	Ornamentals	Striped Mealybug Complex	<i>Ferrisia</i> sp.
Montgomery	Cleyera	Ornamentals	Florida Wax Scale	<i>Ceroplastes floridensis</i>
Lauderdale	Sago Palm	Ornamentals	Cycad Aulacaspis Scale	<i>Aulacaspis yasumatsui</i>
Montgomery	Camellia sinensis	Ornamentals	Barnacle Scale	<i>Ceroplastes cirripediformis</i>
Cullman	Home	Household-Miscellaneous	Wolf Spider Leg	Lycosidae
Walker	Car	Household-Miscellaneous	Gall Midge Larvae	Cecidomyiidae
Washington	Dead Tree Stump	Miscellaneous	Flat-headed Wood Boring Beetle	Buprestidae
Morgan	Japanese Holly	Ornamentals	English Walnut Scale	<i>Diaspidiotus juglansregiae</i>
Jefferson	Blueberries	Fruits & Nuts	Possible Bumblebee or Carpenter Bee Damage	<i>Bombus</i> sp. or <i>Xylcopa</i> sp.
Troup, GA	Muscadine Grape	Fruits & Nuts	Cicada oviposition damage	Cicadidae
Tuscaloosa	Bran	Household-Stored	Mold Mite	<i>Tyrophagus</i>

County	Host	Category	Identification	Scientific Name
		Products		<i>putrescentiae</i>
Marengo	Home	Household-Miscellaneous	May Beetle	<i>Phyllophaga</i> sp.
Calhoun	Home	Household-Miscellaneous	Bottle Fly	<i>Calliphora</i> sp.
Shelby	Home	Household-Stored Product	Varied Carpet Beetle	<i>Anthrenus verbasci</i>
Elmore	Pine Tree	Ornamentals	Eastern Subterranean Termite	<i>Reticulitermes flavipes</i>
Henry	Azalea	Ornamentals	Spider Mites	Tetranychidae
Calhoun	Boxwood	Ornamentals	Chinese Wax Scale	<i>Ceroplastes ceriferus</i>

Disease Possibilities For May

As is usual in May, we are seeing an abundance of turf grass samples. Thus far, many of our turf samples have involved non-infectious situations as is typically the case for turf problems sent to us in April and early May. But, we are seeing brown patch as a fairly common problem on warm-season turf grasses. We received our first sample of tomato spotted wilt virus on tomato on May 1. Also, we received an uncommon rust leaf spot disease on hibiscus. The tentative diagnosis of *Puccinia* sp. is awaiting confirmation by a mycologist.

The list below includes some common disease problems received in the lab during May of the past few years. Comments on control practices are brief. Refer to the Alabama Pest Management Handbook or appropriate fact sheet or disease notes publications for details on disease control.

Table 3. Brief Disease Descriptions and Control Recommendations for Diseases Often Seen in May.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Any	Slime Mold	Not actually a disease; gray, black, pink, or yellow slimy or powdery masses on plants or soil; plasmodium or slime stage may be a watery mass the	Wash off with a strong stream of water; physical removal.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		consistently of a thinnish layer of 'Jello'; the spore stage is often powdery with the coloration of black, pink, yellow or orange from spore production.	
Apple	Armillaria Root Rot	Dieback; thin, white mycelial layer just under bark, appressed to wood.	Removal of tree and roots; maintain healthy trees.
	Botryosphaeria Canker	Cracked, sunken lesions on trunk or branches. If lesion girdles the branch, dieback results.	Prune out cankers making cuts 3 inches beyond the canker edge. Dip shears into alcohol or a 10% solution of bleach between cuts.
	Cedar-Apple Rust (<i>Gymnosporangium</i>)	Bright yellow leaf spots (up to ¼ or ½" diameter).	Protective fungicide sprays; Removal of junipers. See ANR-408.
	Fire Blight (<i>Erwinia</i>)	Small (⅛-¼" diameter) circular-irregular spots with purple margins and brown centers.	Sanitation in the fall; protective sprays during growing season.
	Powdery Mildew (<i>Podosphaera</i>)	Whitish powdery dusting on foliage; leaf deformity; blight.	Sanitation in the fall; protective fungicide sprays.
Armaryllis	Stagnospora Leaf Spot	Fairly large (¼-½ inch long) oval or elliptical or irregularly shaped red spots.	Sanitation. Protective sprays of Cleary's 3336 or Domain.
Ash	Anthrachnose (<i>Discula</i>)	Brown irregular spots blotches.	Sanitation of fallen leaves.
Azalea	Aerial Blight (<i>Rhizoctonia</i>)	Mostly seen in greenhouse-nursery situations; lower leaves become blighted and fall.	Sanitation; protective fungicide sprays.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Colletotrichum Leaf Spot	Small, brown, circular leaf spots develop.	Sanitation; See the AL Pest Management Handbook.
	Exobasidium Gall	Swollen blossoms, leaf, and shoot galls. From mid April-mid May, galls change from a green to a white or pink-white color.	Sanitation; removal of galls while they are still green. Protective fungicide application may be made.
	Phomopsis Blight & Dieback	Sunken cankers develop on twigs and small branches; twig dieback results.	Prune out cankers. Make cuts at least 2 inches beyond the damage.
	Phytophthora Root Rot	Roots brown and water-soaked.	Sanitation. Protective fungicide drenches in nursery situations.
	Powdery Mildew	White, powdery, dusting on leaf & young twigs, blight.	Pruning; protective fungicide sprays; See the AL Pest Management Handbook.
Bahia Grass	Dollar Spot (<i>Sclerotinia</i>)	Silver dollar-sized, bleached-out spot areas in lawn; irregular-shaped whitish leaf spots on grass blades.	No control available. See cultural controls in ANR-493.
	Take-All Patch (<i>Gaeumannomyces</i>)	Dieback, wilt, crown and root rot	See cultural controls ANR-823.
Bean, Garden	Alternaria Leaf Spot	Dark brown angular-circular spots (about 5 mm diameter) scattered on leaf surfaces.	Bravo 720 should control this problem.
	Anthrachnose (<i>Colletotrichum</i>)	Circular-irregular, reddish-brown pod, leaf & stem spots develop.	Sanitation. See the AL Pest Management Handbook.
	Bacterial Blight	Greasy, dark green irregular spots 1/8-1/4" diameter) become black and papery with wet looking margins sometimes	Protective sprays. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		surrounded by a yellow zone.	
	Fusarium Stem Rot	Red-brown sunken lesions on lower stems.	Plant removal, crop rotation away from beans and peas.
	Mosaic Viruses	Distinct pattern of dark green & light green or yellow on leaves.	Control weeds; control beetles and aphids; Do not save seed; remove diseased plants.
	Pythium Root Rot	Roots are soft, brown, water soaked.	Avoid wet conditions.
	Rhizoctonia Soreshin	Dark reddish-brown sunken lesion(s) on lower stem.	See AL Pest Management Handbook for a before-planting soil treatment.
Bee Balm	Powdery Mildew	Powdery white dusting on leaves; blight; distorted new growth.	Cleary's 3336; sanitation.
Begonia	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Dark brown angular spots (about 5 mm diameter or less). Outer edges of spots may appear wet (water soaked); centers of spots may dry and crack. Spots may coalesce.	See the AL Pest Management Handbook; Sanitation.
	Phytophthora Root Rot	Dieback, wilt, wet rot of roots.	Remove damaged plants; reduce water levels. See the AL Pest Management Handbook.
	Pythium Root Rot	Dieback, wilt, wet rot of roots.	Remove damaged plants; reduce water levels. See the AL Pest Management Handbook.
	Rhizoctonia Crown & Root Rot	Dieback; wilt, lower leaves yellow; dry, brown lesions on lower stem.	Remove damaged plants; See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Bentgrass	Anthrachnose (<i>Colletotrichum</i>)	Yellow to brown irregular, elongate spots develop on leaves.	Irrigate in the early morning; collect grass clippings; avoid stress; See the AL Pest Management Handbook.
	<i>Bipolaris cyanodontis</i> Leaf Spot	Small, elliptical brown spots; spots may coalesce into blighted grass blades.	Sanitation of cut grass blades; good fertility practices; see AL Pest Management Handbook for recommended fungicides.
	Pythium Blight	Wet-looking blight of foliage.	See the AL Pest Management Handbook; decrease irrigation.
	Pythium Root Rot	Dieback, wilt, wet rot of roots.	Remove damaged plants; reduce water levels; See AL Pest Management Handbook for protective fungicides.
Bermuda	Algae (<i>Cephaleuros</i>)	Dark green, thin, flat 'sheets' of algae develop on grass foliage.	Promote good drainage and drying out of the area; See the AL Pest Management Handbook.
	<i>Bipolaris</i> Leaf Spot/Blight	Smallish (1/8 inch or less) elongated brown spots on grass blades. Spots may coalesce.	See AL Pest Management Handbook for recommendations under Helminthosporium melting-out.
	Brown Patch (<i>Rhizoctonia</i>)	Circular-irregular patches in lawn become brown. Brown lesions present on individual grass blades.	Reduce nitrogen fertilization. Sanitation. Protective fungicide treatments.
	Dollar Spot (<i>Sclerotinia homeocarpa</i>)	Small pale yellow spots the size of a silver dollar are characteristic; spots often coalesce; leaves show characteristic cream-colored spots	Improve fertilization and provide adequate irrigation. In some situations, protective fungicide treatments are needed. See ANR-493 and the AL Pest

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		with dark brown borders.	Management Handbook.
	Exserohilum Blight	Smallish (1/8 inch or less) elongated brown spots on grass blades. Spots may coalesce.	See AL Pest Management Handbook for recommendations under Helminthosporium melting-out.
	Rust (<i>Puccinia</i>)	Foliage becomes pale green-brown with an orange tint; orange spores masses will rub off on a finger wiped across leaf surfaces.	See the AL Pest Management Handbook.
	Spring Dead Spot (<i>Gaeumannomyces graminis</i> pv <i>graminis</i>)	Large circular areas fail to green-up in the spring.	See the AL Pest Management Handbook.
Bermuda, Coastal	Rust (<i>Puccinia</i>)	Leaf spots, blight; reddish color of blight.	Frequent mowing.
Birch, River	Anthrachnose (<i>Cryptocline</i>)	Brown leaf spots and blotches develop on leaves.	Clean up & remove all leaf litter this fall.
Blackberry	Anthrachnose (<i>Colletotrichum</i>)	Irregularly-shaped brown spots & blotches on leaves and canes.	See the AL Pest Management Handbook.
	Botryosphaeria Canker	Dry, cracked sunken lesions.	Pruning.
	Cane Blight (<i>Leptosphaeria</i>)	Brown, lesions on canes.	Pruning.
	Double Blossom (<i>Cercospora</i>)	Floral canes develop abnormal flowers with thickened petals. Internodes are shortened. Leaf development at nodes is abnormally abundant.	Sanitation as soon as abnormality is discovered. Protective fungicide treatment.
	Phoma Cane Blight	Large brown lesions on canes; dieback.	Sanitation.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Septoria Leaf Spot	Small (usually ¼ inch) angular brown spots.	See fungicide recommendations under anthracnose.
Blueberry	Botryosphaeria Blight	Somewhat inconspicuous sunken cankers develop along branches or on lower trunk areas. Dieback results. (Sometimes this condition is associated with soils excessive in phosphorus & calcium.)	Sanitation. Benlate protective sprays.
	Botrytis Blossom Blight and Stem Canker	Blossoms become brown; stems develop sunken lesions.	Sanitation; Benlate protective sprays.
Boxwood	Macrophoma Blight	Dieback develops; leaves turn yellow or white; black, circular bodies develop on leaf surface.	Prune out damage; avoid stress or other damage. See ANR-0222.
	Phytophthora Root Rot	Dieback, yellowing of lower foliage, wet decay of roots.	Remove damaged plants; reduce water levels; see AL Pest Management Handbook.
	Volutella Blight	Sunken cankers on stems and branches. Surface areas of cankers may be covered with orange spore masses of the fungus.	Pruning of cankered stems. After pruning operations are complete, a protective spray treatment of Cleary's 3336 may be helpful. Avoid stressful situations.
Broccoli	Downy Mildew (<i>Pseudoperonospora</i>)	Light yellow spots/blotches on upper leaf surfaces; gray spots on corresponding lower leaf areas.	See the AL Pest Management Handbook.
Brussel Sprouts	Black Rot (<i>Xanthomonas</i>)	Black V-shaped lesions on leaf edges; black leaf veins; black decay of inner stems; wilt.	Sanitation; See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	<i>Sclerotium rolfsii</i> Crown Rot	Brown decay at soil line. Brown, mustard seed-like structures of fungus may be present at soil line.	Sanitation. See A. Hagan.
Butter Beans	Anthrachnose (<i>Colletotrichum</i>)	Dark brown irregular-circular sunken spots on foliage, pods. Spots become covered with rusty-colored spores.	See AL Pest Management Handbook.
	Cercospora Leaf Spot	Medium-light brown irregular spots.	See AL Pest Management Handbook under Anthracnose.
Camellia	Anthrachnose (<i>Colletotrichum</i>)	Circular or irregular brown or cream-colored spots.	Sanitation of fallen leaves; Cleary's 3336.
	Exobasidium Gall	See azalea.	Sanitation.
Camellia, Sasanqua	Anthrachnose (<i>Glomerella</i> spp.)	Sunken branch lesions, often round or oval; dieback.	Pruning of cankered areas.
Cantaloupe	Bacterial Wilt (<i>Erwinia</i>)	Individual leaves wilt, then lateral shoots also wilt. Eventually whole plant wilts and dies.	Sanitation; use insecticide to control cucumber beetles.
	Cercospora Leaf Spot	Circular-irregular brown leaf spots.	Chlorothalonil protective sprays.
	Gummy Stem (<i>Mycosphaerella</i>)	Brown, elongated, cracked lesions with a gummy exudate.	See the AL Pest Management Handbook.
Catalpa	Alternaria Leaf Spot	Mostly circular brown spots.	Collect and remove all leaf litter this fall.
Cedar, Eastern Red	Kabatina Tip Blight	Branch tips become brown; necrosis spreads down the branch.	Sanitation of diseased plant parts; Cleary's 3336 may provide some protection.
Centipede	Anthrachnose (<i>Colletotrichum</i>)	Yellow-brown, irregular-elongate spots; blight.	Keep the area dry; water early in the day; collect grass clippings; see the AL Pest

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			Management Handbook.
	Brown Patch (<i>Rhizoctonia</i>)	Browning patches or rings of grass. Brown, irregular leaf spots may also occur.	Fungicide sprays; collect clippings; avoid excess nitrogen application.
	Ring Nematode (<i>Criconeoides</i>)	Dieback; poor growth.	See the AL Pest Management Handbook.
	Take-All Patch (<i>Gaeumannomyces</i>) (Possibly a secondary occurrence or weak pathogen.)	Individual plants yellow and thin out. Stolons and roots develop black lesions.	See PP-312. Bayleton is labeled.
Cherry	Armillaria Root Rot	Dieback, lower trunk and root decay. Check for thin layer of white mycelium under the bark, honey-colored mushrooms, and black rhizomorphs.	Tree removal.
	Peach Leaf Curl & Plum Pockets (<i>Taphrina</i> spp.)	Firm irregular galls or concave-convex gall-swellings on leaves.	Fungicide applications in late winter before bud swell, and at bud break.
	Phomopsis Canker	Gray-brown, sunken elliptical lesions on twigs, small branches.	Sanitation.
	Septoria Leaf Spot	Brown, irregularly shaped spots (1/4-1/3 inch diameter) on leaves.	Sanitation.
Cherry Laurel	Algal Leaf Spot (<i>Cephaleuros</i>)	Slightly raised, reddish-green circular spots with irregular edges.	Sanitation. Reduce humidity.
Cherry, Yoshino	Cherry Leaf Spot (<i>Blumerilla</i>)	Brown circular-irregular leaf spots.	Sanitation of fallen leaves.
Chestnut, Chinese	<i>Lasiodiplodia theobromae</i> Canker	Elliptical, sunken lesions develop on trunk and branches.	Prune out damage making cuts 3 inches beyond edge of

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			canker. Dip shears into alcohol or 10% solution of bleach between cuts.
Christmas Cactus	Rhizoctonia Stem Rot	Dry, brown decay of stem.	Sanitation. Protective sprays of Cleary's 3336 or Halt.
Clematis, Evergreen	Pythium Root Rot	Light brown, decayed roots, dieback; wilt.	Remove damaged plants; Keep the area well drained. Allow areas to dry out.
Cleyera	Phytophthora Root Rot	Dieback, wilt; roots become water soaked and rotted.	Sanitation. Reduce irrigation and/or improve soil drainage.
Coleus	Phytophthora Root Rot	Roots are brown and water soaked. Tops collapse.	Sanitation. Avoid poorly drained sites.
	Pythium Root Rot	Plants show poor growth, dieback; roots are discolored to a light brown; root cortex separate easily from the central cylinder.	Sanitation; Improve soil drainage.
Collard	Bacterial Leaf Spot	Dark brown-black angular, vein-bound leaf spots.	Sanitation; do not irrigate overhead. See AL Pest Management Handbook.
Corn	Bacterial Stalk Rot	Brown lesions cause collapse of stem.	Sanitation.
	Fusarium Crown Rot	Crown develops dry, reddish-brown rot; tops wilt and dieback.	Sanitation; crop rotation.
	Ring Nematode (<i>Criconemoides</i>)	Plants become stunted; may appear nutrient deficient.	Crop rotation; consult with Ed Sikora.
	Smut (<i>Ustilago maydis</i>)	Black powdery galls form on ears, tassels, stalks.	Sanitation; crop rotation, resistant hybrids and varieties.
Coral Bells	Black Root Rot (<i>Thielaviopsis</i>)	Foliage wilt and dieback; roots develop black tips	Remove damaged plants. Cleary's 3336 drenches may provide

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		and areas.	some protection.
Cotton	Alternaria Leaf Spot	Medium brown, circular or oval leaf spots, blight.	See Ed Sikora.
	Fusarium Seedling Disease	Red-brown lesions on lower stem near soil line.	Seed Treatment.
	Pythium Root Rot	Roots become brown and water soaked. Foliage becomes yellowed, wilted and browned.	Seed Treatment.
	Rhizoctonia Seedling Disease	Medium-dark brown lesions on lower stem near soil line.	Seed Treatment.
	Thielaviopsis Root Rot	Black spots/lesions along roots.	----
Cowpea	Cercospora Leaf Spot	Brown, circular to irregular leaf spots.	Sanitation.
Crabapple	Botryosphaeria Canker and Frogeye Leaf Spot	Brown circular-irregular leaf spots with dark margins; sunken brown lesions on twigs, branches.	Sanitation; Cleary's 3336, Domain, or a benomyl labeled for ornamentals may help.
	Cedar Apple Rust (<i>Gymnosporangium</i>)	Bright yellow circular spots (1/4-1/2 inch diameter) on leaves; orange spore masses (circular projections) develop on leaves (lower leaf surfaces) and fruit.	Remove junipers or apply protective fungicide sprays to crabapples. See AL Pest Management Handbook.
	Fire Blight (<i>Erwinia amylovora</i>)	Blossom blight; rapid dieback; cankers form at end of the dieback.	Prune off dieback areas. Make cuts 10-12 inches beyond margin of dieback. Remove cuttings. See ANR-542 and the AL Pest Management Handbook.
	Scab (<i>Venturia inaequalis</i>)	Olive, slightly raised small spots develop on leaves and fruit.	Sanitation. See the AL Pest Management Handbook for

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			fungicides.
Crape Myrtle	Powdery Mildew	Initially, a white dusting of the fungus appears on upper leaf surfaces. Eventually, infected leaves turn brown. Leaves may be curled and distorted.	Prune branches to help reduce humidity. See the AL Pest Management Handbook.
Cryptomeria	Cercosporidium Blight	Lower limbs develop a browning of needles. Disease usually starts on inner needles where humidity is higher. Disease will spread upward.	In a commercial situation, trees may be sprayed with Kocide for protection.
	Phytophthora Root Rot	Lower limbs dieback first; roots are water-soaked, brown, rotted.	Remove damaged trees. Reduce water in the area. Check with A. Hagan for fungicide use in commercial situation.
Cucumber	Anthrachnose (<i>Colletotrichum</i>)	Circular, light-medium brown spots.	Sanitation; See the AL Pest Management Handbook.
	Bacterial Wilt (<i>Erwinia</i>)	Individual leaves become wilted, yellowed and finally brown. Eventually, whole stem sections wilt and die.	Control cucumber beetles.
	Cercospora Leaf Spot	Cream-colored, angular spots.	See the AL Pest Management Handbook.
	Pythium Root Rot	Roots become brown and water-soaked; outer cortex slides easily away from root central cylinder.	Sanitation.
	Rhizoctonia Seedling Soreshin	Dieback, wilt, lower leaf yellowing; lower stem dark brown, dry lesions.	Sanitation.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Daisy, Gerbera	Powdery Mildew	White powdery dusting on leaf and blossom surfaces; blight.	See the AL Pest Management Handbook.
Daylily	Kabatiella Streak	Yellow spots and elongated streaks on leaves.	Sanitation of spotted leaves; Cleary's 3336 or Halt may be used.
	Rust (<i>Puccinia hemerocallidis</i>)	Elongated yellow-brown spots, blight on leaves. Orange pustules evident at early stages of disease.	Sanitation. See TI PP-506. For homeowners, Immunox or Ferti-loam System Fungicide may be used.
Dogwood	Anthracoze (<i>Discula</i>)	Large brown circular-irregular spots (1/4-1/2 inch diameter) with dark brown-purple borders develop first on foliage of lower branches. Leaf spots merge. Dieback will result. Cankers develop.	Sanitation. Protective fungicide sprays.
	Botrytis Blight	Foliage shows grey-brown blotches, spots.	See AL Pest Management Handbook under spot anthracnose.
	Phyllosticta Leaf Spot	Small (1/4 inch or less) circular brown spots with dark brown margins.	Sanitation of fallen leaves in the fall. See the AL Pest Management Handbook under Septoria.
	Phytophthora Crown & Root Rot	Crown and root rot tissues brown discolored and water soaked.	Sanitation. See the AL Pest Management Handbook for fungicide recommendations.
	Powdery Mildew	White powdery dusting on leaf and blossom surfaces; blight.	See the AL Pest Management Handbook.
	Spot Anthracnose (<i>Elsinoe</i>)	Small (1/16 inch or less diameter) circular reddish spots	Sanitation in the fall. Protective fungicide sprays recommended

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		develop on bracts; spots on leaves are similar in size but brownish red in color and some leaf spots may be irregular in shape.	only for specimen trees.
Eggplant	Pythium Damping-Off	Seedling plants develop a soft rot with light brown discoloration on stem at soil line; plants collapse and fall over.	Avoid wet situations when planting seeds.
Elm	Powdery Mildew (<i>Phyllactinia</i>)	White powdery patches develop on leaves; some leaf deformity.	Sanitation of fallen leaves. See the AL Pest Management Handbook.
Euonymus	Powdery Mildew	See description above for powdery mildew on elm.	See comments for powdery on elm.
Fescue	Net Blotch (<i>Drechslera</i>)	Brown, irregularly pigmented rectangular shaped blotches (1/8 inch or less diameter) on leaf blades. Spots may merge to cover large areas.	See AL Pest Management Handbook under Helminthosporium melting out.
Forsythia	Crown Gall (<i>Agrobacterium tumefaciens</i>)	Irregular, rough, somewhat spherical galls on branches and lower trunk.	Sanitation; crop rotation.
	Phomopsis Dieback	Branch dieback; sunken cankers form at end of dieback.	Pruning 3-4 inches beyond dieback margin; Remove cuttings.
Gardenia	Anthrachnose	Brown leaf spots and blotches appear.	Remove all fallen leaves from the area. See the AL Pest Management Handbook.
Geranium	Bacterial Blight (<i>Xanthomonas</i>)	Black leaf spots and stem rot lesions. Yellow-brown	Sanitation. Bordeaux Mixture.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		wedges may develop in spotted areas of leaves. Lower leaves may wilt and die due to systemic infections.	
Gerbera Daisy	Powdery Mildew	White, powdery dusting on upper leaf surfaces.	Sanitation. See the AL Pest Management Handbook.
Ginger	Rhizoctonia Crown Rot	Lower stem at soil level & upper root ball develops dry, brown decay.	Remove damaged plants; rotate away from ginger.
Ginseng	Fusarium Root Rot	Plants wilt and dieback or show poor growth.	Crop rotation.
Grape	Black Rot (<i>Guignardia</i>)	Medium-black brown circular spots (1/8-1/4 inch) on leaves and fruit.	Protective fungicide sprays; sanitation.
Greenhouse Crops	Tomato Spotted Wilt Virus	Symptoms are variable but include gray-black leaf spots/blotches, yellow ringspot patterns, stunted growth malformed stunted and/or spotted new growth, and wilt.	Sanitation; control thrips.
Hawthorn	Anthracnose	Circular-irregular shaped, brown/leaf spots.	Remove all fallen leaves; protective sprays of Cleary's 3336, Halt, or OHP 6672 may be used if desired.
	Botryosphaeria Twig Canker	Dry, sunken, cracked lesions on twigs; dieback.	Prune out cankers making cuts at least 2 inches from lesion edge.
Hibiscus	Bacterial Leaf Spot	Dark brown, angular, wet spots; yellow halo may be present.	Sanitation of fallen leaves, spotted leaves; See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Hickory	Downy Leaf Spot (<i>Pseudocercospora</i>)	Cream-yellow spots with indistinct margins on upper leaf surface; corresponding lower leaf surface spots are brown.	See AL Pest Management Handbook for pecan scab fungicides.
Holly	Phytophthora Root Rot	Dieback; roots become water soaked and rotted.	Sanitation. Reduce irrigation and/or improve soil drainage. See the AL Pest Management Handbook.
Holly, Inkberry	Phytophthora Root Rot	Dieback; lower leaves yellowing; wilt; wet, brown rot of roots.	Remove dying plants; reduce water levels; See the AL Pest Management Handbook.
Holly, Japanese 'Compacta'	Botryosphaeria Canker	Small-large sunken lesions with cracking at lesion edges; dieback.	Pruning.
Holly, Yaupon	Armillaria Root Rot	Dieback of lower branches first; brown mushrooms may develop at plant base. See ANR-907.	Remove plant. See ANR-907.
Honeysuckle	Anthrachnose (<i>Colletotrichum</i>)	Circular-irregularly shaped brown leaf spots.	Sanitation.
Horse Chestnut	Leaf Blotch (<i>Guignardia</i>)	Angular, brown leaf spots.	Collect and remove all fallen leaves this fall.
Hosta	Anthrachnose (<i>Colletotrichum</i>) Leaf Spot	Brown, irregularly-shaped leaf spots.	Sanitation; Cleary's 3336 or Domain protective sprays.
Hydrangea	Alternaria Leaf Spot	Circular to oval to irregular brown leaf spots.	Sanitation of fallen leaves; Daconil.
	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Dark brown-black angular spots develop; some reddish discoloration is present.	Remove damaged foliage; do not water over-head; copper protective sprays (Phyton) may help some.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Botrytis Blossom Blight	Irregular brown spots and blotches on flowers.	Sanitation; Cleary's 3336, Domain or benomyl product.
	Colletotrichum Blossom Blight	Irregular brown spots blotches on flowers, sometimes orange patches of spores are present.	Sanitation; Cleary's 3336 or Domain, or benomyl.
	Phyllosticta Leaf Spot	Brown circular spots, sometimes with dark border.	Sanitation; Cleary's 3336 or Domain or benomyl product.
Hydrangea, Oakleaf	Armillaria Root Rot	Dieback; thin, white mycelial layer just under bark, appressed to wood. Black, thread-like structures may be present above or below surface of root or crown; brown mushrooms may be present.	Removal of tree and roots; maintain healthy trees.
	Bacterial Leaf Spot	Black, angular spots that sometimes show water-soaked edges or a yellow halo.	Remove all spotted foliage. Do not water over-head.
Impatiens	Botrytis Foliage Blight	Blight.	Sanitation; See the AL Pest Management Handbook.
	Impatiens Necrotic Spot Virus	Irregularly shaped black spots/blotches (1/8-1/4 inch diameter-usually) which may be confused with bacterial infection. Black ring spots may be present.	Control thrips. Sanitation.
	Rhizoctonia Stem Rot	Lower stems develop brown sunken lesions; plants dieback.	Sanitation; Cleary's protective sprays; crop rotation.
Indian Hawthorn	Entomosporium Leaf Spot	Reddish, irregular-shaped leaf spots sometimes with black, swollen, cushion like bodies scattered on leaf spot	Remove spotted foliage from under plants; prune as needed; See AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		surface.	
Iris	Fusarium Root Rot	Dieback, wilt, yellowing; dry, brown decay of roots.	Removal of dying plants; rotation away from iris and related plants.
	Heterosporium Leaf Spot (<i>Heterosporium=Cladosporium</i>)	Small-large (1/4-1/2 inch long), elliptical or oval shaped medium brown leaf spots.	Sanitation. See AL Pest Management Handbook.
	Rust (<i>Puccinia</i>)	Small, red-orange, powdery, raised spots.	Sanitation.
Ivy, English	Anthrachnose (<i>Colletotrichum</i>)	Irregularly-shaped brown leaf spots.	Sanitation; See the AL Pest Management Handbook.
	Bacterial Leaf Spot	Dark, brown, wet, angular leaf spots.	Sanitation. See the AL Pest Management Handbook.
Jacobinia	Impatiens Necrotic Spot Virus	Stunting, wilt, leaf spots.	Sanitation; thrips control.
Juniper	Cedar Apple Rust (<i>Gymnosporangium</i>)	Hard, spherical twig or branch galls; some galls produce gelatinous, orange, finger-like projections of spore masses.	Remove galls before spore production. Remove cedar or apple trees; protective fungicide sprays. See ANR-468.
	Kabatinia Tip Blight	Juniper twigs dieback; appears similar to Phomopsis tip blight.	Sanitation; refer to fungicides listed for control of Phomopsis tip blight.
	Pestalotiopsis Blight	Blight of inner foliage of stressed plants.	Sanitation. See the AL Pest Management Handbook under Phomopsis Twig Blight for protective fungicides.
	Phomopsis Blight	Tips of twigs on lower branches become brown. The blight spreads down the twig and upward in the plant.	Sanitation. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Phytophthora Root Rot	Roots brown, water soaked.	See AL Pest Management Handbook.
Kiwi	Anthrachnose (<i>Colletotrichum</i>)	Circular-irregular, light to medium brown spots.	Sanitation.
Kudzu	Bacterial Leaf Spot (<i>Pseudomonas</i>)	Small black circular spots with a yellow halo.	---
Lemon	Phytophthora Root Rot	Foliage dieback and wilt; lower leaves may turn yellow; roots develop dark color and wet rot.	Remove dying plants & root associated soil; reduce water levels in soil.
Leyland Cypress	Botryosphaeria Canker	Elongate sunken trunk and branch brown lesions.	Sanitation.
	Cercosporidium Blight	Lower, inner limbs develop blight which gradually spreads up and outward on foliage.	Pruning of damaged limbs; See the AL Pest Management Handbook.
	Pestalotiopsis Canker	Elongate sunken trunk and branch brown lesions.	Sanitation.
	Seiridium Canker	Elongate sunken trunk and branch brown lesions.	Sanitation.
Ligustrum	Cercospora Leaf Spot	Medium brown, irregularly shaped spots (1/4-1/2 inch long).	Sanitation. See the AL Pest Management Handbook.
Lily, Spider	Stagnospora Leaf Spot	Reddish colored leaf edge scorch and leaf spot.	Sanitation.
Liriope	Fusarium Crown Rot	Dry rot of lower stem at the soil line. May follow after stress.	Avoid stress; Cleary's 3336 may provide some protection.
Loropetalum	Botrytis Leaf Spot	Brown-gray irregular leaf spots.	Sanitation of fallen leaves; Cleary's 3336 protective sprays.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Lupin	Rhizoctonia Crown Rot	Brown dried lesions on stems near the soil line.	---
Magnolia, Southern	Algal Leaf Spot (<i>Cephaleuros</i>)	Usually circular, greenish, slightly raised spots. Spot edges slightly more raised than spot center. Edges are irregular.	Sanitation. See the AL Pest Management Handbook.
	Bacterial Leaf Spot (<i>Pseudomonas cichorii</i>)	Small, angular, water soaked spots.	Sanitation. Kocide 101 or Bordeaux mixture.
	Black Mildew (<i>Meliola</i>)	Black surface mold on lower leaf surface.	-----
Mahonia, Chinese	Powdery Mildew	White dusting on foliage; affected area later turns brown.	Prune to reduce humidity; Cleary's 3336 would provide protective disease control.
Mahonia, Leatherleaf	Leaf Spot (<i>Phyllosticta</i>)	Brown, angular leaf spots.	Sanitation of fallen leaves; Cleary's 3336 protective sprays.
Maple	Anthrachnose (<i>Kabatella</i>)	Brown-black spots/blotches which often follow along leaf veins; sometimes blotchy areas occur along leaf edges.	Sanitation. See AL Pest Management Handbook for small trees.
	Purple Eye Spot (<i>Phyllosticta</i>)	Purple circular spots (about ¼ inch diameter) with dark purple, black border.	Sanitation. See the AL Pest Management Handbook.
	Tar Spot (<i>Rhytisma</i>)	Black irregularly-shaped spots.	Sanitation of fallen leaves this fall. See the AL Pest Management Handbook.
Maple, Japanese	Botryosphaeria Canker	Sunken, elliptical cracked, dry lesions on trunk or branches.	Pruning. Make cuts 3-4 inches from lesion edges.
	Phyllosticta Leaf Spot	Cream-light brown circular spots with dark brown or purple	Sanitation of fallen leaves this fall. See the AL Pest

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		margins.	Management Handbook.
	Phytophthora Crown & Root Rot	Initially roots become brown and water-soaked. Later, dead roots dry out. Dieback.	Remove tree. Improve drainage. Plant resistant plant selection, if possible.
Marigold	Anthrachnose (<i>Colletotrichum</i>)	Circular, cream-colored spots (1-3 mm diameter) on leaves.	Sanitation. Cleary's 3336 would control anthracnose and it is labeled on marigold.
	Botrytis Flower/Foliage Blight	A brown-gray foliage blight.	Sanitation; See the AL Pest Management Handbook.
	Fusarium Wilt	Lower leaves wilt and turn yellow. Yellowing and wilt slowly progress up the plant. Vascular tissues turn brown.	Sanitation. Rotate away from marigolds for 10-16 years.
	Pythium Root Rot	Roots become light brown, water-soaked, and tissues pull apart easily.	Sanitation. Reduce soil water content. See the AL Pest Management Handbook.
Mayapple	Rust (<i>Puccinia</i>)	Yellow angular spots may develop; eventually, orange powdery dots of spore pustules appear; leaf blight follows.	Sanitation. Mayapple is only host for this rust.
Mayhaw	Cedar Quince Rust (<i>Gymnosporangium claviceps</i>)	Bright yellow spots may occur on fruit and leaves; tiny black spots may be seen in spots upper leaf surface; orange powdery specks may be seen on lower leaf spots.	If practical, remove junipers in the immediate area.
Million-Bells	Pythium Root Rot	Dieback, wilt, roots are light brown and decayed.	Sanitation. Reduce water in the area.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Mondograss	Anthrachnose (<i>Colletotrichum</i>)	Gray-brown spots, blotches, often at leaf tips, margins.	Sanitation. Cleary's or Domain.
Mulberry	Cylindrosporium Leaf Spot	Brown circular-irregular leaf spots.	Sanitation.
Muscadine	Black Rot (<i>Guignardia</i>)	See Grape.	See AL Pest Management Handbook.
Nandina	Cercospora Leaf Spot	Circular to irregular, brown spots.	Sanitation of fallen leaves; See the AL Pest Management Handbook.
Nectarine	Brown Rot	See Peach.	
Oak	Anthrachnose (<i>Apiognomonina</i>)	Medium-brown blotches along leaf veins, in interveinal areas and along leaf margins.	Sanitation. See AL Pest Management Handbook for small trees.
	Oak Leaf Blister (<i>Taphrina</i>)	Concave-convex spot areas that are slightly swollen and slightly discolored.	Sanitation. See AL Pest Management Handbook.
Oak, Red	Fusiform Rust (<i>Cronartium querauum</i> f. sp. Fusiforme)	Very small yellow to dark brown leaf spot; thin, short, dark brown hair-like structures on lower leaf spots.	Damage to oak is not significant. Remove pines (loblolly & slash) in the area.
Oak, Shumard	Fusiform Rust (<i>Cronartium querauum</i> f. sp. Fusiforme)	Very small yellow to dark brown leaf spots; thin, short, dark brown hair-like structures on lower leaf spots.	Damage to oak is not significant. Remove pines (loblolly & slash) in the area.
Oak, Water	Fusiform Rust	See Shumard Oak above.	
Oats	Loose Smut (<i>Ustilago avenae</i>)	Plants shorter than normal; the grain is replaced by dark-brown to black powdery masses of smut spores. Usually all the grain on an	Treated seed with fungicide.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		infected plant is smutted.	
	Rust (Crown or Leaf) [<i>Puccinia coronata</i>]	Bright orange-yellow round or oblong powdery pustules on leaves, sheaths, stems, panicles. Plants are weakened when infection is severe.	Deep plow.
Okra	Rhizoctonia Crown & Root Rot	A brown, dry decay of crown & roots.	Sanitation. Crop rotation.
Oxalis	Kuehneola Rust	Orange powdery pustules on leaves.	Sanitation. Crop rotation.
Peach	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Small (2-6 mm diameter) circular, brown-black spots with wet-looking margins and dry-sometimes shot-hole centers. Older spots often have reddish margins. Spots may be surrounded with a yellow zone or halo.	Sanitation; protective bactericide sprays for commercial growers.
	Brown Rot (<i>Monilinia</i>)	Twig blight; canker; fruit rot develops as fruits mature.	Protective fungicide sprays. Sanitation.
	Peach Tree Short Life	Dieback of limbs. Poor growth.	Tree removal. Fumigation.
	Root-Knot Nematode (<i>Meloidogyne</i>)	Trees, unthrifty, poor growth; roots have galls.	See AL Pest Management Handbook.
	Twig Blight (<i>Phomopsis</i>)	Twig dieback with brown-black cankers (Some cankers have a zonate pattern.)	Sanitation.
Pear	Entomosporium Leaf Spot (<i>Fabraea</i>)	Roughly circular brownish-black spots (1/8 inch wide).	Sanitation. Spray schedule as described in the AL Pest Management Handbook.
	Fire blight (<i>Erwinia amylovora</i>)	Black discoloration dieback; often infection begins in	Sanitation. See ANR-542 or AL Pest Management

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		the blossoms and moves down into the branch. Blighted branch tips often curve downward. Sunken cankers develop on branches.	Handbook.
	Frogeye Leaf Spot (<i>Botryosphaeria</i>)	Brown, circular spots with dark borders.	Sanitation. See the AL Pest Management Handbook.
	Rust (<i>Gymnosporangium</i> spp.)	Bright yellow spots may occur on pear leaves. Tiny black specks may be seen in upper leaf surface spots; orange/white powdery ‘cups’ may be seen in spots on lower leaf surface.	If practical, remove nearby junipers.
Pear, Callery	Fireblight (<i>Erwinia amylovora</i>)	Flower blight, leaf blight dieback, cankers.	Pruning 10-16” beyond canker edges.
Peanut	Rhizoctonia Seedling Disease & Root Rot	Dry, brown lesions on roots and stems.	Soil or seed treatment before planting.
Pea, Southern	Anthracnose (<i>Colletotrichum</i>)	Gray, brown circular-oval spots on leaves and stems.	See the Alabama Pest Management Handbook.
	Fusarium Crown Rot	Lower stems & crowns become reddish-brown and dry rotted.	Sanitation; crop rotation.
	Rhizoctonia Seedling Disease; Lower Stem Rot	Medium-dark brown blotches or elongated lesions on lower stem near the soil line.	Fungicide treatment prior to planting.
Pecan	Scab (<i>Cladosporium</i>)	Dark olive-green to brown to dark brown slightly raised spots on leaves and nut shucks. Defoliation where disease is severe.	See the AL Pest management Handbook. Sanitation.
Peony	Anthracnose (<i>Colletotrichum</i>)	Circular, light brown spots (2-3mm diameter) on leaves.	Sanitation. See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Peony Ringspot Virus	Concentric bands of alternating dark & light green on leaves; small necrotic spots; plants are not dwarfed.	Remove damaged plants.
Pepper	Bacterial Leaf Spot (<i>Xanthomonas</i>)	Dark, angular green-black, water-soaked spots (1/8-1/4 inch diameter); older spots become dry and white; leaf drop.	Sanitation; protective sprays.
	Bacterial Wilt (<i>Ralstonia solanacearum</i>)	Lower leaves wilt, whole plant wilts.	Sanitation; rotate away from tomato, pepper, eggplant, & Irish potato.
	Cucumber Mosaic Virus	Leaves are malformed and distorted, sometimes shoestring-shaped. Leaves sometimes show yellow patterns.	Sanitation. Control aphids.
	Fusarium Crown Rot	Dry decay of stem at soil line.	Remove and destroy damaged plants. See Ed Sikora for questions.
	Fusarium Wilt	Yellowing and wilt of lower foliage which slowly progresses upward.	Crop rotation away from pepper and other solanaceous crops.
	<i>Phytophthora capsici</i> Crown/Root Rot	Brown lesions on lower stems, crowns mostly; some root rot.	Sanitation; Ridomil.
	Pythium Root Rot	Brown, water-soaked lesions.	Sanitation; Improved soil drainage; See AL Pest Management Handbook.
Persimmon	Phomopsis Dieback	Branch dieback. Cankers form at end of dieback.	Prune out dieback. Make cuts 3-4 inches beyond dieback area.
Persimmon, Japanese	Anthracnose (<i>Colletotrichum</i>)	Circular or irregularly shaped brown leaf spots.	Clean up all fallen leaves in fall; Cleary's 3336 or Halt may be

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			applied as a protective treatment.
	Botryosphaeria Twig Canker	Small twigs develop sunken, cracked spots.	Prune out dieback. Make cuts 3-4 inches beyond end of dieback area.
Petunia	Phytophthora Aerial Blight	Brown lesions on leaves and stems; dieback.	See the AL Pest Management Handbook.
	Phytophthora Root Rot	Roots brown, wet, easily pulled apart.	Sanitation. Avoid wet soils.
	Pythium Root Rot	Wilt, lower leaves turn yellow, plant collapse; a wet, light brown root rot.	Remove dying plants; See the AL Pest Management Handbook.
Phlox	Alternaria Leaf Spot	Dark brown irregularly-shaped spots.	Sanitation; Cleary's 3336 or Domain or an ornamental benomyl product.
Photinia	Entomosporium Leaf Spot	Red-black circular-irregularly shaped spots which often coalesce; leaf drop.	Sanitation. Protective fungicide sprays.
Pine	Needlecast Diseases	Older needles turn brown, drop.	See AL Pest Management Handbook.
	Needle Rust (<i>Coleosporium</i>)	Cream-orange-colored pustules up to 1/8" or 2-3 mm in size occur on needles only of pine.	Remove asters from the area.
	Pitch Canker (<i>Fusarium</i>)	Sunken, cracked lesions on branches, trunks where resin is common.	Sanitation. See the AL Pest Management Handbook for comments.
Pine, Japanese Black Pine	Needlecast (<i>Ploioderma</i> spp.)	Older needles turn brown, drop.	See AL Pesticide Handbook.
Pine, Loblolly	Coleosporium Needle Blight	Needles develop while blisters.	Remove alternate host - asters from the area.
	Fusiform Rust	Elongated, sunken cankers develop on trunk and branches;	Remove oak trees in immediate area.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		orange aeciospores wet masses develop in spring on canker surface. Eventually tree will die back.	
	Rhizosphaeria Needle Blight	Needle blight and drop occurs. Needles develop tiny brown spots in a row.	See the AL Pest Management Handbook.
Plum	Bacterial Leaf Spot (<i>Xanthomonas</i>)	See Peach Bacterial Leaf Spot.	See Peach.
	Black Knot (<i>Plowrightia</i>)	Rough elongated branch galls which begin as greenish swellings but soon develop into hard black elongated branch swellings or galls.	Sanitation; protective sprays of Captan in early spring.
	Botryosphaeria Canker	Dry, sunken, brown lesions that are often cracked around the edges; dieback.	Sanitation.
	Brown Rot (<i>Monilinia fructicola</i>)	A wet, brown discoloration occurs on fruit; a gray mold may develop; small twig cankers.	See the AL Pest Management Handbook.
Potato, Irish	Bacterial Wilt (<i>Ralstonia solanacearum</i>)	Green plants wilt suddenly. Vascular system is brown in the lower stem area. Outer vascular ring of tubers may be discolored.	Rotation away from potato, peppers, tomato, and eggplant for several (4-5 years) years or fumigation.
	Bacterial Soft Rot (<i>Pectobacterium carotovora</i> pv. carotovora)	Irregular, surface, black, wet rot areas on tuber.	Sanitation; avoid wounds/injuries.
	Blackleg (<i>Erwinia</i>)	Large areas of stems become black and water soaked. Plants wilt and die. Tubers are also rotted.	Use disease-free seed pieces; treat seed pieces; clean equipment with solutions of bleach or formaldehyde or allow for good suberization before planting in

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			well-drained areas; crop rotation 2-3 years; control seed corn maggots with insecticides. See the AL Pest Management Handbook.
	Early Blight (<i>Alternaria</i>)	Black, zonate, roughly circular or oval spots (4-10 mm or ¼-½ inch diameter). Spots may also develop on stems & tubers.	Follow spray schedule recommended.
	Fusarium Surface Rot	Dry, dark brown, irregular rot areas on tuber surface.	Sanitation; avoid wounds/injuries.
	Late Blight (<i>Phytophthora infestans</i>)	Brown blotches on leaves and stems; eventually whole leaves and stem sections become severely decayed.	See the AL Pest Management Handbook. See Ed Sikora for commercial plantings.
	Rhizoctonia Stem Rot	Lower stem shows sunken, brown rotted areas.	Sanitation; See AL Pest Management Handbook.
	Scab (<i>Streptomyces scabies</i>)	Slightly raised, scabby circular or circular-irregular lesions on surface of tubers.	It may help to keep soil slightly acid; crop rotation; see Ed Sikora.
	Silver Scurf (<i>Helminthosporium</i>)	Light gray, hard, irregular bodies on tuber surface.	Sanitation.
Privet	Cercospora Leaf Spot	Irregularly-shaped, dark brown leaf spots.	Sanitation of fallen leaves; See the AL Pest Management Handbook.
Rhododendron	Cercospora Leaf Spot	Circular-irregular brown leaf spots.	Sanitation. See the AL Pest Management Handbook.
	Exobasidium Gall	Fleshy, swellings of leaves, flowers. Swellings initially are	Sanitation; Bayleton in some circumstances. See AL Pest

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		green but become white or pink when spore production occurs.	Management Handbook.
	Phytophthora Root Rot	Dieback; roots become water-soaked, rotted, brown.	Remove damaged plants; in a commercial situation, a protective drench treatment may be applied. Reduce water levels in the area.
Rose	Armillaria Root Rot	Dieback of rose; dry, dark brown root rot. See ANR-907.	Remove damaged plants and all roots.
	Black Spot (<i>Diplocarpon</i>)	Black spots (1/8-1/4 inch diameter, 4-8 mm) with feathery margins.	Follow a regular spray schedule; sanitation.
	Botrytis Blossom Blight	Blossoms develop small-large brown blotches. A gray delicate webbing of the fungal mycelium and spores may develop over blotches.	Sanitation. Protective fungicide sprays.
	Brand Canker (<i>Coniothyrium</i>)	Gray, cracked, irregular canker on canes.	Sanitation; fungicides that are recommended for black spot will help.
	Downy Mildew (<i>Peronospora</i>)	Water-soaked, irregular-shaped leaf spots become dark black with age.	See the AL Pest Management Handbook.
	Mosaic	Distinct pattern of dark green & light green or yellow on leaves.	Control weeds; control insects; remove diseased plants.
	Powdery Mildew (<i>Sphaerotheca</i>)	White powdery patches on foliage; some leaf deformity may occur on new growth.	See the AL Pest Management Handbook.
	Tobacco Ringspot Virus	Plant stunted, mottled.	Sanitation; Control nematodes.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Rosemary	Bacterial Leaf Spot & Blight	Black, wet, angular spots.	Sanitation.
Ryegrass, Perennial	Foliar Anthracnose (<i>Colletotrichum</i>)	Light brown/medium brown spots/blotches.	----
	Rust (<i>Puccinia</i>)	Small yellow to red to brown spots/blotches.	Frequent mowing or harvest.
Schefflera, Dwarf	Anthracnose	Whitish or light brown, circular-irregularly shaped spots.	Sanitation of fallen leaves. See the AL Pest Management.
Sedum	Colletotrichum Leaf Spot	Circular-oval, brown leaf spots.	Sanitation; Cleary's 3336 or Halt.
Smilex, Greenbriar	Cercospora Leaf Spot	Medium-brown, irregular-shaped leaf spots.	No control.
Snapdragon	Pythium Root Rot	Roots become light brown and wet rotted. Eventually, roots become dry. Dieback occurs.	Sanitation. Reduce water levels in the area. See the AL Pest Management Handbook.
Spirea	Cylindrosporium Leaf Spot	Brown, irregular-circular spots.	Sanitation.
	Powdery Mildew	White, powdery dusting on surface of leaves; some leaf distortion; leaf blight.	Sanitation; Cleary's 3336.
Spirea, Bridle Wreath	Armillaria Root Rot	Dieback; thin, white mycelial layer just under bark, appressed to wood. Black, thread-like structures may be present above or below surface of root or crown. Brown mushrooms may be present.	Removal of shrub and roots; maintain healthy shrubs.
Squash	Choanephora Wet Rot	Large sections of fruit become wet rotted sometimes with black spore structures covering the fruit surface.	Decrease humidity.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Mosaic Virus	Distinct pattern of dark green and light green or yellow on leaves.	Control weeds; control insects; remove diseased plants; do not save seed.
	Pythium Seeding Disease	Roots become brown and water soaked.	----
Squash, Yellow	Watermelon Mosaic Virus	Patterned patches of light and darker green or yellow pigmentation on leaves.	Sanitation; control aphids.
St. Augustine	Brown Patch (<i>Rhizoctonia</i>)	Leaf blades develop brown blotches; whole leaves become browned.	See the AL Pest Management Handbook.
	Gray Leaf Spot	Gray-brown spots on foliage. Disease may be severe.	See ANR-492 or Pest Management Handbook.
	Slime Mold	An area of grass may be covered by a thin mass of a translucent, watery substance of varying color. As conditions dry out, the watery mass becomes powdery and often black colored with spores. Both growth forms are on the grass blade surface only. See ANR-1117.	Wash off. Avoid wet conditions.
	Take-All (<i>Gaeumannomyces</i>)	Individual plants yellow and areas thin out. Black lesions occurs on roots, stolons.	See PP-312 or AL Pest Management Handbook.
Strawberry	Angular Leaf Spot (<i>Xanthomonas</i>)	Angular, black spots in interveinal areas. Spotting may be severe.	See the AL Pest Management Handbook.
	Anthraxnose-Crown Rot (<i>Colletotrichum</i>)	Lower stems and crowns become brown, and rotted.	Use healthy transplants. Sanitation. See Ed Sikora.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
	Mycosphaerella Leaf Spot	Small (1/8 inch) red-brown circular spots.	See the AL Pest Management Handbook.
	Phytophthora Crown & Root Rot	Wilt, yellowing of lower leaves; dieback; wet, brown rot or crown and roots.	Sanitation; See the AL Pest Management Handbook.
Sweet Potato	Black Rot (<i>Ceratocystis fimbriata</i>)	Black lesions on root surface which extend to but not beyond the vascular ring. Secondary bacteria will sometimes invade tissues and cause rotting beyond the vascular system.	Sanitation. See the AL Pest Management Handbook.
Sycamore	Anthrachnose (<i>Apiognomonina</i>)	Gray-brown blotches on leaves; sometimes blotches follow leaf veins or leaf margins.	Sanitation. Protective fungicide sprays for small trees.
	Scorch (<i>Xylella</i>)	Leaves become brown and dried at the margins; dieback during summer months.	Tree pruning or removal.
Tomato	Bacterial Canker (<i>Clavibacter</i>)	Dark brown, black, slightly sunken, large lesions (1-4 cm or larger) develop on stems.	See the AL Pest Management Handbook.
	Bacterial Spot (<i>Xanthomonas</i>)	Small, dark green or black, wet-looking, angular spots.	Sanitation. Protective sprays.
	Bacterial Wilt (<i>Ralstonia solanacearum</i>)	Plants wilt suddenly while still green. Lower stem areas are black (decayed) beneath the green outer layers.	Rotate the area away from tomato, potato, eggplant and pepper for 4-5 years or fumigate the soil.
	Blossom End Rot	Black, hard, circular, slightly sunken spots at stem end of fruit.	See AL Pest Management Handbook.
	Buckeye Rot (<i>Phytophthora nicotiana</i>)	On fruit, spots are pale brown with concentric rings.	Keep fruit off the soil. See AL Pest Management

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
			Handbook.
	Cucumber Mosaic Virus	Plants become stunted with misshaped leaves (often strap shaped) that may be puckered and show a mosaic pattern.	Sanitation; Aphid control.
	Early Blight (<i>Alternaria</i>)	Black or brown spots (¼-½ inch diameter or 4-10 mm diameter) on leaves, stems, fruit. Spots often have a concentric pattern.	Fungicide sprays; sanitation.
	Fusarium Crown Rot	Lower stem becomes reddish-brown and dry rotted; plant wilts; yellowing of lower leaves; dieback.	Sanitation; rotate away from tomato; resistant varieties.
	Fusarium Wilt	Lower leaves turn yellow and wilt. Yellowing progresses up the plant.	Crop rotates away from tomatoes 10-16 years on resistant varieties.
	Late Blight (<i>Phytophthora infestans</i>)	See Irish Potato.	See the AL Pest Management Handbook.
	Leaf Mold (<i>Fulvia fulva</i>)	Gray fungal growth in blotches on leaves.	See AL Pest Management Handbook.
	Pythium Root Rot	Roots develop brown-colored, soft rot.	Sanitation. Do not irrigate excessively; provide well-drained soil area. See AL Pest Management Handbook
	Rhizoctonia Collar Rot	A dry, brown rot develops on lower stem at soil line.	Plant removal. See Ed Sikora.
	Root-Knot Nematode (<i>Meloidogyne</i>)	Irregular root galls which involve the whole root circumference. Gall size varies from	Soil fumigation (commercial) or rotation to dwarf French marigold or pasture grasses.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		small (2-3 mm diameter) to large (1-1.5 cm diameter).	
	Septoria Leaf Spot	Small (¼ inch diameter) roughly circular gray-brown spots.	Protective fungicide sprays. See AL Pest Management Handbook.
	Southern Blight (<i>Sclerotium rolfsii</i>)	A wet, brown rot develops at the soil line; with humid conditions, a thick white mold may develop, often with hard, brown, spherical sclerotia (over-wintering bodies) present.	Sanitation; see Disease Note ANR-863.
	Timber Rot (<i>Sclerotinia</i>)	Lower stem near soil (crown) becomes brown and decayed. Plant wilts and eventually collapses.	Remove damaged plants. Consult Ed Sikora.
	Tomato Pith Necrosis (<i>Pseudomonas corrugate</i>)	Sections of stem may show a black discoloration; plant parts above the lesion wilt and die; inside the stem, hollow 'chambers' are delimited by 'strands' of pith tissue.	Sanitation.
	Tomato Spotted Wilt Virus	Wilt, ring spots, bronzed leaf surface areas, and wilt may develop. Fruit & new growth is stunted.	Remove plants. Control thrips.
Torenia	Powdery Mildew	A white, powdery dusting develops on leaves and stems; blight follows.	Sanitation; Cleary's 3336.
Tulip Poplar	Cristulariella Leaf Spot	Leaf spots are often large and zonate.	Collect fallen leaves in the fall.
Vegetables	Seedling Diseases (<i>Fusarium</i> , <i>Rhizoctonia</i> , <i>Pythium</i>)	Seedlings fall over at the soil line; small stems are weak and	Soil treatments (in some situations) or seed treatments before

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		limp; brown discoloration may be evident.	planting. See Ed Sikora.
Vinca, Annual Periwinkle (<i>Catharanthus</i>)	Alternaria Leaf Spot	Circular-angular dark brown-black spots.	Sanitation. Cleary's 3336 or Domain should give protection control.
	Botrytis Blight	Brown-gray leaf spots & blight	Sanitation. Make adjustments to decrease humidity levels. Cleary's 3336 or Domain should give protective control.
	Phytophthora Stem Rot	Sections of stems become brown, wet, and collapsed, shriveled. Some water-soaked gray leaf spots. Spots later turn brown.	Sanitation. In some cases Aliette WDG may be used as a protective treatment. See AL Pest Management Handbook and ANR-1023.
	Rhizoctonia Aerial Blight	Dark brown dried cankers on stems. Tops dieback.	Sanitation. Cleary's 3336 may be used in some situations. See the AL Pest Management Handbook.
Violet	Cercospora Leaf Spot	Irregularly shaped brown-black spots.	Sanitation. Cleary's 3336 or Halt may be applied as protective treatments.
Watermelon	Anthrachnose (<i>Colletotrichum</i>)	Foliage spots begin as small yellow or water soaked irregular areas which become larger and eventually black. Whole leaf areas may become covered with the coalesced spotting/ blotches. On the fruit, small water soaked spots become dark green and then brown-black sunken spots (1/4-2 inches x 1/8 inch deep). With high	See the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		humidity the spots may become covered with gelatinous masses of orange spores.	
	Fusarium Root Rot	Roots become dry, rotted; wilt, dieback.	Crop rotates for 10-16 years.
	Fusarium Wilt (<i>F. oxysporum</i>)	Lower leaves turn yellow and wilt slowly develops.	Rotation; resistant varieties.
	Gummy Stem (<i>Mycosphaerella</i>)	Brown, black spots occur interveinally in the lobes of leaves. Usually older leaves near the center of the bill are affected initially. As the season progresses, elongated, water soaked, light-brown to gray lesions occur on vines, often near the crown of the plant. As lesions age, cracking occurs and a reddish-yellow gummy substance oozes around the crack. Plants gradually yellow and turn brown.	See the Alabama Pest Management Handbook.
	Pythium Root Rot	Roots become brown-colored and soft rotted.	Sanitation. Avoid wet conditions, especially with poorly draining soil. Contact Ed Sikora if you have questions.
	Rhizoctonia Seedling Blight	Brown lesions on lower stems; seedlings collapse.	Sanitation. Crop rotation.
	Suspect Ozone Damage	Older leaves with whitish, nitted interveinal areas of upper leaf surfaces. Old damage may become black.	---

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
Wheat	Bipolaris Leaf Spot	Brown irregular spots.	Fungicides in some situations; refer to Bill Gazaway.
	Black Chaff (<i>Xanthomonas</i>)	Gray-brown-black discoloration of seed heads and leaves. Discoloration starts at tip of seed.	Use healthy seed.
	Fusarium Scab (Head Blight)	Head blight with a pinkish cast due to presence of spores.	Use healthy seed. Deep plowing, crop rotation of one year. Some cultivar differences in disease susceptibility have been noted.
	Leaf Rust (<i>Puccinia</i> sp.)	Orange powdery spots.	Fungicides in some situations; refer to Bill Gazaway.
	Powdery Mildew (<i>Erysiphe</i> sp.)	Buff-colored spots and patches.	Fungicides in some situations; refer to Bill Gazaway.
	Septoria Glume Blotch/Leaf Spot	Gray-brown patches of discoloration on glumes; gray-brown leaf spots (1/4-1/3 inch diameter).	---
	Take-All (<i>Gaeumannomyces</i> sp.)	Stem at the soil-line become black. Plants die.	Rotation.
Wild Rocket (Rugula)	Tan Spot (<i>Drechslera tritrici-repentis</i>)	Circular to oval brown leaf spots.	---
	Anthracnose (<i>Colletotrichum</i>)	Round to angular, brown leaf spots.	Sanitation.
Willow	Bacterial Leaf Spot (<i>Pseudomonad syringae</i> pv. <i>syringae</i> & <i>P. cichorii</i>)	Angular, black, wet leaf spots.	Sanitation.
	Leaf Rust (<i>Melampsora</i>)	Yellow orange pustules on leaves. (See weeping willow below.)	Sanitation.
Willow, Weeping	Armillaria Root Rot	Dieback; thin, white mycelial layer just under bark, appressed	Removal of tree and roots; maintain healthy trees.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		to wood. Black, thread-like structures may be present above or below surface of root or crown. Brown mushrooms may be present.	
	Leaf Rust (<i>Melampsora</i>)	Small yellow spots that become orange with powdery rust spores and then later become brown spots.	Sanitation of fallen leaves. Alternate hosts may be fir, saxifrage, hemlock, current, or gooseberry.
Zinnia	Bacterial Leaf Spot	Small, angular, dark gray or black leaf spots. Spot edges may appear wet.	Remove all damaged plants. Do not water over-head. Phyton 27 may be used, if desired. See the AL Pest Management Handbook.
	Rhizoctonia Crown Rot	A dark brown, dry rot develops on the lower stem.	Remove damaged plants. See the AL Pest Management Handbook for commercial situations.
Zoysia	Brown Patch [Large Patch] (<i>Rhizoctonia</i>)	See Centipede.	See Centipede.
	Dollar Spot (<i>Sclerotinia</i>)	Yellow or white spots the size of a silver dollar develop. Spots may coalesce.	See ANR-493.
	Ring Nematode (<i>Criconeoides</i> sp.)	Areas thin; dieback; poor root system.	See AL Pest Management Handbook.
	Rust (<i>Puccinia</i>)	Areas of turf become yellow-brown with tiny spots of orange powdery spores.	See the AL Pest Management Handbook or ANR-621.
	Suspect Winter Damage	Large turf areas do not green up in the spring.	---
	Take-All Patch (<i>Gaeumannomyces</i> sp.)	Individual plants turn yellow and die. Roots show dark	See PP-312 and the AL Pest Management Handbook.

<u>Plant</u>	<u>Disease</u>	<u>Description</u>	<u>Control</u>
		brown or black lesions.	

Lab Notes

As we move into our busiest time of year, please take care when filling out the diagnostic sheet or when advising clients in filling out the information. Soil analysis (pH, minerals) results will come back to the client more quickly if sent directly. If the client would like us to forward soil to Soil Test Lab--if we believe a soil problem exists--he/she must indicate yes on the form. We will not forward soil when a charge is involved unless the client gives approval. Also, please indicate whether the sample is a client charge or an educational (ACES) charge by checking the appropriate blank in the box at the top of the form. Please give your area code with your phone number! Failure to include the above information may delay our response. Thanks!

Soil Samples For Nematode Analysis

Please be sure you are using the current soil nematode sample submission form which indicates a service charge of \$10.

Insects

Usually insects for identification should be sent in isopropyl alcohol. If you send live insects, please clearly indicate this on your paperwork.