

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid* 2F

EPA Reg. No. 62719-442

For Control of Lepidoptera Larvae Infesting Corn (Field, Sweet, Seed)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
European corn borer southwestern corn borer	4 – 8 (0.06 – 0.12 lb ai/acre)	Apply at first sign of egg hatch or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities. Direct application at the whorl for early season (first generation) infestations. Apply as broadcast or multi-nozzle over the row application to mid- and late-season infestations.	<ul style="list-style-type: none"> • Do not apply more than 16 fl oz per acre per application or 64 fl oz of Intrepid 2F (1.0 lb ai) per acre per season. • Pre-harvest Interval-Field Corn: Do not harvest within 21 days of application. • Pre-harvest Interval-Sweet Corn: Do not harvest within 3 days of application for ears and/or green chop (forage) and within 21 days of application for dry fodder. • See Rotational Crop Restrictions below.
true armyworm western bean cutworm		Apply at first sign of egg hatch (field corn), feeding damage (sweet corn), or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities. Under heavy infestations, continuous moth flights, or rapid crop growth and development, reapply at 5-10 day re-treatment interval.	

Specific Use Directions-Field Corn:

Ground Application: Apply a minimum of 5 gallons per acre by conventional ground equipment to young crop or small plants. Higher carrier volumes may be required to provide thorough coverage to larger, more mature crop. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 5 gallons per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

Spray Adjuvants: The use of Latron CS-7* spreader/binder or similar spreader/binder is recommended to maximize coverage and retention of the spray material.

Specific Use Directions-Sweet Corn:

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop or small plants. Apply a minimum of 20 gallons per acre after initiation of tasseling. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre.

Spray Adjuvants: The use of Latron CS-7 spreader/binder or similar spreader/binder is recommended to maximize coverage and retention of the spray material.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days.

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

F3B / Intrepid 2F / Sec 3 Supp Corn FPL / 10-04-02

D06-846-012

EPA-accepted 09/26/02

Initial printing.