



IPM-0429

Grain Sorghum

Insect, Disease, and Weed Control
Recommendations for 2023



INSECT CONTROL (2023 UPDATES)

Grain sorghum is attacked by a number of insect pests, including a new pest, the sugarcane aphid, *Melanaphis sacchari*. Sugarcane aphid is now the key pest of sorghum. DO NOT plant sorghum if you cannot afford to scout the crop weekly and apply at least two applications of insecticide for this and other sorghum pests.

For more information on sugarcane aphid, see Sugarcane Aphid: A New Sorghum Pest in North America (<http://betteryield.agrilife.org/publications/>). Watch the blog on Alabamacrops.com (or download the app) to get the most up-to-date information on sorghum insects and their management including any Section 18 labels for additional insecticides that are effective against sugarcane aphids. Or subscribe to the Alabama IPM Newsletter by visiting www.aces.edu/IPMCommunicator.

Many insecticides are registered for use on grain sorghum for control of whorl worms, headworms, grasshoppers, and sorghum midge. Unfortunately, many of these insecticides disrupt the community of beneficial insects that help control sugarcane aphid. Using these broad spectrum insecticides can flare aphids (cause the number of aphids to increase rapidly). If you are planting a variety that is susceptible to sugarcane aphid (and most of them are), AVOID USING BROAD SPECTRUM INSECTICIDES IF AT ALL POSSIBLE. Insecticides that are more selective and less harsh on natural enemies are highlighted in Table 3.

“Sorghum Insects and Their Management” (http://www.caes.uga.edu/publications/pubDetail.cfm?pk_id=7797) by David Buntin, University of Georgia, provides excellent information on the biology and management of sorghum pests, other than sugarcane aphid. It should be required reading if you plan to grow sorghum, because it provides valuable tips on insect action thresholds as well as on how to reduce risk of insect infestation.

The Sorghum Checkoff Program provides many resources on sorghum pest management, including the latest list of varieties with resistance to sugarcane aphid. See <http://sorghumcheckoff.com/pest-management/>.

Insect Pest Management Tips for Grain Sorghum

- Plant early if you can. The later you plant, the more insect problems you encounter.
- Choose a variety that is resistant to sugarcane aphid.
- Use an insecticide seed treatment to protect the sorghum for a few weeks from various soil insect pests as well as the sugarcane aphid.
- Scout the crop at least once a week.

Organic Insecticides That Can Be Applied to Grain Sorghum

Organic producers may want to consider the following insecticides, most, if not all, of which are OMRI approved. Read the insecticide label to make sure it meets your needs. The following products contain azadirachtin: Azera, Azatrol, Neemix 4.5, and Ecozin Plus. The following products contain *Bacillus thuringiensis*: Biobit HP (subsp. *kurstaki* strain ABTS- 351), Dipel DF (subsp. *kurstaki* strain ABTS-351), Javelin WG (subsp. *kurstaki* strain SA-11), and Xentari (subsp. *aizawai* strain ABTS- 1857). Surround WP contains kaolin. Microthiol Disperss and Yellow Jacket Special Dusting Sulfur contain sulfur. Gemstar LC contains an insecticidal virus. Venerate XC Bioinsecticide contains *Burkholderia* spp. strain A396. Pest Out contains cottonseed, clove, and garlic oils. Ecotec contains rosemary and peppermint oils. Biolink Insect and Bird Repellant contains garlic. Entrust and Entrust SC contain spinosad. Other formulations of spinosad can be found. Mycotrol ESO and BotaniGard ES contain *Beauveria bassiana* strain GHA. Grandevlo contains *Chromobacterium subtsugae* strain PRAA4-1.

PyGanic Crop Protection EC 1.4 and PyGanic Crop Protection EC 5.0 insecticides are OMRI approved and contain natural pyrethrins. Other insecticides contain pyrethrins, but also contain piperonyl butoxide and are not considered organic. Gemstar LC contains polyhedral inclusion bodies of the nuclear polyhedrosis virus of corn earworm. Other organic insecticides may be available. Be sure to check the labels to see if they meet the requirements for your cropping system.

Forage Sorghum + Sudangrass

For information on forage sorghum, sudangrass, and other temporary summer grazing, see Temporary Grazing: Temporary Summer Grazing Insect Control, part of the Georgia Pest Management Handbook at <http://www.caes.uga.edu/departments/entomology/extension/pest-management-handbook.html> or the Pasture and Forage Section of the Louisiana Insect Pest Management Guide at http://www.lsuagcenter.com/portals/communications/publications/management_guides/insect_guide.

Table 1. Seed Treatments for Control of Seedling Insect Pests in Grain Sorghum, 2016

| Insecticide and Formulation | Rate |
|---|-------------------------------------|
| clothianidin | |
| PONCHO 600, NIPSIT INSIDE | 5.1 to 6.4 fl. oz. per 100 lb. seed |
| clothianidin + <i>Bacillus firmus</i> | |
| PONCHO VOTiVO | 6.13 fl. oz. per 100 lb. seed |
| imidacloprid | |
| AXCESS, MACHO 600 ST, DYNASHIELD IMIDACLOPRID 5, NITRO SHIELD, ATTENDANT 480 FS, NITRO SHIELD IV, GAUCHO 600, IMIDACLOPRID 5, SENATOR 600 FS, SHARDA IMIDACLOPRID 5SC, RESONATE 600ST | 6.4 fl. oz. per 100 lb. seed |
| NITROSHIELD IV | 8 fl. oz. per 100 lb. seed |
| imidacloprid + metalaxyd | |
| CONCUR | 5 oz. per 100 lb. seed |
| imidacloprid + metalaxyd+carboxin | |
| LATITUDE | 5 oz. per 100 lb. seed |
| thiamethoxam | |
| CRUISER 5FS | 5.1 to 7.6 fl. oz. per 100 lb. seed |

Table 2. Examples of Broad-Spectrum At-Planting Insecticides for Insect Pests of Sorghum Seeds and Seedlings

| Insecticide (Trade Names) | Rates of Formulated Product | Pests Controlled Include |
|--|-----------------------------|--|
| chlorpyrifos (Lorsban 15G) ¹ | 8 oz./1000 row ft. | Southern corn rootworm, chinch bugs, ants, cutworm, lesser cornstalk borer |
| phorate (THIMET 20-G SMARTBOX or LOCKBOX or EZLOAD) | 4.5–6 oz./1000 row ft. | greenbugs |
| terbufos (Counter 20G Lock'n'Load) ¹ | 4.5–6 oz. 1000 row ft. | Southern corn rootworm, wireworms, white grubs, aphids |

See the insecticide label for specific use instructions, including whether product is to be applied in-furrow, as a T-band, or broadcast.

¹ Other trade names available see Table 4.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control

| Insecticide and Formulation | Acres per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|--|--------------------------|--------------------------------|--------------------------------|--|--|
| APHIDS (SUGARCANE APHID, YELLOW SUGARCANE APHID, CORN LEAF APHID, GREENBUG) | | | | | |
| flupyradifurone SIVANTO PRIME | 18.3–32 | 4–7 fl. oz. | 0.05–0.09 | 7 (forage) 21 (grain) | Controls sugarcane aphid. |
| chlorpyrifos LORSBAN ADVANCED (RESTRICTED USE) | 4–8 | 1–2 pt. | 0.47–0.94 | 30 (1 pt. or less) 60 (> 1 pt.) | Do not apply to sweet sorghum; may give helpful control under light population pressure. |
| malathion CHEMINOVA MALATHION 57% Other trade names ¹ | 53 | 1.5 pt. | 0.93 | 7 | May give helpful control for sugarcane aphid on the head at the end of the season. |
| sulfoxaflor TRANSFORM WG | 5–10 | 0.75–1.5 fl. oz. | 0.023–0.047 | 7 (forage) 14 (grain) | Transform WG was granted a Section 3 label in 2019. |
| WHORL WORMS + HEADWORMS (ARMYWORMS, FALL AND TRUE; CORN EARWORM; SORGHUM WEBWORM) | | | | | |
| alpha-cypermethrin FASTAC EC Other trade names ¹ | 34–71 | 1.8–3.8 fl.oz. | 0.012–0.025 | 14 (grain, stover) 45 (forage) | Fastac EC is a RESTRICTED USE pesticide. |
| beta-cyfluthrin BAYTHROID XL | 46–98 | 1.3–2.8 fl.oz. | 0.01–0.022 | 14 | Baythroid XL is a RESTRICTED USE pesticide. Use highest rate for fall armyworms. |
| carbaryl SEVIN XLR PLUS Other trade names ¹ | 2–4 | 1–2 qt. | 1–2 | 14 (silage, grazing) 21 (fodder, grain) | |
| chlorantraniliprole PREVATHON Other trade names ¹ | 6–9 | 14–20 fl.oz. | 0.05–0.07 | 1 | Does not flare sugarcane aphid. |
| chlorpyrifos LORSBAN ADVANCED LORSBAN 75WG Other trade names ¹ | 4–8 0.75–1.5 A/lb. | 1–2 pt. 0.67–1.33 lb. | 0.47–0.93 0.5–1 | 30 (1 pt. or less) 60 (> 1 pt.) | Use on true armyworms only. See label for detailed instructions. Lorsban Advanced is a RESTRICTED USE pesticide. Lorsban 75WG is not. |
| chlorpyrifos + lambda-cyhalothrin COBALT ADVANCED | 3.4–12 | 11–38 fl.oz. | 0.21–0.74 + 0.011–0.038 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt Advanced is a RESTRICTED USE pesticide. |
| chlorpyrifos + gamma-cyhalothrin COBALT Other trade names ¹ | 3.4–10 | 13–38 fl.oz. | 0.25–0.74 + 0.004–0.013 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt is a RESTRICTED USE pesticide. |

¹ See Table 4 for other trade names.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control (cont.)

| Insecticide and Formulation | Acres per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|---|------------------|--------------------------------|--------------------------------|---|--|
| WHORL WORMS + HEADWORMS (ARMYWORMS, FALL AND TRUE; CORN EARWORM; SORGHUM WEBWORM (cont.)) | | | | | |
| chlorpyrifos + zeta-cypermethrin STALLION | 11–26 | 5–11.75 fl.oz. | 0.11–0.25 + 0.01–0.025 | 30 (grain, stover) 45 (forage) | Stallion is a RESTRICTED USE pesticide. |
| cyfluthrin TOMBSTONE Other trade names ¹ | 46–98 | 1.3–2.8 fl.oz. | 0.020–0.044 | 14 | Tombstone is a RESTRICTED USE pesticide. Use high rate for fall armyworm. |
| deltamethrin DELTA GOLD 1.5 EC | 67–128 | 1.0–1.9 fl.oz. | 0.012–0.022 | 14 | Delta Gold is a RESTRICTED USE pesticide. Use higher rate for <i>Spodoptera</i> armyworms. |
| esfenvalerate ASANA XL Other trade names ¹ | 13–22 | 5.8–9.6 fl.oz. | 0.03–0.05 | 21 | Asana is a RESTRICTED USE pesticide. |
| gamma-cyhalothrin DECLARE Other trade names ¹ | 83–125 | 1.02–1.54 fl.oz. | 0.01–0.015 | 30 | Declare is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin WARRIOR II with Zeon Technology Other trade names ¹ | 67–100 | 1.28–1.92 fl.oz. | 0.02–0.03 | 30 | Warrior II is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin + chlorantraniliprole BESIEGE | 13–21 | 6–10 fl.oz. | 0.02–0.03 + 0.04–0.06 | 30 | Besiege is a RESTRICTED USE pesticide. |
| methomyl LANNATE LV Other trade names ¹ | 5–11 | 0.75–1.5 pt. | 0.22–0.45 | 14 | Lannate is a RESTRICTED USE pesticide. |
| methoxyfenozide INTREPID 2F Other trade names ¹ | 13–16 | 8–10 fl.oz. | 0.12–0.16 | 21 (grain or stover) 3 (forage or sweet sorghum stalk) | |
| novaluron DIAMOND 0.83 EC Other trade names ¹ | 10.7–14.2 | 9–12 fl.oz. | 0.06–0.08 | 7 (forage) 14 (grain, stover) | |
| spinosad BLACKHAWK Other trade names ¹ | 4.8–10.6/lb. | 1.5–3.3 oz. | 0.034–0.074 | 21 (grain) 3 (forage, fodder) | Use higher rates for heavier infestations. Time applications to peak egg hatch. Does not flare sugarcane aphid. |
| spinosad + gamma-cyhalothrin CONSERO | 42–64 | 1–1.5 fl.oz. of each product | 0.03–0.046 + 0.01–0.015 | 30 | Consero is a RESTRICTED USE pesticide. It is a co-pack of two insecticides that must be applied together. |
| zeta-cypermethrin MUSTANG MAXX EC INSECTICIDE Other trade names ¹ | 32–73 | 1.76–4 fl.oz. | 0.011–0.025 | 14 (grain, stover) 45 (forage) | Mustang Maxx is a RESTRICTED USE pesticide. |
| CHINCH BUGS | | | | | |
| General Comments: Apply insecticide as a directed spray to the base of plants in at least 15 to 20 gallons of water per acre. See Table 2 for suggested seed treatments and Table 4 for at-planting insecticides for control of chinch bugs. | | | | | |
| alpha-cypermethrin FASTAC EC Other trade names ² | 34–40 | 3.2–3.8 fl.oz. | 0.02–0.025 | 14 (grain, stover) 45 (forage) | Fastac EC is a RESTRICTED USE pesticide. |
| beta-cyfluthrin BAYTHROID XL | 46–64 | 2.0–2.8 fl.oz. | 0.019–0.022 | 14 | Baythroid XL is a RESTRICTED USE pesticide. |

¹ See Table 4 for other trade names.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control (cont.)

| Insecticide and Formulation | Acres Per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|--|--------------------------|--------------------------------|--------------------------------|--|--|
| CHINCH BUGS (cont.) | | | | | |
| carbaryl SEVIN XLR PLUS Other trade names ¹ | 2–4 | 1–2 qt. | 1–2 | 14 (silage, grazing) 21 (grain, fodder) | |
| chlorpyrifos LORSBAN ADVANCED LORSBAN 75 WG Other trade names ¹ | 4–8 0.75–1.5 A/lb. | 1–2 pt. 0.67–1.33 lb. | 0.47–0.93 0.5–1 | 30 (1 pt. or less) 60 (> 1 pt.) | See label for detailed instructions. Lorsban Advanced is a RESTRICTED USE pesticide. Lorsban 75WG is not. |
| chlorpyrifos + lambda-cyhalothrin COBALT ADVANCED | 3–12 | 11–38 fl.oz. | 0.21–0.74 + 0.011–0.013 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt Advanced is a RESTRICTED USE pesticide. |
| chlorpyrifos + gamma-cyhalothrin COBALT Other trade names ¹ | 3.4–10 | 13–38 fl.oz. | 0.25–0.74 + 0.004–0.013 | 30 (26 fl. oz. or less) 60 (> 26 fl. oz.) | Cobalt is a RESTRICTED USE pesticide. |
| chlorpyrifos + zeta-cypermethrin STALLION | 11–14 | 9.25–11.75 fl.oz. | 0.2–0.25 + 0.02–0.025 | 30 (grain, stover) 45 (forage) | Stallion is a RESTRICTED USE pesticide |
| cyfluthrin TOMBSTONE Other trade names ¹ | 46–64 | 2.0–2.8 fl.oz. | 0.038–0.044 | 14 | Tombstone is a RESTRICTED USE pesticide. |
| deltamethrin DELTA GOLD 1.5 EC | 67–98 | 1.3–1.9 fl.oz. | 0.015–0.022 | 14 | Delta Gold is a RESTRICTED USE pesticide. |
| esfenvalerate ASANA XL Other trade names ¹ | 13–22 | 5.8–9.6 fl.oz. | 0.03–0.05 | 21 (harvest) — | Asana is a RESTRICTED USE pesticide. |
| gamma-cyhalothrin DECLARE Other trade names ¹ | 83 | 1.54 fl.oz. | 0.015 | 30 | Declare is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin WARRIOR II with Zeon Technology Other trade names ¹ | 67 | 1.92 fl.oz. | 0.03 | 30 | Warrior II is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin + chlorantraniliprole BESIEGE | 13 | 10 fl.oz. | 0.03 + 0.06 | 30 | Besiege is a RESTRICTED USE pesticide. |
| spinosad + gamma-cyhalothrin CONSERO | 42 | 1.5 fl.oz. of each product | 0.046 + 0.015 | 30 | Consero is a RESTRICTED USE pesticide. It is a co-pack of two insecticides that must be applied together. |
| zeta-cypermethrin MUSTANG MAXX EC INSECTICIDE Other trade names ¹ | 32–40 | 3.2–4 fl.oz. | 0.02–0.025 | 14 (grain, stover) 45 (forage) | Mustang Maxx is a RESTRICTED USE pesticide. |
| CUTWORMS | | | | | |
| See Tables 2 and 4 for suggested seed treatments and broad-spectrum preplant and at-planting insecticides for control of cutworms. | | | | | |
| alpha-cypermethrin FASTAC EC Other trade names ² | 34–98 | 1.3–3.8 fl.oz. | 0.008–0.025 | 14 (grain, stover) 45 (forage) | Fastac EC is a RESTRICTED USE pesticide. |

¹ See Table 4 for other trade names.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control (cont.)

| Insecticide and Formulation | Acres Per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|---|--------------------------|--------------------------------|--------------------------------|--|---|
| CUTWORMS (cont.) | | | | | |
| <i>See Tables 2 and 4 for suggested seed treatments and broad-spectrum preplant and at-planting insecticides for control of cutworms.</i> | | | | | |
| beta-cyfluthrin BAYTHROID XL | 98–128 | 1.0–1.3 fl.oz. | 0.008–0.01 | 14 | Baythroid XL is a RESTRICTED USE pesticide. |
| carbaryl SEVIN XLR PLUS Other trade names ¹ | 2 | 2 qt. | 2 | 21 (grain, fodder) 14 (grazing, silage) | |
| chlorpyrifos LORSBAN ADVANCED LORSBAN 75 WG Other trade names ¹ | 4–8 0.75–1.5 A/lb. | 1–2 pt. 0.67–1.33 lb. | 0.47–0.93 0.5–1 | 30 (1 pt. or less) 60 (> 1 pt.) | See label for detailed instructions. Can be applied preplant, at plant, or preemergence. Lorsban Advanced is a RESTRICTED USE pesticide. Lorsban 75WG is not. |
| chlorpyrifos + lambda-cyhalothrin COBALT ADVANCED | 3–12 | 11–38 fl.oz. | 0.21–0.74 + 0.01–0.038 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt Advanced is a RESTRICTED USE pesticide. |
| chlorpyrifos + gamma-cyhalothrin COBALT Other trade names ¹ | 3.4–10 | 13–38 fl.oz. | 0.25–0.74 + 0.004–0.013 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt is a RESTRICTED USE pesticide. |
| chlorpyrifos + zeta-cypermethrin STALLION | 11–34 | 3.75–11.75 fl.oz. | 0.08–0.25 + 0.008–0.025 | 30 (grain, stover) 45 (forage) | Stallion is a RESTRICTED USE pesticide. |
| cyfluthrin TOMBSTONE Other trade names ¹ | 98–128 | 1.0–1.3 fl.oz. | 0.016–0.020 | 14 | Tombstone is a RESTRICTED USE pesticide. |
| deltamethrin DELTA GOLD 1.5 EC | 85–128 | 1–1.5 | 0.012–0.018 | 14 | Delta Gold is a RESTRICTED USE pesticide. Can be applied pre- or postemergence. |
| esfenvalerate ASANA XL Other trade names ¹ | 13–22 | 5.8–9.6 fl.oz. | 0.03–0.05 | 21 | Asana is a RESTRICTED USE pesticide. |
| gamma-cyhalothrin DECLARE Other trade names ¹ | 125–166 | 0.77–1.02 fl.oz. | 0.0075–0.01 | 30 | Declare is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin WARRIOR II with Zeon Technology Other trade names ¹ | 100–133 | 0.96–1.28 fl.oz. | 0.015–0.020 | 30 | Warrior II is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin + chlorantraniliprole BESIEGE | 21–26 | 5–6 fl.oz. | 0.016–0.02 + 0.03–0.04 | 30 | Besiege is a RESTRICTED USE pesticide. |
| novaluron DIAMOND 0.83 EC Other trade names ¹ | 10.7–14.2 | 9–12 fl. oz. | 0.06–0.08 | 7 (forage) 14 (grain, stover) | |
| spinosad + gamma-cyhalothrin CONSERO | 42–64 | 1–1.5 fl.oz. of each product | 0.03–0.046 + 0.01–0.015 | 30 | Consero is a RESTRICTED USE pesticide. It is a co-pack of two insecticides that must be applied together. |

¹ See Table 4 for other trade names.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control (cont.)

| Insecticide and Formulation | Acres Per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|---|---------------------|--------------------------------|--------------------------------|--|---|
| CUTWORMS (cont.) | | | | | |
| zeta-cypermethrin MUSTANG MAXX EC INSECTICIDE Other trade names ¹ | 32–100 | 1.28–4 fl.oz. | 0.008–0.025 | 14 (grain, stover) 45 (forage) | Mustang Maxx is a RESTRICTED USE pesticide. Can be applied at planting or prior to planting. |
| GRASSHOPPERS | | | | | |
| alpha-cypermethrin FASTAC EC Other trade names ¹ | 32–40 | 3.2–3.9 fl.oz. | 0.017–0.025 | 14 (grain, stover) 45 (forage) | Fastac EC is a RESTRICTED USE pesticide. |
| beta-cyfluthrin BAYTHROID XL | 46–64 | 2.0–2.8 fl.oz. | 0.019–0.022 | 14 | Baythroid XL is a RESTRICTED USE pesticide. |
| carbaryl SEVIN XLR PLUS Other trade names ¹ | 3–8 | 0.5–1.5 qt. | 0.5–1.5 | 14 (silage, grazing) 21 (grain, fodder) | Use lower rate of Sevin for young grasshoppers or sparse vegetation; use higher rate for larger grasshoppers or thicker vegetation. |
| chlorpyrifos LORSBAN ADVANCED LORSBAN 75 WG Other trade names ¹ | 8–16 1.5–3 A/lb. | 0.5–1 pt. 0.33–0.67 lb. | 0.23–0.47 0.25–0.5 | 30 (1 pt. or less) 60 (> 1 pt.) | See label for detailed instructions. Lorsban Advanced is a RESTRICTED USE pesticide. Lorsban 75WG is not. |
| chlorpyrifos + lambda-cyhalothrin COBALT ADVANCED | 10–21 | 6–13 fl.oz. | 0.1–0.25 + 0.006–0.013 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt Advanced is a RESTRICTED USE pesticide. |
| chlorpyrifos + zeta-cypermethrin STALLION | 11–14 | 9.25–11.75 fl.oz. | 0.2–0.25 + 0.02–0.025 | 30 (grain, stover) 45 (forage) | Stallion is a RESTRICTED USE pesticide. |
| chlorpyrifos + gamma-cyhalothrin COBALT Other trade names ¹ | 10–18 | 7–13 fl.oz. | 0.14–0.25 + 0.002–0.004 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt is a RESTRICTED USE pesticide. |
| cyfluthrin TOMBSTONE Other trade names ¹ | 46–64 | 2.0–2.8 fl.oz. | 0.038–0.044 | 14 | Tombstone is a RESTRICTED USE pesticide. |
| deltamethrin DELTA GOLD 1.5 EC Other trade names ¹ | 85–128 | 1.0–1.5 fl.oz. | 0.012–0.018 | 14 | Delta Gold is a RESTRICTED USE pesticide. |
| dimethoate DIMETHOATE 4E Other trade names ¹ | 8 | 16 fl.oz. | 0.5 lb. | 28 | |
| gamma-cyhalothrin DECLARE Other trade names ¹ | 83–125 | 1.02–1.54 fl.oz. | 0.01–0.015 | 30 | Declare is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin WARRIOR II with Zeon Technology Other trade names ¹ | 67–100 | 1.28–1.92 fl.oz. | 0.02–0.03 | 30 | Warrior II is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin + chlorantraniliprole BESIEGE | 13–21 | 6–10 fl.oz. | 0.02–0.03 + 0.04–0.06 | 30 | Besiege is a RESTRICTED USE pesticide. |
| spinosad + gamma-cyhalothrin CONSERO | 42–64 | 1–1.5 fl.oz. of each product | 0.03–0.046 + 0.01–0.015 | 30 | Consero is a RESTRICTED USE pesticide. It is a co-pack of two insecticides that must be applied together. |

¹ See Table 4 for other trade names.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control (cont.)

| Insecticide and Formulation | Acres Per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|---|------------------|--------------------------------|--------------------------------|--|---|
| GRASSHOPPERS (cont.) | | | | | |
| zeta-cypermethrin MUSTANG MAXX EC INSECTICIDE Other trade names ¹ | 32–46 | 2.8–4 fl.oz. | 0.017–0.025 | 14 (grain, stover) 45 (forage) | Mustang Maxx is a RESTRICTED USE pesticide. |
| GREEN JUNE BEETLE GRUBS (IN FIELDS WHERE BROILER LITTER HAS BEEN USED) | | | | | |
| carbaryl SEVIN XLR PLUS Other trade names ¹ | — | 1–1.5 qt. | 1–1.5 | 21 (grain, fodder) 14 (grazing, silage) | Treat on the surface of the soil when there is more than one grub per square foot in the fall prior to planting. Spring treatment when soil temperature is cold is not as effective as an application made in the fall. |
| SORGHUM MIDGE | | | | | |
| alpha-cypermethrin FASTAC EC Other trade names ¹ | 34–98 | 1.3–3.8 fl.oz. | 0.008–0.025 | 14 (grain, stover) 45 (forage) | Fastac EC is a RESTRICTED USE pesticide. |
| beta-cyfluthrin BAYTHROID XL | 98–128 | 1.0–1.3 fl.oz. | 0.008–0.01 | 14 | Baythroid XL is a RESTRICTED USE pesticide. |
| chlorpyrifos LORSBAN ADVANCED LORSBAN 75 WG Other trade names ¹ | 16 3 A/lb. | 0.5 pt. 0.33 lb. | 0.23 0.23 | 30 (1 pt. or less) 60 (> 1 pt.) | See label for detailed instructions. Lorsban Advanced is a RESTRICTED USE pesticide. Lorsban 75WG is not. |
| chlorpyrifos + lambda-cyhalothrin COBALT ADVANCED | 10–21 | 6–13 fl.oz. | 0.12–0.25 + 0.006–0.013 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt Advanced is a RESTRICTED USE pesticide. |
| chlorpyrifos + gamma-cyhalothrin COBALT Other trade names ¹ | 10–18 | 7–13 fl.oz. | 0.14–0.25 + 0.002–0.004 | 30 (26 fl.oz. or less) 60 (> 26 fl.oz.) | Cobalt is a RESTRICTED USE pesticide. |
| chlorpyrifos + zeta-cypermethrin STALLION | 11–34 | 3.75–11.75 fl.oz. | 0.08–0.25 + 0.008–0.025 | 30 (grain, stover) 45 (forage) | Stallion is a RESTRICTED USE pesticide. |
| cyfluthrin TOMBSTONE Other trade names ¹ | 98–128 | 1.0–1.3 fl.oz. | 0.016–0.020 | 14 | Tombstone is a RESTRICTED USE pesticide. |
| deltamethrin DELTA GOLD 1.5 EC | 67–98 | 1.3–1.9 fl.oz. | 0.015–0.022 | 14 | Delta Gold is a RESTRICTED USE pesticide. |
| dimethoate DIMETHOATE 4E Other trade names ¹ | 16–32 | 0.25–0.5 pt. | 0.125–0.25 | 28 | |
| esfenvalerate ASANA XL Other trade names ¹ | 22–44 | 2.9–5.8 fl.oz. | 0.015–0.03 | 21 | Asana is a RESTRICTED USE pesticide. |
| gamma-cyhalothrin DECLARE Other trade names ¹ | 125–166 | 0.77–1.02 fl.oz. | 0.0075–0.01 | 30 | Declare is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin WARRIOR II with Zeon Technology Other trade names ¹ | 100–133 | 0.96–1.28 fl.oz. | 0.015–0.02 | 30 | Warrior II is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin + chlorantraniliprole BESIEGE | 21–26 | 5–6 fl.oz. | 0.016–0.02 + 0.03–0.04 | 30 | Besiege is a RESTRICTED USE pesticide. |

¹ See Table 4 for other trade names.

Table 3. Suggestions for Postemergence Grain Sorghum Insect Control (cont.)

| Insecticide and Formulation | Acres Per Gallon | Amount of Formulation per Acre | Lb. Active Ingredient Per Acre | Minimum Days from Last Application to Harvest or Grazing | Comments |
|---|-------------------|--------------------------------|--------------------------------|--|--|
| SORGHUM MIDGE (cont.) | | | | | |
| methomyl LANNATE LV Other trade names ¹ | 5–11 | 0.75–1.5 pt. | 0.22–0.45 | 14 | Lannate is a RESTRICTED USE pesticide. |
| novaluron DIAMOND 0.83EC Other trade names ¹ | 11–14 | 9–12 fl.oz. | 0.12–0.16 | 14 (grain or stover) 7 (forage) | |
| spinosad BLACKHAWK Other trade names ¹ | 4.8–10.6 A/lb. | 1.5–3.3 oz. | 0.034–0.074 | 21 (grain) 3 (forage, fodder, hay) | |
| spinosad + gamma-cyhalothrin CONSERO | 42–64 | 1–1.5 fl.oz. of each product | 0.03–0.046 + 0.01–0.015 | 30 | Consero is a RESTRICTED USE pesticide. It is a co-pack of two insecticides that must be applied together. |
| zeta-cypermethrin MUSTANG MAXX EC INSECTICIDE Other trade names ¹ | 32–100 | 1.28–4 fl.oz. | 0.008–0.025 | 14 (grain, stover) 45 (forage) | Mustang Maxx is a RESTRICTED USE pesticide. |
| MITES | | | | | |
| dimethoate DIMETHOATE 4E Other trade names ¹ | 8 | 1 pt. | 0.5 | 28 | |
| propargite COMITE | 3.5–5.3 | 24–36 fl.oz. | 1.1–1.7 | 60 (grain) 30 (silage) | Comite is a RESTRICTED USE pesticide. |
| STINK BUGS, LEAF FOOTED BUGS, FALSE CHINCH BUGS ON GRAIN HEADS | | | | | |
| alpha-cypermethrin FASTAC EC Other trade names ¹ | 34–71 | 1.8–3.8 fl.oz. | 0.012–0.025 | 14 (grain, stover) 45 (forage) | Fastac EC is a RESTRICTED USE pesticide. |
| beta-cyfluthrin BAYTHROID XL | 46–98 | 1.3–2.8 fl.oz. | 0.010–0.022 | 14 | Baythroid XL is a RESTRICTED USE pesticide. |
| cyfluthrin TOMBSTONE Other trade names ¹ | 46–98 | 1.3–2.8 fl.oz. | 0.020–0.044 | 14 | Tombstone is a RESTRICTED USE pesticide. |
| deltamethrin DELTA GOLD 1.5 EC | 67–85 | 1.5–1.8 fl.oz. | 0.018–0.022 | 14 | Delta Gold is a RESTRICTED USE pesticide. |
| gamma-cyhalothrin DECLARE Other trade names ¹ | 83–125 | 1.02–1.54 fl.oz. | 0.01–0.015 | 30 | Declare is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin WARRIOR II with Zeon Technology Other trade names ¹ | 67–100 | 1.28–1.92 fl.oz. | 0.02–0.03 | 30 | Warrior II is a RESTRICTED USE pesticide. |
| lambda-cyhalothrin + chlorantraniliprole BESIEGE | 13–21 | 6–10 fl.oz. | 0.02–0.03 + 0.04–0.06 | 30 | Besiege is a RESTRICTED USE pesticide. |
| spinosad + gamma-cyhalothrin CONSERO | 42–64 | 1–1.5 fl.oz. of each product | 0.03–0.46 + 0.01–0.015 | 30 | Consero is a RESTRICTED USE pesticide. It is a co-pack of two insecticides that must be applied together. |
| zeta-cypermethrin MUSTANG MAXX EC INSECTICIDE Other trade names ¹ | 32–73 | 1.76–4 fl.oz. | 0.011–0.025 | 14 (grain, stover) 45 (forage) | Mustang Maxx is a RESTRICTED USE pesticide. |

¹ See Table 4 for other trade names.

Table 4. Insecticides Labeled for Use on Sorghum, Including Worker Protection and Harvest or Grazing Intervals

| Insecticide and Trade Name | A.I./Formulated Product | Formulation | Restricted Entry Interval (hr) | Minimum Days from Last Application to Harvest or Grazing | Bee Hazard Restriction |
|---|--------------------------|--|--------------------------------|--|--------------------------------------|
| alpha-cypermethrin MoA Group 3A* (grain sorghum + millet) | | | | | |
| FASTAC CS (Restricted Use) | 0.83 lb./gal. | capsule suspension | 12 | 14 (grain, stover) 45 (forage) | yes ¹ |
| FASTAC EC (Restricted Use) | 0.83 lb./gal. | emulsifiable concentrate | 12 | 14 (grain, stover) 45 (forage) | yes ¹ |
| beta-cyfluthrin MoA Group 3A (sorghum) | | | | | |
| BAYTHROID XL (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 12 | 14 | yes ³ |
| carbaryl MoA Group 1A (grain sorghum) | | | | | |
| SEVIN 4F | 4 lb./gal. | liquid suspension | 12 | 21 (grain, fodder) 14 (grazing, silage) | yes ² |
| SEVIN XLR PLUS CARBARYL 4L, others | 4 lb./gal. 4 lb./gal. | liquid suspension liquid suspension | 12 12 | Same as above Same as above | yes ² yes ² |
| chlorantraniliprole MoA Group 28 (grain sorghum, forage sorghum, sudangrass) | | | | | |
| DUPONT CORAGEN | 1.67 lb./gal. | suspension concentrate | 4 | 1 | no |
| DUPONT PREVATHON | 0.43 lb./gal. | suspension concentrate | 4 | 1 | no |
| chlorpyrifos MoA Group 1B (grain sorghum) Do not treat sweet varieties of sorghum. | | | | | |
| LORSBAN 4E (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | 30 (1 pt. or less) 60 (> 1 pt./A) | yes ¹ |
| LORSBAN ADVANCED (Restricted Use) | 3.755 lb./gal. | water emulsion | 24 | Same as above | yes ¹ |
| GOVERN 4E (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| HATCHET (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| CHLORPYRIFOS 4E AG, others (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| NUFOS 4E (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| YUMA 4E (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| WARHAWK (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| WHIRLWIND (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| LORSBAN 75WG | 12 oz./lb. | water dispersable granule | 24 | 30 (0.67 lb./A or less) 60 (> 0.67 lb./A) | yes ¹ |
| LORSBAN 15G | 2.4 oz./lb. | granular | 24 | not specified | yes ¹ |
| LORSBAN 15G SMARTBOX | 2.4 oz./lb. | granular | 24 | not specified | yes ¹ |
| SAURUS 15G | 2.4 oz./lb. | granular | 24 | not specified | yes ¹ |
| CHLORPYRIFOS 15G | 2.4 oz./lb. | granular | 24 | not specified | yes ¹ |
| CPF 4E (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | 30 (1 pt. or less) 60 (>1 pt./A) | yes ¹ |

¹ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

² BEE CAUTION: May kill honey bees and other bees in substantial numbers. Do not apply this product to target crops or weeds in bloom. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds in bloom. This product may show residual toxicity to honeybees, especially in humid climates and under slow drying conditions. Notifying beekeepers within 1 mile of treatment area at least 48 hours before product is applied will allow them to take additional steps to protect their bees. Limiting applications to times when bees are least active, e.g., within 2 hr of sunrise or sunset, will minimize risk to bees.

³ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

* MoA = Mode of Action classification from the Insecticide Resistance Action Committee (www.irac-online.org). Insecticides with different MoAs should be used for insecticide resistance management.

Table 4. Insecticides Labeled for Use on Sorghum, Including Worker Protection and Harvest or Grazing Intervals

| Insecticide and Trade Name | A.I./Formulated Product | Formulation | Restricted Entry Interval (hr) | Minimum Days from Last Application to Harvest or Grazing | Bee Hazard Restriction |
|---|-------------------------|--------------------------|--------------------------------|--|------------------------|
| chlorpyrifos MoA Group 1B (grain sorghum) Do not treat sweet varieties of sorghum.(cont.) | | | | | |
| ERASER (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| WARHAWK CLEARFORM (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| VULCAN (Restricted Use) | 4 lb./gal. | emulsifiable concentrate | 24 | Same as above | yes ¹ |
| chlorpyrifos + gamma-cyhalothrin MoA Group 1B + MoA Group 3A (grain sorghum) Do not use on sweet sorghum. | | | | | |
| BOLTON (Restricted Use) | 2.5 + 0.083 lb./gal. | emulsifiable concentrate | 24 | 30 (26 fl.oz./A or less) 60 (> 26 fl.oz./A) | yes ¹ |
| COBALT (Restricted Use) | 2.5 + 0.045 lb./gal. | emulsifiable concentrate | 24 | 30 (26 fl.oz./A or less) 60 (> 26 fl.oz./A) | yes ¹ |
| chlorpyrifos + lambda-cyhalothrin MoA Group 1B + MoA Group 3A (grain sorghum) Do not use on sweet sorghum. | | | | | |
| COBALT ADVANCED (Restricted Use) | 2.5 + 0.13 lb./gal. | emulsifiable concentrate | 24 | 30 (26 fl.oz./A or less) 60 (> 26 fl.oz./A) | yes ¹ |
| chlorpyrifos + zeta-cypermethrin MoA Group 1B + MoA Group 3A (grain sorghum) Do not use on sweet sorghum. | | | | | |
| STALLION BRAND (Restricted Use) | 2.75 + 0.275 lb./gal. | emulsifiable concentrate | 24 | 30 (grain, stover) 45 (forage) | yes ¹ |
| clothianidin MoA Group 4A | | | | | |
| PONCHO 600 | 5 lb./gal. | seed treatment | Not specified | Not specified | minimize planter dust |
| ACCELERON IC-609 | 5 lb./gal. | seed treatment | Not specified | Not specified | |
| NIPSIT INSIDE | 5 lb./gal. | seed treatment | 12 | Not specified | yes ⁴ |
| clothianidin + <i>Bacillus firmus</i> MoA Group 4A | | | | | |
| PONCHO VOTiVO | 4.17 + 0.84 lb./gal. | seed treatment | Not specified | Not specified | yes ⁴ |
| cyfluthrin MoA Group 3A | | | | | |
| TOMBSTONE (Restricted Use) | 2 lb./gal. | emulsifiable concentrate | 12 | 14 | yes ¹ |
| TOMBSTONE HELIOS (Restricted Use) | 2 lb./gal. | emulsifiable concentrate | 12 | 14 | yes ¹ |
| deltamethrin MoA Group 3A | | | | | |
| DELTA GOLD 1.5 EC (Restricted Use) | 1.5 lb./gal. | emulsifiable concentrate | 12 | 14 | yes ¹ |
| dimethoate MoA Group 1B (grain sorghum) | | | | | |
| DIMETHOATE 4E | 4 lb./gal. | emulsifiable concentrate | 48 | 28 | yes ¹ |
| DIMATE 4E | 4 lb./gal. | emulsifiable concentrate | 48 | 28 | yes ¹ |
| DIMETHOATE 400, others | 4 lb./gal. | emulsifiable concentrate | 48 | 28 | yes ¹ |
| DIMETHOATE 4EC | 4 lb./gal. | emulsifiable concentrate | 48 | 28 | yes ¹ |
| DIMETHOATE LV-4 | 4 lb./gal. | emulsifiable concentrate | 48 | 28 | yes ¹ |
| esfenvalerate MoA Group 3A (grain sorghum) | | | | | |
| ASANA XL (Restricted Use) | 0.66 lb./gal. | emulsifiable concentrate | 12 | 21 | yes ¹ |
| S-FENVALOSTAR (Restricted Use) | 0.66 lb./gal. | emulsifiable concentrate | 12 | 21 | yes ¹ |
| ZYRATE (Restricted Use) | 0.66 lb./gal. | emulsifiable concentrate | 12 | 21 | yes ¹ |
| flupyradifurone MoA Group 4D (sorghum) | | | | | |
| SIVANTO PRIME | 1.67 lb./gal. | soluble liquid | 4 | 7 (forage, hay) 21 (grain, stover) | yes ⁵ |

¹ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

⁴ This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

⁵ Toxic to adult bees in laboratory studies via oral exposure, however, not toxic to bees through contact exposure. Field studies conducted with this product have shown no effects on honeybee colony development.

Table 4. Insecticides Labeled for Use on Sorghum, Including Worker Protection and Harvest or Grazing Intervals

| Insecticide and Trade Name | A.I./Formulated Product | Formulation | Restricted Entry Interval (hr) | Minimum Days from Last Application to Harvest or Grazing | Bee Hazard Restriction |
|---|--------------------------------|-------------------------------|--------------------------------|--|------------------------|
| gamma-cyhalothrin MoA Group 3A (grain sorghum) | | | | | |
| DECLARE (Restricted Use) | 1.25 lb./gal | microencapsulated suspension | 24 | 30 | yes ¹ |
| PROAXIS (Restricted Use) | 0.5 lb./gal. | microencapsulated suspension | 24 | 30 | yes ¹ |
| imidacloprid MoA Group 4A (sorghum, millet) | | | | | |
| NITRO SHIELD | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ⁶ |
| MACHO 600 ST | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ¹ |
| GAUCHO 600 | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | no |
| AXCESS | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | no |
| NITRO SHIELD IV | 4 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ⁶ |
| SHARDA IMIDACLOPRID 5SC | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ⁴ |
| DYNA-SHIELD IMIDACLOPRID 5 | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ⁴ |
| RESONATE 600 ST | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ⁶ |
| SENATOR 600 FS | 5 lb./gal. | liquid used as seed treatment | 12 | 45 | no |
| ATTENDANT 480 FS | 4 lb./gal. | liquid used as seed treatment | 12 | 45 | yes ⁴ |
| imidacloprid + metalaxyl MoA Group 4A | | | | | |
| CONCUR SEED TREATMENT | 4 oz./lb. + 0.16 oz./lb. | dust used as seed treatment | 24 | Not specified | no |
| imidacloprid + carboxin and metalaxyl MoA Group 4A | | | | | |
| LATITUDE | 4 oz. + 2.2 oz. + 0.16 oz./lb. | dust used as seed treatment | 24 | 45 | no |
| lambda-cyhalothrin MoA Group 3A (grain sorghum) | | | | | |
| WARRIOR II with ZEON TECHNOLOGY (Restricted Use) | 2.08 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| KENDO (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| LAMCAP | 1 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| WILLOWWOOD LAMBDA-CY 1EC (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| LAMDA CY AG (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| LAMDA-CY EC (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| LAMBDASTAR (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| LAMBDASTAR 1CS (Restricted Use) | 1 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |

¹ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

⁴ This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

⁶ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift if bees are visiting the treatment area. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust, which can drift to blooming crops or weeds.

Table 4. Insecticides Labeled for Use on Sorghum, Including Worker Protection and Harvest or Grazing Intervals

| Insecticide and Trade Name | A.I./Formulated Product | Formulation | Restricted Entry Interval (hr) | Minimum Days from Last Application to Harvest or Grazing | Bee Hazard Restriction |
|---|-----------------------------|---|--------------------------------|---|------------------------|
| lambda-cyhalothrin MoA Group 3A (Grain sorghum) (cont.) | | | | | |
| LAMBDASTAR PLUS (Restricted Use) | 2 lb./gal. | aqueous-based formulation | 24 | 30 | yes ² |
| LAMBDA-T (Restricted Use) | 1 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| L-C INSECTICIDE | 1 lb./gal. | emulsifiable concentrate | 24 | 30 (grain) Do not graze or harvest for fodder, silage, or hay | yes ¹ |
| NUFARM LAMBDA CYHALOTHIN IEC (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| PARADIGM (Restricted Use) | 1 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| PROVINCE (Restricted Use) | 1 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| PROVINCE II (Restricted Use) | 2.08 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| RAVAGE (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 30 | 30 | yes ¹ |
| SILENCER (Restricted Use) | 1 lb./gal. | emulsifiable concentrate | 24 | 30 | yes ¹ |
| SILENCER VZN (Restricted Use) | 1 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| GRIZZLY TOO (Restricted Use) | 2.08 lb./gal. | capsule suspension | 24 | 30 | yes ¹ |
| lambda-cyhalothrin + chlorantraniliprole MoA Group 3A + MoA Group 28 (Grain Sorghum) | | | | | |
| BESIEGE (Restricted Use) | 0.417 lb. + 0.835 lb./ gal. | capsule suspension plus soluble concentrate | 24 | 30 | yes ¹ |
| malathion MoA Group 1B (Grain Sorghum) | | | | | |
| MALATHION 5, 5E, 57EC, others | 5 lb./gal. | emulsifiable concentrate | 12 | 7 | yes ¹ |
| GOWAN MALATHION, 8 others | 8 lb./gal | emulsifiable concentrate | 12 | 7 | yes ¹ |
| CHEMINOVA MALATHION 57% | 5 lb./gal. | emulsifiable concentrate | 12 | 7 | yes ¹ |
| FYFANON | 5 lb./gal. | emulsifiable concentrate | 12 | 7 | yes ¹ |
| FYFANON ULV AG | 9.9 lb./gal. | emulsifiable concentrate | 12 | 7 | yes ¹ |
| FYFANON 8 LB EMULSION | 8 lb./gal. | emulsifiable concentrate | 12 | 7 | yes ¹ |
| methomyl MoA Group 1A (Sorghum including sudangrass, not for sweet sorghum) | | | | | |
| DUPONT LANNATE LV (Restricted Use) | 2.4 lb./gal. | water soluble liquid | 48 | 14 | yes ¹ |
| DUPONT LANNATE SP (Restricted Use) | 14.4 oz./lb. | water soluble packet | 48 | 14 | yes ¹ |
| NUDRIN LV (Restricted Use) | 2.4 lb./gal. | water soluble liquid | 48 | 14 | yes ¹ |
| NUDRIN SP (Restricted Use) | 14.4 oz./lb. | water soluble packet | 48 | 14 | yes ¹ |
| ANNIHILATE LV (Restricted Use) | 2.4 lb./gal. | water soluble liquid | 48 | 14 | yes ¹ |
| ANNIHILATE SP (Restricted Use) | 14.4 oz./lb. | water soluble bags | 48 | 14 | yes ¹ |
| CORRIDA 29 SL (Restricted Use) | 2.4 lb./gal. | water soluble liquid | 48 | 14 | yes ¹ |
| CORRIDA 90 WSP (Restricted Use) | 14.4 oz./lb. | water soluble bags | 48 | 14 | yes ¹ |

¹ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.² In order to minimize the possibility of developmental effects on pollinator larvae, including honey bee brood, do not use Diamond 0.83 EC on blooming crops.

Table 4. Insecticides Labeled for Use on Sorghum, Including Worker Protection and Harvest or Grazing Intervals

| Insecticide and Trade Name | A.I./Formulated Product | Formulation | Restricted Entry Interval (hr) | Minimum Days from Last Application to Harvest or Grazing | Bee Hazard Restriction |
|--|-------------------------------|--|--------------------------------|---|------------------------|
| methoxyfenozide (Grain and Sweet Sorghum) MoA Group 18 | | | | | |
| INTREPID | 2 lb./gal. | flowable liquid | 4 | 2 (grain or stover) 3 (forage or sweet sorghum stalk) | no |
| TROUBADOUR | 2 lb./gal. | flowable liquid | 4 | 21 (grain or stover) 3 (forage or sweet sorghum stalk) | |
| novaluron MoA Group 15 (Grain Sorghum) | | | | | |
| DIAMOND 0.83 EC, others | 0.83 lb./gal. | emulsifiable concentrate | 12 | 7 (forage) 14 (grain and stover) | yes ⁷ |
| phorate MoA Group 1B (Grain Sorghum) | | | | | |
| THIMET 20-G SMARTBOX OR LOCK'N'LOAD OR EZLOAD (Restricted Use) | 3.2 oz./lb. | granular | 48 | 30 | no |
| propargite MoA Group 12C | | | | | |
| COMITE II (Restricted Use) | 6 lb./gal. | emulsifiable concentrate | 48 | 60 (grain) 30 (silage) | no |
| spinosad MoA Group 5 (sorghum, millet) | | | | | |
| BLACKHAWK | 5.8 oz./lb. | wettable powder | 4 | 21 (grain, straw) 3 (forage, fodder, hay) | yes ⁸ |
| ENTRUST | 12.8 oz./lb. | wettable powder | 4 | Same as above | yes ⁸ |
| ENTRUST SC | 2 lb./gal. | soluble concentrate | 4 | Same as above | yes ⁸ |
| SPINTOR 2 SC | 2 lb./gal. | soluble concentrate | 4 | Same as above | yes ⁸ |
| spinosad + gamma-cyhalothrin MoA Group 5 + MoA Group 3A (sorghum) | | | | | |
| CONSERO (Restricted Use) | 4 lb./gal. + 1.25 lb./gal. | co-pack containing 0.5 gallon of each insecticide | 24 | 30 | yes ¹ |
| terbufos MoA Group 1B (Grain sorghum) | | | | | |
| COUNTER 20G LOCK'N'LOAD OR SMARTBOX (Restricted Use) | 3.2 oz./lb. | granular | 48 | 50 (forage) 100 (grain or fodder) | no |
| thiamethoxam MoA Group 4A (sorghum) | | | | | |
| CRUISER 5FS | 5 lb./gal. | seed treatment | 12 | Not specified | yes ⁹ |
| zeta-cypermethrin MoA Group 3A (Grain sorghum) | | | | | |
| MUSTANG MAXX EC INSECTICIDE (Restricted Use) | 0.8 lb./gal. | emulsifiable concentrate | 12 | 14 (grain, stover) 45 (forage), silage | yes ¹ |
| RESPECT INSECTICIDE OR RESPECT EC INSECTICIDE (Restricted Use) | 0.8 lb./gal. | emulsifiable concentrate | 12 | 14 (grain, stover) 45 (forage) | yes ¹ |

¹ This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

⁷ Because of its mode of action as an insect growth regulator and since it is not systemic, Diamond 0.83EC has no direct effect on fully developed adult stages, such as bees and other beneficial pollinators. However, in order to minimize the possibility of transient effects on honeybee brood development, do not use Diamond 0.83EC on blooming crops when bees are actively foraging.

⁸ This product is toxic to bees exposed to treatment during the 3 hours following treatment. Do not apply to blooming, pollen-shedding, or nectar-producing parts of plants if bees may forage on the plants during this time period.

⁹ Thiamethoxam is highly toxic to bees exposed to direct treatment and effects may be possible as a result of exposure to translocated residues in blooming crops.

Other products may be available. Always read the label to make sure the specific crop is listed and to determine what rate to use. Insect Pest Management section prepared by Katelyn Kesheimer, Extension Specialist, Assistant Professor, Department of Entomology and Plant Pathology, Auburn University.

DISEASE CONTROL (2023 UPDATES)

Diseases may have a significant impact on the yield and grain quality of sorghum. While the risk of yield loss is greatest with the leaf blight and stalk (peduncle) rot phases of anthracnose, other diseases that can impact sorghum yield include Fusarium stalk rot and gray leaf spot. Forage sorghum varieties are particularly susceptible to anthracnose. Anthracnose, rough leaf spot, and zonate leaf spot are common and sometimes damaging diseases on sweet sorghum. Other than frequent showers, factors that influence crop susceptibility to diseases include sorghum cropping frequency, tillage practices, planting dates, and variety selection. During periods of extended drought stress, charcoal rot may reduce the yield and increase lodging in grain sorghum.

Descriptions of the diseases commonly found on grain sorghum in Alabama are listed below:

Anthracnose can be a serious problem where grain sorghum has been grown continuously and attacks both the foliage and stalks. Initially, red to maroon colored lesions with tan centers form along the mid rib of lower leaves. Lesions will increase in size and number giving affected leaves a severely blighted appearance. Rapid blighting can occur during wet, humid weather. Anthracnose can also cause a stalk rot that can be identified by the appearance of red to purple tissue on the stalk's surface and in its pith just below the seed head. Anthracnose develops most rapidly after flowering as the grain matures.

Charcoal rot is most common during periods of severe drought. It is caused by soil-borne fungus that infects plants at the soil line as the grain begins to mature. Stems of diseased plants will have a dry, stringy appearance near the soil line. Pith tissue will be stringy with a salt and pepper appearance. Diseased stalks may lodge.

Fusarium head mold occurs following periods of humid, hot weather at or shortly after bloom. The fungus infects the

panicles and rachis of the seed head and then the seed stalk. Red to purple spots are found on the diseased panicles and rachis. The most obvious symptom of Fusarium head mold is the appearance of a white to pink cotton-like growth on infected seed heads. Tissues of the infected seed stalk are a solid dark red. Diseased stalks may lodge and swine are highly sensitive to toxins produced by the fungus.

Gray Leaf Spot is a late-season disease of grain sorghum. The disease is characterized by the appearance of small, red spots on the leaves. Spots will enlarge into dark red to purple rectangular areas bordered by leaf veins, which may merge to form elongated stripes. The spots produce a gray cast as the causal fungus sporulates during moist, humid weather.

Maize Dwarf Mosaic Virus (MDMV) is a disease of corn and grain sorghum spread by aphids. This disease forms a mosaic of fine, yellow-green and dark green streaks that are most apparent on the younger leaves in the whorl. When exposed to cool nighttime temperatures, red streaks may appear on leaves of some infected sorghum varieties. The disease is most common on early grain sorghum planted in johnsongrass infested fields.

Rough leaf spot is a common disease of grain sorghum in Alabama, but rarely causes serious problems. Rough leaf spot lesions are elliptical, reddish in color with well defined margins. Leaf spots are rough to the touch due to hard, raised, black fruiting bodies of the causal fungus. Spots will grow together and killing areas of the leaf.

Zonate leaf spot is a common disease of grain sorghum, but rarely has a significant affect on grain quality or yield. The disease occurs primarily on the lower leaves and leaf sheaths. Spots on the leaves are circular, reddish-purple bands separated by tan-colored areas that form a concentric or zonate pattern ranging up to several inches in diameter. This disease develops mainly during wet weather.

Table 1. Grain Sorghum Disease Control

| Fungicide and Formulation | Rate per 1,000 Row Ft. | Comments |
|---|------------------------|---|
| SEED ROT AND SEEDLING DISEASE | | |
| azoxystrobin AFRAME AZOXY 2SC AZOXYSTAR DEXTER SC QUADRIS FLOWABLE | 0.4–0.8 fl.oz. | At-plant in-furrow spray for control of rhizoctonia and pythium seedling blight. Apply in 3 to 7 gallons of water at planting and direct spray into the seed furrow before seed is covered. Use higher rate when conditions favor disease. May help suppress charcoal rot. |
| LEAF SPOTS AND BLIGHTS | | |
| azoxystrobin AFRAME AZOXY 2SC AZOXYSTAR DEXTER SC QUADRIS FLOWABLE | 6–15.5 fl.oz. | For control of anthracnose, charcoal rot, and gray leaf spot on forage and grain sorghum. Begin applications prior to disease development and repeat 14 to 21 days later as needed. Use higher rate when conditions are favorable for disease on susceptible host. DO NOT make more than two consecutive applications of Quadris Flowable or other Group 11 fungicide (QoI). See label for additional information concerning resistance management with QoI fungicides. See label for additional instructions. |
| azoxystrobin + propiconazole AFRAME PLUS AZOXY PROP XTRA QUILT XCEL | 10.5–14 fl.oz. | For control of anthracnose, ergot, gray leaf spot, northern corn leaf blight, ladder spot, and zonate leaf spot on sorghum. Apply at first sign of disease and repeat after 14 days as needed. DO NOT apply more than 56 fluid ounces per acre of Quilt Xcel. |

| LEAF SPOTS AND BLIGHTS (cont.) | | |
|--|-------------|---|
| fluxapyroxad + pyraclostrobin PRIAXOR XEMIUM | 4–8 fl.oz. | For control of anthracnose, gray and cercospora leaf spot, northern leaf blight, rust, southern leaf blight, and bipolaris leaf spot. Make an application prior to disease development. Limit one application per season. |
| flutriafol TOPGUARD | 7–14 fl.oz. | For control of anthracnose, gray leaf spot, leaf blight, rough leaf spot, zonate leaf spot. Apply when conditions favor disease development and again 14 days later as needed. DO NOT make more than two applications per season. |
| flutriafol + azoxystrobin TOPGUARD EQ | 5–7 fl.oz. | For control of anthracnose, gray leaf spot, leaf spot, latter leaf spot, rough spot, target spot, and zonate leaf spot. Make first application when conditions favor disease development and repeat as needed if conditions remain favorable for disease development. DO NOT make more than two applications per year. DO NOT apply within 30 days of harvest. |
| fluxapyroxad + propiconazole + pyraclostrobin NEXICOR XEMIUM | 7–13 fl.oz. | For control of anthracnose, northern leaf blight, rust, and southern leaf blight. For optimal disease control, apply prior to disease development. Limit one application per season. |
| mefentrifluconazole + pyraclostrobin VELTYMA | 7–10 fl.oz. | For control of anthracnose, gray leaf spot, northern leaf blight, rust, and southern leaf blight. Apply before disease development. Make no more than one application per year. |
| picoxydostrobin APPROACH | 6–12 fl.oz. | For control of anthracnose, gray leaf spot, and rust on sorghum and sorghum spp. Apply prior to disease development and continue at 7- to 14-day intervals when disease pressure is high. DO NOT apply after flowering. DO NOT make more than two sequential applications before switching to a fungicide with a different mode of action. |
| pyraclostrobin HEADLINE HEADLINE SC | 6–12 fl.oz. | For control of anthracnose and gray leaf spot on forage and grain sorghum. Apply no later than at 25% flowering. DO NOT make more than application of Headline per year. If additional fungicide applications are needed, then use a labeled non-Group 11 fungicide with a different mode of action. |

Disease Control section prepared by **Amanda Strayer-Scherer**, Extension Plant Pathologist, Assistant Professor, Entomology and Plant Pathology, Auburn University.

WEED CONTROL (2023 UPDATES)

Weeds compete with grain sorghum for light, water, and nutrients. If weeds are not properly controlled, they will reduce yields and quality of the crop and be problematic during harvest.

Grain Sorghum and Johnsongrass

Johnsongrass is in the same family as grain sorghum, which means it is closely related to grain sorghum, making it difficult, if not impossible, to control. Do not plant grain sorghum in fields that are heavily infested with johnsongrass. No herbicides are available that will selectively control johnsongrass in grain sorghum.

Atrazine and Water Quality

Atrazine is a Restricted Use Pesticide because of ground and surface water concerns. Atrazine can travel through the soil and can enter groundwater, which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable (i.e., well-drained). These restrictions apply to all formulations and all atrazine-containing products.

Mixing, Loading, and Application

Atrazine must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes, and reservoirs or any wells, including abandoned wells, drainage wells, and sink holes. Atrazine must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If atrazine is applied to highly erodible land, the 66-foot buffer or setback from runoff entry points must be planted to a crop or seeded with grass or other suitable crop. See atrazine label for additional restrictions.

Application Rates

** Highly erodible soils (as defined by NRCS)—using conservation tillage practices (at least 30% plant residue coverage at planting), apply a maximum of 4 pt./A (2 lbs a.i./A). If residue is less than 30%, apply a maximum of 3.2 pt./A (1.6 lb. a.i./A).

** Not highly erodible soils. With no limit on the amount of plant residue, apply a maximum of 4 pt./A (2 lb. a.i./A).

*MOA=Mechanism of action. Herbicides with different MOAs should be used in weed resistance management. See Table 4.

Table 2. Grain Sorghum Weed Control

| Hericide Trade Name | Hericide Common Name | REI/PHI (Hours or Days) | Formulation | Rate/Acre Broadcast Active Ingredient | Hericide Group | Time of Application | Weeds Controlled | Comments |
|--|------------------------------|-------------------------|--|---------------------------------------|----------------|---|--------------------------------------|--|
| BURNDOWN (NO-TILL/REDUCED TILL) | | | | | | | | |
| Clarity/Vision/Banvel/ Sterling Blue/Clash/ Detonate/Diablo/ Dicamba DMA salt/ Rifle/Strut | dicamba | 24 hr/30 d | 8 fl. oz. | 0.25 lb. | | 4 Apply at least 15 days before sorghum planting. | Annual broadleaf weeds. | See label for tank-mixing partners and adjuvant requirements. Do not apply in tank mixtures with Lorsban herbicide. It may be applied in split applications: burndown followed by postemergence or preharvest; or postemergence followed by preharvest; Do not exceed 8 fl. oz./A per application or a total of 16 fl. oz./A per season. |
| Gramoxone SL/others | paraquat | 24 hr/48 d | 2–4 pt. | 0.5–1.0 lb. | | 22 Apply before sorghum emerges | Annual broadleaf weeds and grasses. | Use a non-ionic surfactant and apply in 20–30 gallons of spray solution per acre. Use the higher rate for harder-to-kill weeds. Complete coverage is essential for good weed desiccation. See label for specific application instructions and tank-mixing partners. |
| Sharpen | saflefencil | 12 hr/70 d | 1–2 fl. oz. | 0.022–0.045 lb. | | 14 Apply for burndown, preplant, or PRE control before crop emerges. | Horseweed and other broadleaf weeds. | An MSO is a required adjuvant for use with this product. See label for specifics. Tank-mix with glyphosate or Gramoxone SL for best burndown results. |
| Various* | glyphosate | 4 hr/7 d | 32–64 fl. oz. (3 lb. ae) 22–43 fl. oz. (4.5 lb. ae) | 0.75–1.5 lb. ae | 9 | Apply before, at planting, or immediately after planting but before crop emerges. | Broadleaf and grass weeds. | Adjuvant recommendations vary by glyphosate brand. See label for tank mixing partners. Tank mixes must follow the most restrictive REI/PHIs. Use lower rate on smaller, easy-to-kill weeds, increase rate with larger weeds and perennials. |
| Verdict | saflefencil + dimethenamid-P | 12 hr/70 d | 5–10 fl. oz. | 0.218–0.436 lb. | 14 + 15 | Apply burndown, preplant, or PRE to grain sorghum that has been treated with an approved chloroacetamide seed safener such as Concep III. | Horseweed and other broadleaf weeds. | Tank-mix with glyphosate or Gramoxone SL for best burndown results. See label for additional tank-mix partners and adjuvant system required. Verdict rate depends on soil texture and organic matter content. |
| PREEMERGENCE | | | | | | | | |
| Atrazine | atrazine | 12 hr/45 d | 1.6 qt if 4L or 1.8 lb. if 90% formulation | 1.6 lb | 5 | Apply prior to weed emergence. | Annual grass and broadleaf weeds. | Do not use on sand, loamy sand, or sandy loam or any soil with less than 1% organic matter. Do not plant crops other than corn or sorghum until the year after treatment. If applied after June 10, do not rotate with crops other than corn or sorghum the next year or crop injury may occur. |

Table 2. Grain Sorghum Weed Control (cont.)

| Herbicide Trade Name | Herbicide Common Name | REI/PHI (Hours or Days) | Rate/Acre Broadcast Formulation | Active Ingredient | Herbicide Group | Time of Application | Weeds Controlled | Comments |
|---|--------------------------|-------------------------|---|--|-----------------|---|---|--|
| PREEMERGENCE (cont.) | | | | | | | | |
| Bicep II Magnum | atrazine + S-metolachlor | 24 hr/ N/A | Coarse soils: do not use Medium soils: 2.1–2.33 qt. Fine soils: 2.33–2.58 qt. | Coarse soils: do not use Medium soils: 0.045–0.106 lb. Fine soils: 0.106–0.177 lb. | 5 + 15 | Apply preplant, PPI, or PRE to grain sorghum. | Annual grasses and broadleaf weeds. | Do not use on medium soils (loam, silt loam, silt) with less than 1.0% organic matter. For preplant applications, apply within 14 days before planting. Do not use unless your seed has been treated with Concep or Screen seed safener. See label for tank-mixing partners. |
| Bullet, Lariat | alachlor + atrazine | 12 hr/N/A | Coarse soils: 2.5 qt. Medium soils: 2.75 qt. Fine soils: 3.0 qt. | Coarse soils: 2.5 lb. Medium soils: 2.75 lb. Fine soils: 3.0 lb. | 5 + 15 | Apply before crop emergence. | Annual grasses and broadleaf weeds. | See label for tank-mixing partners. |
| Callisto | mesotrione | 12 hr/60 d | 6.0–6.4 fl.oz. | 0.012–0.0125 lb. | 27 | Apply more than 7d before planting, but not more than 21, will reduce risk of injury. Can apply PRE or PPI. | Annual broadleaf weeds | Do not apply to coarse-textured soils. |
| Cinch/Brawl II/ Charger Max/ Dual II Magnum/Medal II/ | S-metolachlor | 24 hr/75 d | Coarse soils: 1.0–1.33 pt. Medium soils: 1.33–1.5 pt. Fine soils: 1.33–1.67 pt. | Coarse soils: 0.96–1.27 lb. Medium soils: 1.27–1.43 lb. Fine soils: 1.27–1.6 lb. | 15 | Apply at planting. | Annual grasses and broadleaf weeds. | Do not use unless your seed has been treated with Concep or Screen seed safener. Tank mixtures are permitted only in states where the tank-mix partner is registered. Check label for tank-mix partners. |
| Cinch ATZ | atrazine + S-metolachlor | 24 hr/N/A | Coarse soils: do not use Medium soils: 3.2–4.2 pt. Fine soils: 3.2–4.2 pt. | Coarse soils: do not use Medium soils: 2.2–2.9 lb. Fine soils: 2.2–2.9 lb. | 5 + 15 | Apply before crop emergence. | Annual grasses and broadleaf weeds. | Do not use unless your seed has been treated with Concep or Screen seed safener. Tank mixtures are permitted only in states where the tank-mix partner is registered. Check label for tank-mix partners. |
| Degree Xtra | acetolachlor + atrazine | 12 hr/60 d | 2.0–2.9 qt. | 0.124 - 0.18 lb. | 15 + 5 | Preplant, PPI, and PRE | Annual grass and small-seeded broadleaf control | Do not use unless seed is treated with a safener for grain sorghum. |

Table 2. Grain Sorghum Weed Control (cont.)

| Herbicide Trade Name | Herbicide Common Name | REI/PHI (Hours or Days) | Rate/Acre Broadcast Formulation | Active Ingredient | Herbicide Group | Time of Application | Weeds Controlled | Comments |
|-----------------------------|---------------------------------------|-------------------------|---|--|-----------------|--|--|--|
| PREEMERGENCE (cont.) | | | | | | | | |
| Lexar, Lumex, Lumax EZ | S-metolachlor + atrazine + mesotrione | 24 hr/ | 3 qt | 2.78 lb. | 5 + 27 + 15 | Premetgerence up to 21 d before planting. | Grasses and broadleaf weeds. | This is a Restricted Use Pesticide. Do not apply on sandy soils. Sorghum seed must be treated with Concep III herbicide safener before planting. Add an NIS at 0.25% v/v or a COC at 1% v/v if weeds are present at time of application. |
| Outlook | dimethenamid-P | 12 hr/80 d | Coarse soils: 12–14 fl.oz. Medium- to fine-textured soils: 14–18 fl.oz. | 0.56–0.66 lb. 0.66–0.84 lb | 15 | Preplant, PPI, and pre-emergence | Grasses and small-seeded broadleaves | See label for approved tank mixes. May use a split application. |
| Sharpen | safufenacil | 12 hr/70 d | 1–2 fl. oz. | 0.022– 0.045 lb. | 14 | Apply PRE control before crop emerges. | Small- and large-seeded broadleaf weeds. | If weeds are present at the time of application, add glyphosate or gramoxone. An MSO is a required adjuvant for use with this product if weeds are present. See label for specifics. |
| Verdict | safufenacil + dimethenamid-P | 12 hr/70 d | 5–10 fl. oz. | 0.218–0.436 lb. | 14 + 15 | Apply PRE control before crop emerges. | Broadleaf and grass weeds. | If weeds are present at the time of application, add glyphosate or gramoxone. An MSO is a required adjuvant for use with this product if weeds are present. See label for specifics. Seed must have been treated with an approved chloroacetamide seed safener such as Concep III. |
| Warrant | Acetochlor | 12 hr/60 d | 3–6 pt. | 1.13–2.25 lb. | 15 | Apply preplant incorporated or preemergence prior to weed emergence. | Annual grasses and small-seeded broadleaf weeds. | Sorghum seed must be treated with an approved chloroacetamide herbicide safener. Do not exceed 8 pints per acre per year when making multiple applications. |
| POSTEMERGENCE | | | | | | | | |
| Aim EC, Aim EW | carfentrazone-ethyl | 12 hr/21 d | 0.5 fl. oz. (Aim EC) 0.5–0.8 fl. oz. (Aim EW) | 0.008 lb. (Aim EC) 0.008–0.013 lb. (Aim EW) | 14 | Apply from 4 inches tall to just prior to boot stage. | Broadleaf weeds. | Always add a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Do not use crop oil concentrates or methylated seed oils. See label for tank-mixing partners. |

Table 2. Grain Sorghum Weed Control (cont.)

| Herbicide Trade Name | Herbicide Common Name | REI/PHI (Hours or Days) | Rate/Acre Broadcast Formulation | Active Ingredient | Herbicide Group | Time of Application | Weeds Controlled | Comments |
|--|-------------------------|---|---------------------------------|-------------------|---|---|---|---|
| POSTEMERGENCE (cont.) | | | | | | | | |
| Atrazine | atrazine | 12 hr/45 d 2.2 lb.–90% | 2.0 qt.–4L 2.0 lb. | | 5 | Apply after sorghum is completely emerged but before 12 inches tall and before weeds are more than 1.5 inches tall. | Most annual grasses and broadleaf weeds. | Do not apply to sand or loamy sand, or injury may occur. Do not exceed 2 pounds of active ingredient per acre per year. Do not use when sorghum is under stress or crop is wet and succulent from recent rainfall. See label for restrictions. Follow directions for crop rotation. |
| Banvel, Clarity, Sterling Blue, Vision | dicamba | 24 hr/30 d | 8 fl. oz. | 0.25 lb. | 4 | Apply over the top from emergence to 15 inches tall. Best performance will be when sorghum is in 3- to 5-leaf stage and weeds are smaller than 3 inches tall. | Most broadleaf weeds. | Use caution to prevent drift and injury to sensitive crops. Do not use adjuvants. Do not apply by air. Use drop nozzles on 8–15" grain sorghum to avoid injury from herbicide collecting in the whorl. |
| Basagran | bentazon | 48 hr/N/A 1.5–2 pt. (4 lb./gal.) 1.2–1.6 pt. (5 lb./gal.) | 0.75–1 lb. | 6 | Apply after sorghum has completely emerged but before boot stage. | Most broadleaf weeds less than 4 inches tall. | Refer to label for specific weed sizes. Rainfall within 4 hours after application may reduce effectiveness. See label or adjuvant requirements and tank-mixing partners. | |
| Buctril | bromoxynil | 24 hr/ 0.5–0.75 pt. | 0.25–0.38 lb. | 6 | Apply between the 3-leaf stage but prior to the preboot stage (growth stage 4). | Most broadleaf weeds in the 2- to 4-leaf stage. | Do not add spray additives or mix with fluid fertilizers. See label for tank-mixing partners. Applications prior to the 3-leaf growth stage may result in increased crop leaf burn. Less drift potential than dicamba or 2,4-D. | |
| Degree Xtra | acetolachlor + atrazine | 12 hr/60 d | 2.0–2.9 qt. | 0.124 - 0.18 lb. | 15 + 5 | Apply post to grain sorghum before the crop exceeds 11 inches in height (in general, 5–6 leaf stage). | Grasses and small-seeded broadleaves. | This is a Restricted Use Pesticide. See label for approved tank mixes. May use a split-application. |

Table 2. Grain Sorghum Weed Control (cont.)

| Herbicide Trade Name | Herbicide Common Name | REI/PHI (Hours or Days) | Rate/Acre Broadcast Formulation | Active Ingredient | Herbicide Group | Time of Application | Weeds Controlled | Comments |
|------------------------------|----------------------------|-------------------------|--|--|---|--|---|---|
| POSTEMERGENCE (cont.) | | | | | | | | |
| Facet L | quinchlorac | 12 hr/ 22-32 fl. oz. | 0.115 - 0.151 lb. 0.21 - 0.26 lb. 12.8 - 16 fl. oz. | 4 27 + 6 15 | Apply POST (plants up to 12 inches tall) Apply between 3-leaf up to 30 inches and/or prior to flag leaf emergence. Postemergence up to 12 inches in grain sorghum | Annual grass and broadleaf weeds Certain broadleaf weeds. | Annual grass and broadleaf weeds Grasses and small-seeded broadleaves. | For best annual grass control, apply with atrazine at 0.5-1.0 lb ai/A when weeds are less than 2 inches tall. Do not use liquid fertilizer as a carrier for POST application of Facet L to grain sorghum. Huskie is an EC so see label for tank mix partners. For optimal activity, add 1 pound of AMS per acre. |
| Huskie | pyrasulfotole + bromoxynil | 24 hr/60 d | 0.56-0.66 lb. 12-14 fl.oz. Medium- to fine-textured soils: 14-18 fl.oz. | 0.66-0.84 lb. | | | | This is a Restricted Use Pesticide. See label for approved tank mixes. May use a split-application. |
| Outlook | dimethenamid-P | 12 hr/80 d | | | | | | Prowl H2O will not control emerged weeds. Rainfall/irrigation of 0.5" to 1.0" can be used instead of mechanical cultivation. Herbicide must be applied to weed-free soil. |
| Prowl H2O | pendimethalin | 24 hr/60 d | Coarse soil: 1.5 pt Medium soil: 2.0 pt Fine soil: 3.0 pt | 0.5-0.74 lb. 0.74-1.0 lb. 0.74-1.5 lb. | 3 | Cultivate to throw soil around stems and protect brace roots when sorghum is a minimum of 4" in height and immediately spray with Prowl. Use drop nozzles if sorghum foliage will prevent uniform coverage of the soil surface. Can spray as late as the last cultivation. | Annual grasses and small-seeded broadleaf weeds. | |
| Sandeia, Permit, Profine 75 | halosulfuron | 12 hr/30 d | 0.67 - 1 oz. | 0.031 - 0.476 lb. | 2 | Apply at the 2-leaf through layby stage (before grain head emerges) | nutsedges, grasses, broadleaves | Do not make more than one application per use per season. See label for tank-mix partners. Use an NIS, COC in the spray mixture. |

Table 2. Grain Sorghum Weed Control (cont.)

| Herbicide Trade Name | Herbicide Common Name | REI/PHI (Hours or Days) | Rate/Acre Broadcast | | Herbicide Group | Time of Application | Weeds Controlled | Comments |
|------------------------------|-----------------------|-------------------------|---------------------|-------------------|-----------------|---|---|---|
| | | | Formulation | Active Ingredient | | | | |
| POSTEMERGENCE (cont.) | | | | | | | | |
| Warrant | acetochlor | 12 hr/60 d | 3–6 pt. | 1.13–2.25 lb. | 15 | Postemergence up to 11 inches tall and with up to 5–6 leaves. | Annual grasses and small-seeded broadleaf weeds. | Sorghum seed must be treated with an approved chloroacetamide herbicide safener. This treatment will not control emerged weeds. Do not exceed 8 pints per acre per year when making multiple applications. Use lower rates on coarse-textured soils and low organic matter. |
| Zest WDG | nicosulfuron | 4 hr/see label | 0.67-1.33 oz. | 0.031 - 0.062 lb. | 2 | Apply to emerged grain sorghum containing the INZEN technology that is up to 20 inches tall. Applications from fifth leaf stage. | crabgrass, broadleaf signalgrass, johnsongrass, panicums. | May only be used on INZEN grain sorghum. Applications made at approximately five leaf to flag leaf visible provides the best crop tolerance. Must use a COC at 1% v/v or NIS at 0.25% v/v. |
| 2,4-D (various trade names) | 2,4-D | 48 hr/30 d | 1.0 - 1.5 pt | 0.5 - 0.7 lb. | 4 | Apply over the top of grain sorghum that is 5-15 inches tall. Use drop nozzles if sorghum is more than 8 inches in height. Use a non-ionic surfactant at 0.25% v/v or a crop oil concentrate at 1% v/v. | Most annual broadleaf weeds. | Use caution to prevent drift and injury to sensitive crops. Do not apply during boot, bloom, or early dough stages or apply in fluid fertilizers. Do not use the ester formulations to prevent drift to susceptible crops. |

Table 2. Grain Sorghum Weed Control

| Herbicide Trade Name | Herbicide Common Name | REI/PHI (Hours or Days) | Rate/Acre Broadcast | | Herbicide Group | Time of Application | Weeds Controlled | Comments |
|----------------------------------|-----------------------|-------------------------|--|-------------------|-----------------|---|---|---|
| | | | Formulation | Active Ingredient | | | | |
| HARVEST AID | | | | | | | | |
| Aim | carfentrazone-ethyl | 12 hr/3 d | 1 fl. oz. | 0.016 lb. | | 14 | Apply when grain is fully mature, when black layer has formed, and when kernels are hard. | Desiccation of broadleaf weeds, such as pigweeds and morningglories |
| glyphosate | glyphosate | 4 hr/7 d | 32–64 oz. 24–48 oz. 19–38 oz. | 19–38 oz. | 9 | Apply when grain reaches 30% moisture or less and black layer has formed. | Desiccation of broadleaf weeds, grasses, and green vegetation. | Allow a minimum of 7 days before harvest. Do not use on sorghum grown for seed as a reduction in seed germination or vigor may occur. See label for use as all glyphosate formulations may be labeled for this use. |
| Sodium Chlorate, Defol 5, others | sodium chlorate | | 4.8 qt. (5 lb./gal.) 3.2 qt. (7.5 lb./gal.) | 5–7.5 lb. | | Apply when sorghum has reached 25% or less moisture. | Desiccation of grasses. | Make application on a sunny day when air temperature is above 85 degrees and humidity is below 65%. Apply 7 to 10 days before planned harvest date. |

Rate of herbicides are given for broadcast application. Band application reduces the amount needed per acre of crop and can be determined by the formula:

$$\frac{\text{Band Width}}{\text{Row Width}} \times \text{Broadcast Rate} = \text{Band Rate}$$

For example, the amount of AATrex 4L needed for a broadcast application to a light sandy soil is 2 quarts per acre. The amount needed to treat a 20-inch band on a 30-inch row would be $20/30 \times 2 \text{ qt./A} = 1.33 \text{ qt./A}$ on a 20-inch band

| Table 3. Weed Response to Grain Sorghum Herbicides | | | | | | | | |
|--|----------|--------------|------------|---------|---------|-----------------|----------------|---------|
| Herbicides | Clarity | Gramoxone SL | Glyphosate | Sharpen | Verdict | Bicep II Magnum | Dual II Magnum | Warrant |
| Application Timing | BURNDOWN | | | | | PRE | | |
| Site of Action Group | 4 | 22 | 9 | 14 | 14 + 15 | 5 + 15 | 15 | 15 |
| Weeds | | | | | | | | |
| PERENNIAL WEEDS | | | | | | | | |
| bermudagrass | N | P | G | N | N | N | N | N |
| johnsongrass, rhizome | N | P | E | N | N | N | N | N |
| nutsedge, purple | N | P | F | N | N | N | N | N |
| nutsedge, yellow | N | P | F | N | P | F-G | F-G | N |
| ANNUAL GRASSES | | | | | | | | |
| crabgrass, large | N | F | G | N | G | E | E | G |
| crowfootgrass | N | F | E | N | F | | E | G |
| goosegrass | N | F | E | N | F | E | E | G |
| fall panicum | N | F | G | N | F | E | E | G |
| foxtail | N | F | E | N | G | E | | G |
| johnsongrass (seedling) | N | F | E | N | G | G | G | F |
| signalgrass, broadleaf | N | F | G | N | G | G | G | G |
| Texas panicum | N | F | G | N | F | F-G | P-F | F |
| BROADLEAF WEEDS | | | | | | | | |
| beggarweed, Florida | G | E | E | | | | F | F |
| bristly starbur | E | E | E | | | | P | F |
| black nightshade | E | E | E | E | E | G | G | G |
| cocklebur | E | E | E | E | E | F | P | N |
| crotalaria, showy | G | E | E | E | E | P | P | - |
| croton, tropic | G | E | E | | | | P | - |
| groundcherries | E | E | E | E | E | G | G | F |
| jimsonweed | E | E | E | E | E | | P | - |
| lambsquarters, common | E | E | E | E | E | E | F | G |
| morningglories sp. | E | E | F | E | E | G | P | P |
| pigweed | G-E | E | E | E | E | E | G | F |
| prickly sida | E | E | E | E | E | F | F | F |
| pusley, Florida | E | E | E | | | | G-E | G |
| ragweed, common | E | E | E | E | E | E | F | - |
| ragweed, giant | E | E | E | G | G | F | P | - |
| sicklepod | E | E | E | P | P | F | P | - |
| smartweed | G | E | E | E | E | E | G | F |
| velvetleaf | G | E | E | E | E | F | P | - |

N = No control

P = < 70% control

F = 70–80% control

G = 80–90% control

E = 90% control

¹ Will not control ALS-resistant ryegrass² Weeds must not be larger than 2 inches at time of application.

Table 3. Weed Response to Grain Sorghum Herbicides (cont.)

| Herbicides | Cinch ATZ | Atrazine | Basagran | Buctril | 2,4-D | Warrant | Zest WDG |
|-------------------------|-----------|----------|----------|---------|-------|---------|----------|
| Application Timing | PRE | POST | | | | | |
| Site of Action Group | 5 + 15 | 5 | 6 | 6 | 4 | 15 | 2 |
| Weeds | | | | | | | |
| PERENNIAL WEEDS | | | | | | | |
| bermudagrass | N | N | N | N | N | N | G-E |
| johnsongrass, rhizome | N | N | N | N | N | N | G |
| nutsedge, purple | N | N | P | N | N | N | N |
| nutsedge, yellow | F-G | N | G | N | N | N | N |
| ANNUAL GRASSES | | | | | | | |
| crabgrass, large | E | F | N | N | N | G | E |
| crowfootgrass | | | N | N | N | G | F |
| goosegrass | E | P | N | N | N | G | F |
| fall panicum | E | P | N | N | N | G | E |
| foxtail | E | | N | N | N | G | E |
| johnsongrass (seedling) | G | P | N | N | N | F | E |
| signalgrass, broadleaf | G | F | P | N | N | G | E |
| Texas panicum | F-G | | N | N | N | F | E |
| BROADLEAF WEEDS | | | | | | | |
| beggarweed, Florida | | G | G | G | P | F | N |
| bristly starbur | | E | E | G | G | F | N |
| black nightshade | G | G | P | G | G | G | N |
| cocklebur | F | E | E | E | E | N | N |
| crotalaria, showy | P | G | P | | G | - | N |
| croton, tropic | | G | P | | G | - | N |
| groundcherries | G | F | P | G | G | F | N |
| jimsonweed | | E | E | | E | - | N |
| lambsquarters, common | E | E | P | G | G | G | N |
| morningglories sp. | G | E | F | F | E | P | N |
| pigweed | E | E | P | G | G | F | N |
| prickly sida | F | E | F-G | G | F | F | N |
| pusley, Florida | | G | P | G-E | G | G | N |
| ragweed, common | E | E | P | F | G | - | N |
| ragweed, giant | F | F | P | N | E | - | N |
| sicklepod | F | F-G | N | P | G | - | N |
| smartweed | E | G | F | G | P | F | N |
| velvetleaf | F | F | G | G | G | - | N |

N = No control

P = < 70% control

F = 70–80% control

G = 80–90% control

E = 90% control

¹ Will not control ALS-resistant ryegrass² Weeds must not be larger than 2 inches at time of application.

Table 3. Weed Response to Grain Sorghum Herbicides (cont.)

| | |
|----------------------|--------------------|
| Herbicides | Aim |
| Application Timing | HARVEST AID |
| Site of Action Group | 14 |

Weeds

| PERENNIAL WEEDS | |
|-------------------------|---|
| bermudagrass | N |
| johnsongrass, rhizome | N |
| nutsedge, purple | N |
| nutsedge, yellow | N |
| ANNUAL GRASSES | |
| crabgrass, large | N |
| crowfootgrass | N |
| goosegrass | N |
| fall panicum | N |
| foxtail | N |
| johnsongrass (seedling) | N |
| signalgrass, broadleaf | N |
| Texas panicum | N |
| BROADLEAF WEEDS | |
| beggarweed, Florida | |
| bristly starbur | |
| black nightshade | N |
| cocklebur | F |
| crotalaria, showy | |
| croton, tropic | |
| groundcherries | |
| jimsonweed | |
| lambsquarters, common | G |
| morningglories sp. | G |
| pigweed | G |
| prickly sida | P |
| pusley, Florida | |
| ragweed, common | |
| ragweed, giant | P |
| sicklepod | N |
| smartweed | F |
| velvetleaf | E |

N = No control

P = < 70% control

F = 70–80% control

G = 80–90% control

E = 90% control

¹ Will not control ALS-resistant ryegrass² Weeds must not be larger than 2 inches at time of application.

Weed Control section prepared by David Russell, Assistant Extension Professor, Assistant Research Professor, Crop, Soil, and Environmental Sciences, Auburn University.



FOR MORE INFORMATION on pesticides, pesticide safety, or submitting samples for analysis, see the following publications in the IPM series:

- IPM 1293, "Safety." Safety contact information; worker protection standards; the safe use, handling, and storage of pesticides
- IPM 1294, "Submitting Samples." Procedures for submitting samples for diagnosis, analysis, and identification
- IPM 1295, "General Pesticide Information." Federal and state restricted use pesticide lists; pesticides and water quality
- IPM 1317, "Appendix." Pesticide guidelines for agronomic crops, including preharvest intervals; rain-free requirements; grazing restrictions; crop rotation guidelines; and the names, classifications, and toxicities of pesticides.

2023 IPM-0429

Trade and brand names used in this publication are given for information purposes only. No guarantee, endorsement, or discrimination among comparable products is intended or implied by the Alabama Cooperative Extension System.

In accordance with Federal law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, this institution is prohibited from discriminating on the basis of race, color, national origin, sex (including gender identity and sexual orientation), age, disability, and reprisal or retaliation for prior civil rights activity.

Program information may be made available in languages other than English. Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, and American Sign Language) should contact the Alabama Cooperative Extension System Human Resources Department at (334) 844-5531 or the State of Alabama Governor's Office on Disability (GOOD) at (888) 879-3582 or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339.

To file a program discrimination complaint, a complainant should complete a Form AD-3027, USDA Program Discrimination Complaint Form, which can be obtained online at www.usda.gov/oascr/how-to-file-a-program-discrimination-complaint from any USDA office, by calling (866) 632-9992, or by writing a letter addressed to USDA. The letter must contain the complainant's name, address, telephone number, and a written description of the alleged discriminatory action in sufficient detail to inform the Assistant Secretary for Civil Rights (ASCR) about the nature and date of an alleged civil rights violation. The completed AD-3027 form or letter must be submitted to USDA by: U.S. Department of Agriculture | Office of the Assistant Secretary for Civil Rights | 1400 Independence Avenue, SW | Washington, D.C. 20250-9410 | Fax: (833) 256-1665 or (202) 690-7442; or Email: program.intake@usda.gov.

This institution is an equal opportunity provider.

© 2023 by the Alabama Cooperative Extension System. All rights reserved.

www.aces.edu